College research projects. I am in a well-organized, up-to-date group practice of four. We have a rota system. Despite these advantages we work a 74- to 80-hour week. In my opinion the publication of this report at this time was ill advised. To release it to the press before doctors generally had had time to assess it was, to say the least, misguided.

It should have been realized that the worst possible construction would be placed upon any adverse comments in it. A delay of one or two months, pending the completion of negotiations with the Minister of Health, would have been to the profession's advantage. Instead a disservice has now been rendered which is practically irreparable, conclusions having been drawn which in my opinion are inaccurate, hasty, and statistically invalid.

I agree that much of the general practitioner's time is taken up with trivia, but one of the qualities that makes a good general practitioner is his ability to sort out the serious from the trivial. Are we going to delegate this vital function of differential diagnosis to someone less qualified? But I agree that something will have to be done to aid in coping with this problem.

To analyse work load into actual time spent face to face with the patient is, of course, fallacious. How much time during an eight-hour day does an architect spend with his client, a colliery worker cut coal, or a solicitor spend in court? It is well known that physiologically a human being can work continuously and efficiently only for a period of a few hours without having a pause. Finally I would point out that the work load of general practitioners in South Wales is twice that of the south of England. How many practices in South Wales, the industrial north, or industrial Scotland took part in this survey? It is in these areas where the present problem is most acute.—I am, etc.,

Port Talbot.

D. HUBERT JONES.

SIR,—It is almost unbelievable that Dr. Fry and the College of General Practitioners should publish their report at the present stage.

In one blow they have, perhaps, succeeded in demolishing the general practitioners' case, which the Ministry have never succeeded in doing.

The folly of it is almost beyond belief, the motives impossible of deduction, the result probably catastrophic.—I am, etc.,

London S.W.4.

F. V. GRIFFITHS.

Certification

SIR,—I notice that the Blyth with Morpeth motion (*Supplement*, 24 July, p. 78), that all certification for N.H.I. purposes be abolished, was carried. I assume, therefore, that this is now the policy of the B.M.A., though I am very surprised at the lack of publicity over this quite radical departure.

I should like to suggest some ways of starting the ball rolling in this direction, and it is quite clear that our negotiators have no idea of the radical steps required, or they would not be so pleased with their reduction of four certificates a week.

Firstly, I think it is essential that the National Insurance Service should be completely separated from the National Health Service.

This I think is a fundamental error in the present National Health Service agreement. Separation of these two services would have a large and immediate beneficial effect on the Health Service. I do not mind patients coming in for trivial illnesses if they are worried about them, but I do object when they come in for trivial illnesses because all they want is a certificate. Furthermore, it would greatly improve the doctor-patient relationship, and it would be quite clear to the patient that the doctor's job was to give him advice or treatment on medical matters and not just to dish out certificates. It would also improve relationships, as there would be no need for arguments about whether the patient was or was not fit for work, which, though uncommon in my practice, do take up a long time when they occur.

This is a practical proposition. There is no reason why certificates should not be issued by medical auxiliaries such as health visitors, who, in cases of doubt, could call upon the R.M.O. I am sure a health visitor is just as well able to say whether somebody has diarrhoea or the "flu" from the certification angle as is the doctor. The treatment side, of course, would remain with the doctor. It is these short-period illnesses which tend to take up unnecessary time in certification. For illnesses over two weeks in duration, if the patient found his G.P.'s advice conflicted with the R.M.O. he could appeal for a general-practitioner certificate.

It would be up to the Minister to make alternative arrangements for his National Insurance Service, but I think we must stand out and make it quite clear that we consider it no part of the National Health Service.—I am, etc.,

Newport, Mon.

MICHAEL WADE.

Remuneration

SIR,—May I have this opportunity of correcting a misquotation of my speech on remuneration to the A.R.M? (*Supplement*, 17 July, p. 66.) I am reported as saying that the remuneration of a doctor with a list of 2,200 patients should be not less than £3,000 a year net. The figure I gave was £4,000 a year net. This is, of course, a very conservative revaluation in 1965 terms of the income suggested by the Spens Committee for the average doctor, which income was fully accepted by the Government as appropriate in 1948.—I am, etc.,

Low Fell,
Gateshead-on-Tyne.

J. C. ARTHUR.

Abuse of the Service

SIR,—Last Saturday at 6 p.m. a patient, who had visited me earlier in the week and who was not responding to treatment for a sore throat, asked me to call. She had not reattended, nor sent a message before 10 a.m., as she had hoped it would get better. I agreed to visit that evening and was asked when I would arrive. I wanted to know why she asked, and she replied that she was going out for dinner that evening.

This morning at 4 a.m. I was called to a woman with "chest pain." On arrival I found the patient had attended the surgery before with premenstrual breast pain, which had been investigated, her chest x-rayed, and she had been advised. This morning she had recognized the pain as the usual one and had not even tried an aspirin for relief.

This type of abuse cannot be stopped by "educating the public," and I urge our negotiators to press for a complaints machinery with penalties, which can be used against that small minority of patients who misuse the Health Service.—I am, etc.,

London N.W.10.

R. LAW.

"The Right Word"

SIR,—In answer to the letter by Dr. T. P. Lineham, Jun. (17 July, p. 174), which was also the opinion of "Pertinax," that the general practitioner's name gave the wrong image to the general public in this age of free-for-all under the National Health Service, I do not think that his word "physician" is well chosen. Granted the word "physician" denotes the knowledge of medicine and surgery (according to *Cassell's Dictionary*), but the name implies to most of us that the practitioner has chosen medicine as his "raison d'être." However, in *Nuttall's Standard English Dictionary* "physician" is defined as "a doctor of medicine; one whose profession is to prescribe remedies for diseases and not to perform surgical operations." Unfortunately for my interpretation of definition it continues "a licentiate as well as member or fellow of the Royal College of Physicians," and most general practitioners have the L.R.C.P. as we know. Gunnar Björck (2 July, p. 7) uses the word continuously to describe all medical practitioners, translating from his Swedish tongue.

Let us keep "physician" as the definitive name for a medical specialist. Certain epithets are not outworn. Surely the general practitioners should be proud of the surgical half of their degrees, which the physician prefers to discard?—I am, etc.,

London W.1.

MARJORIE M. DOBSON.

Scared to Death?

SIR,—I was very interested to read the letter under the above heading (7 August, p. 363).

It would appear that a not unlikely diagnosis here would be fulminating meningococcal septicaemia or Friderichsen-Waterhouse syndrome. Practitioners who were working in infectious diseases hospitals in Great Britain during the epidemic of cerebrospinal fever in 1940 would have seen a few cases of this.

I saw well over 300 cases of cerebrospinal fever at the Cardiff City Isolation Hospital (as it was then called) in that year, and they included five admissions of the fulminating variety of the disease, all of whom succumbed within a few hours of admission; if my memory serves me correctly they were adolescent or adult, one being an elderly man in the 60's.

The outstanding features were the state of shock with multiple petechiae and ecchymoses