Current Practice

CHILD CARE IN GENERAL PRACTICE

Emotional and Behavioural Disorders: Part II

JOHN APLEY,* M.D., F.R.C.P

Sleep Disorders

It is true that in the early weeks most babies hardly wake between feeds and that the amount of sleep required decreases as they get older. But even from the earliest days some infants seem to do with little sleep, and some children seem to sleep really soundly only when they are old enough to exhaust themselves by running about. Each child has his own pattern of sleep. Its quality and adequacy are best judged when he is awake, by how he is and what he does then.

When a child is asleep he may be restless, breathe heavily or noisily, grind his teeth, rock, bang his head, or perspire. None of these is important, and the parents will be happier if they understand and follow the doctor's advice and do not look for trouble.

The infant who does not readily fall asleep may be hungry or cold or he may be uncomfortable because of wet clothes, though many a baby sleeps perfectly well in a batter of excreta. A new or temporary disturbance of previously regular sleep may be due to pain or to a feverish illness.

If the infant unexpectedly wakes at night and cries the mother will naturally want to see him. But she should be advised that an immediate resort to feeding and picking up is better avoided. If he is cold he can be covered; if he is not hungry or in pain he can be soothed in his cot. The mother who then goes out and closes the door behind her and waits for 10 minutes (timed by the clock) will nearly always find that the baby has fallen asleep and she has avoided the formation of a troublesome habit.

The commonest cause for habitual screaming, either when the child is put to bed or on awaking in the night, is habit formation based on attention-seeking. An infant even only a few months old is quick to learn that crying will bring his mother, who may pick him up from his cot, make a fuss of him, and possibly take him into bed with her. The habit is easier to prevent than to break. The infant should not be picked up at the first cry, and if he needs soothing it can usually be done in the cot. The habit may need to be broken with the help of a sedative, such as chloral hydrate (1-2 gr., 60-120 mg.) given to the child for a few nights only. The mother, with whom the rationale has been discussed, may need a sedative too.

In the older child many of the sleep disturbances will have started with faulty habits acquired in infancy. If the child lies awake for hours after going to bed he may be insufficiently tired through lack of exercise, or uncomfortable through being tucked in too tight; he may have been over-stimulated by excitement caused by watching television before bedtime or reading horrific stories in bed. He may be provoked by the excessive noise of older members of the family obviously enjoying themselves downstairs while he is "punished" by being

banished to bed. He may be afraid of the dark or of the shadows cast by a flickering light.

Children learn best by example. In a family where going to bed is taken for granted and welcomed sleep problems are not common or serious. The parents who are obsessed by their child's need for "enough" sleep are those whose child will have sleep problems. They should not be blamed but helped. Going to bed is not to be used as a threat or a punishment. The child who lies awake for hours may sleep more readily if he goes to bed later rather than earlier. Undue excitement before bedtime is best avoided, and a simple routine of a meal, a little play, bath, and reading a familiar story is often effective. A "good-night cuddle" is fine, but it should be accepted also as final. It is unnecessary to go into the child's room at night to see if he is all right, or breathing, or even alive. The child who has a fear of the dark should, of course, be given a light or have immediate access to a light-switch. He may be happier in a room with a brother or sister. A favourite doll or a glass of water by the bedside, though it may never be drunk, may act as a soother. The older child should be capable of understanding that his parents need sleep too and that they should be called only for an important reason. When parents are still unduly worried the doctor can draw to their notice the child's good health and healthy reactions while he is awake.

Appetite Disorders

Here we do not discuss the child with a poor appetite, who looks poorly, and is ill with an acute or chronic infection, renal disease, or neoplasm and may also have lost weight. Habitual thinness is another problem. We need not hold with the political agitator who based his campaign on the fact that half the children in this country are below the average weight. Some children are always thin, even if they eat well. But a loss of weight associated with poor appetite is strongly suggestive of illness, which may be either physically or psychologically determined, for the emotions can influence the appetite as well as the digestion.

Food Refusal

The problem of the child who looks well, is active and growing, has no evidence of disease, but is brought for advice because "nothing passes his lips," or "he doesn't eat enough to keep a bee alive," is a common one. Faced with a pink-cheeked child, whose nutrition is obviously adequate, and an inevitably anxious mother the doctor needs diplomacy and patience. The remedy lies not in a blunt statement of fact, or in criticism of parental management, but in discussion and explanation.

Food refusal or faddiness in the child does, indeed, often reflect errors in management by the parents, but "realization

* Consultant Paediatrician, United Bristol Hospitals and Bath Clinical Area, Bristol Royal Hospital for Sick Children.

that the mother's love for her child is the underlying cause of most cases of anorexia will prevent one from being critical of the mother's mistaken methods" (Illingworth). Contrary to many popularized views, my own impression is that in many instances the seeds of these disorders are sown in the first days of life or even before birth. In a considerable proportion of cases parental anxiety has been provoked or exaggerated by a long wait for the arrival of the child, by previous miscarriages, by illness during pregnancy or a difficult labour, or by death or serious illness in the family. The resulting parental over-anxiety is often canalized in the direction of feeding, for the child's acceptance of food and of love are equated. The doctor can sympathize with the mother who, against a background of grandmotherly and neighbourly commentary, lovingly prepares meals which her child spurns.

The most important single causative factor in anorexia and food refusal is the forcing of food. Another common factor is the natural development of independence and negativism in the child. Managed wrongly, mealtimes become battlefields, and the intervals between meals restless armistices punctuated with milk, snacks, and sweets.

The first step is for the doctor to convince the mother that he appreciates her feelings. An abrupt statement that there is nothing wrong gets nowhere. If he encourages her to express her fears he will obtain important evidence on which to base further discussion and explanation. He may need, not once but repeatedly, to emphasize the absence of organic disease. He can then discuss the functions of food (for providing energy, body-building, and so on) and the wide variety of foods that are suitable. Healthy negativism in children and the importance of teaching eating habits by example will find their place in the discussions.

When the child will not eat a meal there should be no threats or punishment, no forcing or fighting, no panic or fussing, no coaxing or bribing. Nobody should take much notice. The child should go without food until the next meal, or the one after that. The doctor is justified in insisting that the child is healthy, and he must explain that a healthy child cannot possibly come to harm just because he is allowed to wait until he wants to eat. He soon will. If the doctor prescribes a "tonic" or vitamins he may undermine his own explanations. He will reinforce the parents' belief that their child is physically unwell, and distract attention from the all-important emotional background.

Sweets and snacks between meals for everybody in the family are best abolished, and the fact that milk is not a drink but a food should not just be stated but explained. The child should not sit at table indefinitely while he dawdles or refuses. The food given should be varied and attractive, without encouraging faddism. A meal a day taken at school or at home in the company of his peers often provides enough food to reassure the most anxious mother that her child is getting enough.

It will almost certainly take more than one discussion to get the parents (and even more the grandparents) to agree and to carry out the doctor's suggestions. There is no easy way, but, if they are guided and supported, their anxieties and the child's feeding difficulties are likely to be lessened or abolished, and he will be less likely to remain a "fusspot" as he grows up.

Over-eating

Over-eating may lead to obesity, the disadvantages and dangers of which are becoming more widely recognized. Over-eating is often a habit which is more easily prevented than remedied.

Plumpness in a child is taken to be a sign of good maternal care; this is in contrast to thinness and illness, which are often considered as synonymous. Food is continually pressed on the child whose health has given rise to anxiety. The overworked

or neglected mother may buy peace or affection by continually offering food as a bribe; the mother who goes out to work may salve her conscience in the same way. The child who has had rheumatic fever or some other illness for which he was taken to hospital may over-eat because of his feeling of insecurity.

The role of the family is important in over-eating. To families which are in economic difficulties food may be a symbol of prosperity, and the family which is going up in the world may celebrate its success daily with large meals. Genetic factors may play a part in obesity, but family habits of eating and imitation are certainly important.

If the parents can be brought to understand that over-eating and consequent obesity can prove harmful, they will be more likely to modify the eating habits not only of the child but of the whole family. They can show their affection in other ways than by pressing food. They may realize that it is not a punishment but a kindness to discourage the child's excessive intake of food. Nobody wants to see a child go hungry, but ordering an active child to eat less is ineffective, and drugs to depress the appetite are equally so. The tempting carbohydrates can be restricted and replaced at meals; sweets and food should be banished, with very occasional exceptions, between meals; and the child should not be rewarded with sweets for good behaviour. But the only satisfactory and lasting effects are obtained through the family.

Continued support by the family doctor may produce a modification of the habit of over-eating, but sometimes the child (or the family or environment) is so emotionally disturbed that the help of a child psychiatrist will be needed.

Anorexia Nervosa

In anorexia nervosa a pathological loss of appetite results in self-starvation which may prove fatal. It is commonest in adolescent girls, when the repugnance to food may start with an attempt to slim or after an emotional disturbance. But there is often a history of earlier feeding difficulties and of neurotic traits.

The management of the patient with anorexia nervosa cannot be discussed here in detail. It is a serious condition for which a team approach is essential, with devoted nursing, urgent attention to nutrition and metabolism, and skilled psychiatry.

Pica

This perversion of appetite usually occurs under the age of 3. The commonly held belief that it is caused by intestinal parasites has no foundation. The cause is not established, but some children with pica have behavioural disorders and in a few the intelligence is subnormal.

Earth, faeces, wood, paint, clothing, and other substances ingested may cause surprisingly little harm, but if anaemia is present or lead poisoning occurs it will need treating.

The child should be given safe objects to "mouth" and be distracted by play. Fortunately, pica is a self-limiting disorder.

The Delicate Child

The doctor's most regular and faithful clientele includes many children whose mothers describe them as "delicate." He sees them for a variety of given reasons: repeated infections, catarrh, allergies, recurrent pains, anaemia or "suddenly going pale," poor appetite or weak stomach, fatigue, lack of concentration—singly or in combination. They often miss a great deal of schooling. They are seldom really ill but seldom really well.

To confine the attention to the current symptom does little but gain a temporary respite. One of the most rewarding

experiences in the daily practice of medicine is to put this sort of problem in perspective; to consider not the symptom in isolation but the symptom in relation to the child as a person against the background of the family and environment.

He may have a cardiac murmur, or even a small congenital cardiac lesion, or a weak leg after poliomyelitis, or an occasionally wheezy chest ("Is it asthma, doctor?"). But often the physical abnormalities are quite inadequate to explain the perpetual "delicacy." The doctor will need to explore further, bearing in mind particularly the probability of "over-protection." The patient may be an only child or the youngest in a grown-up family. He may have been born to previously divorced or elderly parents, perhaps after years of waiting, or after previous miscarriages or a difficult pregnancy or labour, or he may have been premature. Another child may have died or a parent may suffer from a serious illness.

The diagnosis of an organic disorder that has persisted for years should rarely offer difficulty. A perpetual series of changing diagnoses and treatments, sometimes bolstered by witch-hunts in the laboratory, may suggest that the common denominator is to be found in the atmosphere of the home. The child's health may be a barometer of the domestic climate. It takes a very robust individual to resist the constant erosion of over-anxiety and over-protection. The real patient may be the mother or father. With sympathy and an understanding of their own fears, rather than blame or criticism, the parents may be able to modify their attitudes. If they do, the "delicacy" and the symptoms in the child may improve surprisingly and he will have a better chance to grow up robust in mind and body.

(The third part of this article will appear in next week's issue.)

ANY QUESTIONS?

We publish below a selection of questions and answers of general interest.

Apparent Resistance to Antimalarial Drugs

Q.—Expatriate patients, particularly children, are occasionally found to have mild *Plasmodium falciparum* infections despite a regular daily dosage of 100 mg. of proguanil hydrochloride. What is the reason for this? Is it just an individual variation in the blood level of the drug following oral ingestion or could it be an indication that the particular *Plasmodium* strain is becoming resistant to proguanil hydrochloride. If so, what alternative form of suppression should be employed?

A.—There are several possible explanations of this situation. It is unlikely to be due to individual variation in the blood level of the drug after ingestion, though interference with absorption by parasites, particularly worms, cannot be entirely dismissed as a possibility. It is also very unlikely that resistance to proguanil will have developed if 100 mg. has regularly been taken daily. When resistance to antimalarial drugs develops it usually does so after small irregular dosage.

Antimalarials such as proguanil are much more reliable than those who take them, and the commonest cause of trouble is observer error, the observer responsible in most instances being the patient. It is extraordinarily difficult for individuals to ensure that they take drugs regularly: memory is very fallible, and written records of drugs taken are seldom kept. In cases of this kind such recording is to be recommended. A second kind of observer error is the interpretation of artifacts as malarial parasites, which results in non-malarial pyrexias being diagnosed as malaria. This regrettably occurs much more frequently than is commonly believed.

Careful attention to regularity of drug administration and rechecking of blood films by a second and highly trained microscopist reduces almost to nil the number of cases of the kind reported by the questioner. Any encountered after these precautions have been taken should be investigated for drug resist-

ance, preferably by subinoculation. Help with facilities for this would be provided by the local medical administration or by regional malaria eradication headquarters. If resistance is proved a change to a 4-amino quinoline compound, such as chloroquine, should be considered.

Procidencia of Cervix in the Elderly

Q.—Is there anything that can be done for a severe procidencia of the cervix in a woman aged 83 who is unfit for operation, who cannot retain any sort of ring pessary, and who finds a cup-and-stem pessary so uncomfortable as to be unacceptable?

A.—The problem of intractable prolapse in old women can prove a very difficult one. Recently there have been new thoughts about fitness for operation. With careful pre-operative preparation many elderly patients can be safely submitted to a simple, quickly done repair operation without ill effect.

For those in whom operation is genuinely contraindicated the Simpson shelf pessary made in vulcanite is sometimes an appliance which succeeds when others fail. The Simpson pessary is made in various sizes. It is quite easily inserted and removed, and, provided the correct size is found, it may succeed in really difficult cases.

Powdered Alfalfa Seed

Q.—Are there any toxic factors in powdered alfalfa (lucerne) seed such as there are in soya bean and cotton seed? Its manganese and protein content is said¹ to stimulate the growth of lactobacilli in the intestine.

A.—Jacobson² states that alfalfa contains a saponin which has no pharmacological action when taken orally but which causes severe local irritation and death when given subcutaneously. It is strongly sternutatory in the dry state. Other constituents are two

proteins and a bitter principle. Wren³ states that an infusion of 1 oz. (28 g.) of the whole herb in 1 pint (0.6 litre) is prescribed in cupful doses for increasing the weight. Lucerne is also given to cattle for fattening purposes.

REFERENCES

- 1 Snell, E. E., *Nutritional Requirements of Lactic Acid Bacteria*, 1948, 11, 81. Wallerstein Laboratories Communications.
- 2 Jacobson, C. A., *J. Amer. chem. Soc.*, 1919, 41, 640.
- 3 Wren, R. W., *Potters New Cyclopaedia of Botanical Drugs and Preparations*, 7th ed., 1956. Pitman, London.

Drugs for Tremor

Q.—Efforts to speak by a child suffering from cerebral palsy are impeded by tremor of the jaw. Is there any drug therapy which would help?

A.—Involuntary movements due to organic nervous disease are very resistant to drug therapy. Atropine and its analogues, for example, relieve the rigidity of Parkinsonism, but the tremulousness of this disorder is often little influenced even by sedation to the point of drowsiness. The chance of relieving the jaw tremor in this child by drugs is remote, but there is, of course, nothing against empirical trial of amylobarbitone, benzhexol, chlorthalidate, or Valium.

Risk of Congenital Deformity

Q.—The first-born male child of a young healthy couple had clinical symptoms of congenital pyloric stenosis but recovered without surgical treatment. Their second baby, which survived 72 hours, had malformations of the heart and several other organs. Chromosomal studies on this baby showed trisomy of E₁₈ chromosome. There is no family history of congenital deformities on either side. What is the risk of subsequent children having deformities?

A.—There is no connexion between pyloric stenosis in the first child and the trisomy E₁₈ in the second child of this family.

Assuming the first child did have congenital pyloric stenosis, the risk of recurrence of the condition is in the order of 5%.¹ The risk of recurrence of the trisomy E₁₈ is prob-