Published in final edited form as:

Tetrahedron Lett. 2007 February 19; 48(8): 1395–1398.

Facile epoxidation of α,β -unsaturated ketones with cyclohexylidenebishydroperoxide

Kavitha Jakka, Jinyun Liu, and Cong-Gui Zhao*

Department of Chemistry, University of Texas at San Antonio, One UTSA Circle, San Antonio, Texas 78249-0698

Abstract

Cyclohexylidenebishydroperoxide was successfully used as the oxygen source for the oxidation of α,β -Unsaturated ketones for the first time. The corresponding epoxides were obtained in excellent yields under the Weitz-Scheffer reaction conditions.

Keywords

Enone; Epoxide; Dihydroperoxide; Hydroperoxide; Hydrogen peroxide; Epoxidation

Hydrogen peroxide (H_2O_2) is an environmentally benign oxidant, as the only by-product after oxidation is water. Unfortunately, its oxidation power towards organic compound is pretty low and, therefore, for synthetic purpose, it is normally used together with a catalyst¹ or converted to hydroperoxides² to enhance its oxidation power. *gem*-Dihydroperoxides, which have received considerable interest in recent years because of their relevance to peroxidic antimalarial agents,³ are structurally similar to hydroperoxides. However, to the best of our knowledge, they have never been utilized as oxidants in organic reactions. The reason was probably due to their difficult synthesis.⁴

Most recently, Iskra and co-workers reported a novel method for the preparation of *gem*-dihydroperoxides from ketones and H_2O_2 under iodine catalysis.⁴ This method uses readily available starting materials and is easy to operate, which paves the way for the synthetic applications of these interesting peroxides as stoichiometric oxidants. Because of our continued interest in the epoxidation chemistry,⁵ we became interested in using these peroxides as potential oxidants for epoxidation. Herein, we wish to report the first oxidation application of cyclohexylidenebishydroperoxide (**1**), which was prepared by following the reported procedure in high yield (Eq 1),⁴ in the epoxidation of α,β -unsaturated ketones.

By using *trans*-chalcone (**2a**) as the model compound and 1.0 M aqueous KOH as the base, we studied the Weitz-Scheffer epoxidation reaction in various solvents at room temperature (Table 1). It was found that nonpolar hexanes and toluene are bad solvents for this reaction, since no epoxide could be obtained in these two solvents (entries 1–2). Nonetheless, the reaction did happen in polar solvents, and the desired epoxide (**3a**) was obtained in 67%, 51% and 75% yield in MeOH (entry 3), CH_2Cl_2 (entry 4), and DMF (entry 5), respectively. Water-miscible polar solvent CH_3CN produced the epoxide in an even better 83% yield within 6 h

*Corresponding author. Tel.: +1-210-458-5432; fax: +1-210-458-7428; e-mail: cong.zhao@utsa.edu..

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

(entry 6). Further screening of other water-miscible solvents revealed dimethoxyethane (DME) and 1,4-dioxane are excellent solvents for this oxidation. The reaction in DME requires 2.5 h and the epoxide **3a** was generated in 88% yield (entry 7), whereas the reaction in dioxane requires only 2 h and the formation of the product is almost quantitative (95%, entry 8). Thus, 1,4-dioxane was identified as the best solvent for this reaction, whereas DME and CH₃CN are also very good solvents. For comparison purpose, the oxidation was also carried out with hydrogen peroxide as the oxidant in dioxane and lower yields of the epoxide were obtained with either 1 (53%, entry 9) or 2 equiv. (82%, entry 10) of the oxidant. It should be pointed out that cyclohexanone may be recovered in high yield after the reaction (91%, entry 8), which means that this is a green oxidation. (**Caution:** since compound **1** is potentially explosive, it should be handled with care. All the reactions in this study were carried out behind a safety shield inside a hood.)

To understand the scope of this reaction, we studied the reaction of various α,β -unsaturated ketones under the optimized conditions (rt, 1.0 M aqueous KOH in 1,4-dioxane),⁶ and the results are summarized in Table 2.

As is evident from Table 2, besides *trans*-chalcone (entry 1), substituted *trans*-chalcones are also good substrates for this reaction. For example, *trans*-4'-methoxychalcone yielded the desired product (**3b**) in 79% yield (entry 2). Similarly, the epoxide of *trans*-4-methoxychalcone (**3c**) was obtained in 83% yield (entry 3). Excellent yield (93%) was also obtained for the product of *trans*-2'-chlorochalcone (**3d**, entry 4). However, the reaction of *trans*-2'-hydroxychalcone did not yield the desired epoxide. Instead, the epoxide-ring-opening product (**4**) was obtained as a minor product (28%, entry 5). The major product of this reaction was the intramolecular Michael addition product **5** (53%, entry 5). These results are not surprising, because similar results are also obtained under the Weitz-Scheffer epoxidation of this substrate by using hydrogen peroxide.⁷

Acyclic enones other than chalcones are also good substrates. *trans*-3-Penten-2-one gives the epoxide product (**3f**) in quantitative yield according to GC analysis (entry 6).⁸ Likewise, the epoxide (**3g**) of *trans*-4-phenyl-3-buten-2-one was obtained in 86% yield (entry 7).

Some cyclic enones were also studied. 2-Cyclohexenone produced the epoxide (**3h**) in 89% yield (entry 8). Similarly, the epoxide of chromone (**3i**) was obtained in 86% yield (entry 9). An enedione (1,4-naphthalenedione) was also successfully epoxidized, and the product (**3j**) was obtained in excellent yield (97%, entry 10). Good yield was also obtained for the product of perinaphthenone (**3k**) (86%, entry 11). The oxidation of (*R*)-carvone generates only the *trans* diastereomers (**3l**)⁹ in 91% yield (entry 12).

In summary, we have shown that *gem*-dihydroperoxide can be used as the stoichiometric oxidant for the Weitz-Scheffer epoxidation of α,β -unsaturated ketones. Excellent yields of the epoxides may be obtained under very mild reaction conditions within short reaction times. The development of an asymmetric version of this reaction is currently under way.

Acknowledgements

The generous financial support of this research from the Welch Foundation (Grant No. AX-1593; AX-0026) and the NIH-MBRS program (Grant No. S06 08194) is gratefully acknowledged.

References and notes

1. For reviews, see: (a) Nardello V, Aubry J-M, De Vos DE, Neumann R, Adam W, Zhang R, ten Elshof JE, Witte PT, Alsters PL. *J Mol Catal A: Chem* 2006;251:185–193. (b) Lane BS, Burgess K. *Chem Rev* 2003;103:2457–2473. [PubMed: 12848577]
2. For a review, see: Kratz T, Zeiss W, Adam W. *Peroxide Chemistry* Wiley-VCH Weinheim 2000;4159
3. For leading examples, see: (a) Iskra J, Bonnet-Delpon D, Begue JP. *Tetrahedron Lett* 2003;44:6309–6312. (b) Tsuchiya K, Hamada Y, Masuyama A, Nojima M, McCullough KJ, Kim HS, Shibata Y, Wataya Y. *Tetrahedron Lett* 1999;40:4077–4080.
4. Žmitek K, Zupan M, Stavber S, Iskra J. *Org Lett* 2006;8:2491–2494. [PubMed: 16737296] and references cited therein
5. (a) Bez G, Zhao CG. *Tetrahedron Lett* 2003;44:7403–7406. (b) Adam W, Saha-Möller CR, Zhao CG. *Tetrahedron: Asymmetry* 1999;10:2749–2755. (c) Adam W, Zhao C-G. *Tetrahedron: Asymmetry* 1997;8:3995–3998.
6. General experimental procedure (Caution: Although we did not encounter any problem with compound 1, dihydroperoxides are potential explosives and should be handled with care; all reactions should be carried out behind a safety shield inside a hood): To a mixture of trans-chalcone (2a, 208 mg, 1.0 mmol) and cyclohexylidenebishydroperoxide (1, 148 mg, 1.0 mmol) in 1,4-dioxane (5.0 mL) was added 1.0 M aqueous KOH solution (1.0 mL). The reaction mixture was stirred at room temperature for 2 h, diluted with diethyl ether (15 mL) and washed with water (4 mL). The organic layer was dried over MgSO₄ and then the solvent was removed in vacuo. The crude product obtained was purified by column chromatography (95:5 hexane/ethyl acetate) to yield the pure epoxide 3a (213 mg, 95%). Cyclohexanone (89 mg, 91%) may also be recovered.
7. Loeser R, Chlupacova M, Marecek A, Opletalova V, Guetschow M. *Helvetica Chim Acta* 2004;87:2597–2601.
8. This product is very volatile and an accurate isolated yield could not be obtained.
9. Gonzalez MA, Ghosh S, Rivas F, Fischer D, Theodorakis EA. *Tetrahedron Lett* 2004;45:5039–5041.

Table 1Solvent screening in the epoxidation of *trans*-chalcone (**2a**) using cyclohexylidenebishydroperoxide (**1**).^a

Entry	Solvent	Time (h)	Yield (%) ^b
1	Hexane	24	0
2	Toluene	24	0
3	MeOH	10	67
4	CH ₂ Cl ₂	24	51
5	DMF	3	75
6	CH ₃ CN	6	83
7	MeOCH ₂ CH ₂ OMe	2.5	88
8	1,4-Dioxane	2	95 ^c
9	1,4-Dioxane ^d	2	53
10	1,4-Dioxane ^e	2	82

^a Unless otherwise specified, all reactions were carried out with *trans*-chalcone (1.0 mmol), cyclohexylidenebishydroperoxide (**1**, 1.0 mmol), aqueous KOH (1.0 M, 1.0 mL) and the solvent (5.0 mL) at room temperature.

^b Yield of isolated product after chromatographic purification.

^c 91% cyclohexanone may be recovered and reused.

^d With 30% H₂O₂ (1.0 mmol) as the oxidant.

^e With 30% H₂O₂ (2.0 mmol) as the oxidant.

Table 2

Epoxidation of α,β -unsaturated ketones (**2a-1**) using cyclohexylidenebishydroperoxide (**1**) under the optimized conditions^a

Entry	α,β -Unsaturated ketone (2)	Time (h)	Product [Yield (%) ^b]
1	(2a)	2	(3a) [95]
2	(2b)	6	(3b) [79]
3	(2c)	2.5	(3c) [83]
4	(2d)	1.5	(3d) [93]
5	(2e)	3.5	(4) [28] + (5) [53]
6	(2f)	0.5	(3f) [100] ^c
7	(2g)	3	(3g) [86]
8	(2h)	0.5	(3h) [89]
9	(2i)	5	(3i) [86]

Entry	α,β -Unsaturated ketone (2)	Time (h)	Product [Yield (%) ^b]
10	(2j)	1	(3j) [97]
11	(2k)	4	(3k) [81]
12	(2l)	2.5	(3l) [91]

^a All reactions were carried out with the α,β -unsaturated ketone (1.0 mmol), cyclohexylidenebishydroperoxide (**1**, 1.0 mmol) and aqueous KOH (1.0 M, 1.0 mL) in 1,4-dioxane (5.0 mL) at room temperature.

^b Yields of isolated products after chromatographic purification.

^c GC yield.