LIVER DISEASE

In vivo imaging of cerebral "peripheral benzodiazepine binding sites" in patients with hepatic encephalopathy

A Cagnin, S D Taylor-Robinson, D M Forton, R B Banati

Gut 2006;55:547-553. doi: 10.1136/gut.2005.075051

See end of article for authors' affiliations

Correspondence to:
Professor R B Banati,
School of Medical
Radiation Sciences and
Ramaciotti Centre for Brain
Imaging, University of
Sydney, East St PO Box
170, Lidcombe, NSW
1825, Australia;
r.banati@fhs.usyd.edu.au

Revised version received
1 September 2005
Accepted for publication
28 September 2005
Published online first
12 October 2005

Background and aim: One proposed mechanism whereby hepatic encephalopathy (HE) leads to loss of brain function is dysregulated synthesis of neurosteroids. Mitochondrial synthesis of neurosteroids is regulated by "peripheral benzodiazepine binding sites" (PBBS). Expressed in the brain by activated glial cells, PBBS can be measured in vivo by the specific ligand [^{11}C](R)-PK11195 and positron emission tomography (PET). Recently, it has been suggested that PBBS expressing glial cells may play a role in the general inflammatory responses seen in HE. Therefore, we measured PBBS in vivo in the brains of patients with minimal HE using [^{11}C](R)-PK11195 PET.

Methods: Five patients with minimal HE and biopsy proven cirrhosis of differing aetiology were assessed with a neuropsychometric battery. Regional expression of PBBS in the brain was detected by [^{11}C](R)-PK11195 PET. **Results:** All patients showed brain regions with increased [^{11}C](R)-PK11195 binding. Significant increases in glial [^{11}C](R)-PK11195 binding were found bilaterally in the pallidum, right putamen, and right dorsolateral prefrontal region. The patient with the most severe cognitive impairment had the highest increases in regional [^{11}C](R)-PK11195 binding.

Conclusion: HE is associated with increased cerebral binding of [^{11}C](R)-PK11195 in vivo, reflecting increased expression of PBBS by glial cells. This supports earlier experimental evidence in rodent models of liver failure, suggesting that an altered glial cell state, as evidenced by the increase in cerebral PBBS, might be causally related to impaired brain functioning in HE.

Chronic hepatic encephalopathy (HE) is a characteristically reversible neuropsychiatric disorder without overt structural brain damage, which occurs on a background of chronic liver disease. Minimal HE is characterised by a subtle impairment of cognitive function, leading to deficits in attention and impaired reaction times in driving or operating machinery, whereas confusion, stupor, and coma are the manifestations of overt disease.¹

One change observed consistently in the brain of mice with induced hyperammonaemia² or in post-mortem brain tissues of patients with portal-systemic encephalopathy³ is a significant increase in expression of "peripheral benzodiazepine binding sites" (PBBS). Originally discovered as additional binding sites for certain benzodiazepines, such as diazepam, PBBS are a heteromeric complex that is largely, although not exclusively, localised in the outer mitochondrial membrane.⁴ PBBS were however found to be structurally and functionally unrelated to the central benzodiazepine receptor, which is associated with gamma-aminobutyric acid (GABA) regulated channels. Subsequently, the isoquinoline PK11195, a specific high affinity ligand, has been used to pharmacologically characterise PBBS.^{5,6} Therefore, we will refer to PBBS either as "PK11195 binding (sites)" or "(R)-PK11195 binding", where the data are obtained using the R enantiomer, for which a slightly higher affinity for PBBS has been reported.⁷

Unlike in peripheral organs and cell types, normal brain parenchyma has only minimal binding of PK11195. However, as soon as brain disease gives rise to active tissue pathology, de novo expression of PK11195 binding sites is observed. At least three cellular sources have been discussed: (i) astrocytes,^{3,8} (ii) invading blood borne cells of mononuclear macrophage lineage, if the blood-brain barrier is disrupted,⁹ or (iii) activated microglia, the intrinsic population of the normally dormant brain macrophages, if the blood-brain

barrier has remained intact.¹⁰⁻¹² A role for PBBS in the pathogenesis of HE has been hypothesised, based on their regulatory effect on mitochondrial cholesterol transport and thus the altered synthesis of neurosteroids, which may be responsible for the neuronal inhibition observed in HE.^{6,13}

Additionally, recent evidence points to systemic inflammatory immune responses in minimal HE as important mediators or at least enhancers of ammonia toxicity.^{14,15} The fact that glial cells, notably microglia, the dominant immune effector cell of the brain, can rapidly express PBBS de novo in response to even the most subtle pathological stimuli is thus of added significance.^{10-12,15} The exclusively non-neuronal PBBS directly regulate many immune functions, such as release of cytokines and reactive oxygen intermediates, and are strongly regulated even in pathological conditions that are not characterised by overt tissue damage or inflammatory signs, such as recruitment of blood borne immune cells.¹¹ There is therefore the intriguing possibility that PBBS expressing glial cells are local participants in the general inflammatory responses seen in HE. Labelled with carbon-11, (R)-PK11195 is a ligand for measurement of PK11195 binding for PBBS in vivo by positron emission tomography (PET). To address the fundamental question of whether there is a change in PBBS expression in the brains of patients with minimal HE, we report measurement of cerebral [^{11}C](R)-PK11195 binding in five patients with HE and biopsy proven cirrhosis of differing aetiology. We relate the regional pattern of increased [^{11}C](R)-PK11195 binding to the underlying severity of liver disease and psychometric test analysis.

Abbreviations: HE, hepatic encephalopathy; PBBS, peripheral benzodiazepine binding sites; PET, positron emission tomography; GABA, gamma-aminobutyric acid; HCV, hepatitis C virus; NCT, number connection tests; EEG, electroencephalogram; MRI, magnetic resonance imaging; MRS, magnetic resonance spectroscopy; BP, binding potential

Table 1 Summary of the clinical characteristics for each patient with hepatic encephalopathy

	Age (y)	Sex	Aetiology	Disease duration (y)	Child grade	Bilirubin ($\mu\text{mol/l}$)	Albumin (g/l)	NH ₃ ($\mu\text{mol/l}$)	Varices	Surgical shunt (y)
Patients										
1	57	F	Autoimmune	22	B	72	34	91	—	20*
2	45	F	Alcohol	4	A	8	41	286	—	2†
3	78	F	Hepatitis C	14	A	22	37	75	+	—
4	47	M	Hepatitis C/alcohol	3	A	16	44	181	—	—
5	75	M	Alcohol	10	C	31	24	57	+	—
Control group n = 10	62.7 (12.7)‡	6F/4M								

*Surgical portocaval anastomosis.

†Transjugular intrahepatic portosystemic stent shunt.

‡Mean (SD).

SUBJECTS AND METHODS

Subjects

Five patients with HE and biopsy proven cirrhosis (mean (SD) age 60.4 (15.4) years) were recruited to the study. The patient cohort comprised two patients with alcohol related cirrhosis, one with autoimmune chronic active hepatitis/cirrhosis, one with post-viral cirrhosis secondary to hepatitis C virus (HCV) infection, and one with a dual HCV and alcohol related cirrhosis aetiology. The patient with autoimmune chronic active hepatitis/cirrhosis (table 1, patient 1) had a surgically placed portocaval shunt of more than 20 years' duration and a history of well documented minimal HE. This patient was taking maintenance oral prednisolone at 4–5 mg daily with intermittent courses of azathioprine over this 20 year period. All patients were clinically stable at the time of the study and had been abstinent from alcohol for a minimum of six months prior to the PET examination. None was receiving psychoactive medication and none was thiamine deficient. Patients with a prior history of alcohol excess had started thiamine supplementation at the time of first diagnosis.

Blood was drawn for estimation of standard biochemical and haematological parameters of liver function and for serum electrolytes. A Pugh's score and Child's grade (as modified by Pugh),¹⁶ reflecting the severity of hepatic dysfunction, were calculated for each subject (table 1). Functionally, three were Child's grade A, one Child's grade B, and one was classified as having Child's grade C disease (table 1).

A full neurological, psychometric, and electrophysiological assessment was performed on each patient. Mental state was assessed using West Haven criteria.¹⁷ Psychometric performance was assessed under standardised conditions, using a battery of four tests comprising number connection tests (NCT) A and B,¹⁸ the digit symbol subtest of the Wechsler adult intelligence scale,¹⁹ and the digit copying subscore of the Kendrick battery.²⁰ Results are shown in table 2.

Electroencephalograms (EEG) were performed using conventionally placed electrodes and mean cycle frequency was obtained. Capillary ammonia concentrations from a finger prick were measured from the pulp of the index finger, using

a blood ammonia checker II (Kyoto Daiichi Kagaku Co, Ltd, Kyoto, Japan). All patients underwent baseline T1 weighted volumetric magnetic resonance imaging (MRI) on the same day of the PET study with [¹¹C](R)-PK11195 to rule out significant structural brain pathology. Additionally, patients were characterised within one week of the PET study by in vivo cerebral ¹H magnetic resonance spectroscopy (MRS) and ³¹P MRS. Methods of data acquisition and analysis have previously been reported.^{21–23} Exclusion criteria for PET study were the presence of focal brain lesions or severe atrophy detected by MRI, presence of a past history of neurological or psychiatric disorders, or episodes of acute HE or gastrointestinal bleeding in the previous three months.

The cohort of control subjects enrolled in the PET study comprised 10 aged matched healthy volunteers (mean (SD) age 62.7 (12.7) years; range 41–80 years). Also, except for the thalamus, there was no region in the brain in which a significant age related increase in baseline binding of [¹¹C](R)-PK11195 was observed.²⁴ Each subject underwent an extensive clinical, neurological, and psychometric investigation and a T1 weighted volumetric MRI scan on the same day of the PET study with [¹¹C](R)-PK11195, to rule out systemic and neurological diseases.

The study conformed to the guidelines set out in the Declaration of Helsinki of 1975 and prior ethics approval was obtained from the Ethics Committee of the Imperial College School of Medicine, London. All subjects provided written informed consent.

PET study

The PET study was performed on a CTI/Siemens ECAT 953B PET scanner operated in three dimensional acquisition mode. [¹¹C](R)-PK11195 was injected as a bolus, 30 seconds after the acquisition scan started. Mean tracer dose was 360 (30) MBq with a specific activity of 37 (1) GBq/mmol. Dynamic data were collected over 60 minutes as 18 temporal frames. Attenuation correction factors were determined using a 15 minute transmission scan acquired before the dynamic scan. Scatter correction was achieved using a dual energy

Table 2 Results of four psychometric tests for individual patients. The normal range derives from a data set obtained from a cohort of healthy age matched controls

Patient No	NCT A (s)	NCT B (s)	Digit symbol	Digit copying
1	32	100	76	128
2	55	Not completed	50	154
3	59	Not completed	45	118
4	31	90	67	143
5	72	Not completed	19	80
Normal range	15–37 s	31–81 s	55–90	124–208

NCT, number connection tests.

Figure 1 Magnetic resonance imaging (MRI) and [^{11}C](R)-PK11195 positron emission tomography (PET) images of patients with hepatic encephalopathy. All images follow the radiological convention: the left side of the image corresponds to the subject's right side. (A, B) Transverse and coronal orientation of T1 weighted MRI images (first and third) and co-registered [^{11}C](R)-PK11195 images overlaid on the MRI (second and fourth) of patient No 1, showing MRI hyperintensity in the basal ganglia and normal [^{11}C](R)-PK11195 binding (only constitutive binding in the thalamus). (C, D) [^{11}C](R)-PK11195 PET images of patient No 2, (E, F) patient No 3, and (G, H) patient No 4. Increases in [^{11}C](R)-PK11195 binding sites were localised in the frontal lobe, particularly in the anterior cingulate cortex (F, G) and in the white matter along fibre tracts such as in the corpus callosum (C) or following projections connecting the frontal cortex with subcortical structures (E). (H) A sagittal view of patient No 4, demonstrating the spatial pattern of increase in [^{11}C](R)-PK11195 binding involving the basal ganglia and frontal lobe (arrows). In patient No 2 (D), MRI hyperintensity in the basal ganglia overlapped with the regions with an increase in [^{11}C](R)-PK11195. cc, corpus callosum; ac, anterior cingulate; wmt, white matter tract. The colour scale is calibrated for binding potential values from 0 to 1. White indicates values >1 .

window method.²⁵ Data were reconstructed with a ramp filter at Nyquist cut off, producing an image resolution of 5.8 mm (full width half maximum) at the centre of the field of view. A three dimensional T1 weighted MRI scan for the purpose of co-registration was acquired on a 1.0T Philips HPQ MRI scanner (voxel size $1 \times 1 \times 1.3$ mm; 128 contiguous slices; repetition time (TR) 35 ms, echo time (TE) 6 ms, flip angle 35°) on the same day of the PET scan.

Regional binding of [^{11}C](R)-PK11195, expressed as binding potential (BP), a measure of specific binding of the tracer, was calculated using a basis function implementation of a

simplified reference tissue model.^{26–28} Selection of an anatomically defined reference region may introduce errors since an a priori assumption that the chosen reference region is devoid of specific binding has to be made. This is particularly true under conditions where global changes in binding may occur, such as in metabolic encephalopathy. Therefore, cluster analysis^{27–29} was employed as an alternative approach for extraction of a normal ligand kinetic to serve as the reference input function, as previously described.^{28–30} For each patient the appropriate ligand kinetic was selected by χ^2 test ($p < 0.05$) comparison against a population input kinetic

previously created from the ligand kinetics of the normal cortex in healthy control patients.^{28,30}

For calculation of regional mean BP values, the following anatomical volumes of interest were defined on the individuals' volumetric MRI prior to spatial co-registration with the regional [¹¹C](R)-PK11195 binding potential map³¹: right and left temporal gyrus (superior, inferior, and middle), insula, inferior parietal lobule, anterior and posterior cingulate gyrus, dorsolateral prefrontal cortex, as well as right and left pallidum, putamen, and thalamus. The cerebellum, seen within the restricted field of view (10.65 cm) of the PET camera only to a varying extent, was excluded from formal analysis.

Statistical analysis

The Student's *t* test was used to determine the significance of the differences in regional mean [¹¹C](R)-PK11195 binding between normal control subjects and patients. Z values were calculated to test for significant increases in [¹¹C](R)-PK11195 binding in the brains of individual patients, compared with normal control brain. As [¹¹C](R)-PK11195 binding changes are unidirectional (that is, only increases) a z value of 1.6 in a one tailed z test represents a level of significance of *p* < 0.05.³² Potential error due to multiple comparisons was examined using the Hochberg correction and p-plot graphical method for estimation of the number of "true" null hypotheses.³³ Analysis of the set of *p* values revealed a widespread pattern of statistically significant changes. The estimated number of "true" null hypotheses was <3. This being the theoretical default minimum of "true" null hypotheses, it indicates that correction for multiple comparisons would not be meaningful for this data set.

RESULTS

All patients were clinically normal on examination but showed slowing of the alpha rhythm in the EEG below the reference range of 8.9 cycles per seconds and/or impaired performance in at least two of the four psychometric tests compared with the reference range for normal healthy volunteers. None had structural brain abnormalities on T1 weighted MRI, apart from hyperintensity in the basal ganglia in three of the five patients (patient Nos 1, 2, and 4), a finding common in chronic liver disease (fig 1).³⁴ All patients had cerebral metabolite abnormalities on ¹H MRS (increased glutamine/glutamate and reduced choline resonances) and on ³¹P MRS (reduced phosphomonoester, phosphodiester, and β-nucleoside triphosphate resonances), which have previously been described in patients with hepatic encephalopathy.^{21–23}

With regard to the PET study, group analysis of the patients revealed increases in [¹¹C](R)-PK11195 binding bilaterally in the pallidum (right BP: HE 0.39 (0.08), controls 0.27 (0.07), *p* < 0.01; left: HE 0.35 (0.10), controls 0.25 (0.07), *p* < 0.05), in the right putamen (BP: HE 0.31 (0.06), controls 0.25 (0.04), *p* < 0.05) and the right dorsolateral prefrontal region (BP: HE 0.27 (0.08), controls 0.15 (0.06), *p* < 0.01) (fig 2). Individual analysis demonstrated marked heterogeneity in the regional pattern of increased [¹¹C](R)-PK11195 binding (fig 1, 3). Three patients (patient Nos 2, 3, and 5) showed widespread increase in [¹¹C](R)-PK11195 signal, which was particularly pronounced in the pallidum and the frontal cortex (fig 3).

The observation appeared to be unrelated to the aetiology of the disease or other clinical characteristics, apart from the severity of cognitive decline. The three patients (patient Nos 2, 3 and 5) with significantly increased [¹¹C](R)-PK11195 binding were those impaired in the majority of the psychometric tests administered. In contrast, the two patients who

Figure 2 Mean [¹¹C](R)-PK11195 binding potential values for all volumes of interest drawn for the right (A) and left (B) hemispheres. Mean values for control subjects are expressed as empty histograms with the standard deviation bar on the top. Black dots represent individual patient values. DLPC, dorsolateral prefrontal cortex. **p* < 0.05, ***p* < 0.01.

were impaired in one test (that is, NCT B) had either normal regional mean [¹¹C](R)-PK11195 binding in all regions (patient No 1) or increased [¹¹C](R)-PK11195 binding only in the anterior cingulate cortex (patient No 4) (fig 3).

DISCUSSION

Increased PK11195 binding has been shown in experimental animal studies² and in post-mortem brain samples from patients with HE.³⁵ Our study is the first attempt to localise and quantify expression of PK11195 binding sites in vivo in patients with HE.

Regional pattern

Although increases in [¹¹C](R)-PK11195 binding were widespread in some HE patients, group analysis suggested the presence of a regional pattern, whereby the highest level of [¹¹C](R)-PK11195 binding was found in the basal ganglia, dorsolateral prefrontal regions, and anterior cingulate gyrus. Involvement of the pallidum in liver disease is widely known and T1 weighted MRI signal hyperintensities are well documented and have been ascribed to manganese deposition.³⁶ Abnormal MRI signal intensities in the pallidum correlate with poor motor performance in tasks involving speed, while cognitive decline is associated with measures of cortical atrophy.³⁷ In our study, three out of the five patients showed T1 weighted MRI signal hyperintensities in the pallidum and only one matched with a significant increase in [¹¹C](R)-PK11195 binding, indicating that increases in PBBS appear to be unrelated to MRI signal alteration.

The regionality detected by [¹¹C](R)-PK11195 PET is, instead, in keeping with other neuroimaging observations, such as decreased glucose metabolism and cerebral blood flow in the frontal-limbic-basal ganglia circuits that correlate with neuropsychological deficits.^{38,39} It is thus possible that widespread increased [¹¹C](R)-PK11195 binding with a

Figure 3 Overview of individual patient mean $[^{11}\text{C}](\text{R})\text{-PK11195}$ binding potential and standard deviation for each region of interest. The aetiology of each patient's disease is indicated in parentheses. The degree of cognitive impairment (cognitive score) is expressed as the number of impaired psychometric tests out of the 4 tested in the battery. Patient Nos 2, 3, and 5 showed a widespread increase in mean $[^{11}\text{C}](\text{R})\text{-PK11195}$ binding potential (expression of increased peripheral benzodiazepine binding sites) throughout the cortical and subcortical regions, with predominance in frontal areas and basal ganglia. These patients (Nos 2, 3 and 5) showed significant cognitive impairment.

superimposed regional pattern involving the basal ganglia and frontal regions is one pathological correlate of the cognitive impairment found in HE.

Clinical correlations

In our study, neither duration or severity of liver disease, expressed as Child-Pugh score, nor blood ammonia levels correlated with $[^{11}\text{C}](\text{R})\text{-PK11195}$ binding. However, we found instead that patients with the highest $[^{11}\text{C}](\text{R})\text{-PK11195}$ binding were the cognitively most impaired on psychometric testing.

Pathophysiological relevance of increased PK11195 binding

The apparent link between upregulation of $[^{11}\text{C}](\text{R})\text{-PK11195}$ binding and cognitive decline, the key symptom for minimal HE, supports the hypothesis that PBBS in the brain could indeed participate in the pathogenic mechanism of HE. This further implies that glial cell dysfunction is pivotal in this process, as the binding sites for PK11195 binding are

exclusively non-neuronal and, given an intact blood-brain barrier, have been shown to be expressed only by activated or reactive glia (microglia and astrocytes). One theory, the "neurosteroid hypothesis", suggests that by virtue of the regulatory influence of PBBS on the transport of cholesterol across the mitochondrial membrane,^{40–41} higher expression of PBBS increased the synthesis of neurosteroids, such as pregnenolone. Neurosteroids in turn, are potent positive allosteric modulators of GABA and/or glutamate neurotransmission.³ Several experimental studies have confirmed that conditions inducing hyperammonaemia result in an increase in PK11195 binding and brain pregnenolone synthesis and are concomitant with the clinical presentation of HE.⁴²

Although our study does not resolve the issue of the relative contribution of the different glial subtypes to the increased $[^{11}\text{C}](\text{R})\text{-PK11195}$ signal in vivo, binding sites of this ligand have been shown to be expressed only by glial cells in a reactive or dysregulated state. There is now a substantial body of evidence that the responses in a wide range of disease conditions of both astrocytes and microglia

are accompanied by profound functional changes, such as expression of cytokines, their respective receptors, and other molecules characteristic of an inflammatory tissue response, such as reactive oxygen intermediates. These changes are often referred to as "glial inflammation or neuroinflammation" as they largely occur without any recruitment of blood borne immune cells seen in classical inflammation.⁴³ Microglia and the regulatory influence that astrocytes exert over them are increasingly recognised as the cellular link between the peripheral immune system and the local immune system in the brain.⁴³ Long overlooked, microglial responses to a large variety of pathological stimuli (including subtle toxic ones) are now known to set in early and rapidly.⁴³ They are regularly found well before overt pathological changes are noted, such as in cortical spreading depression, a condition that has been referred to as "a pathology without pathology" and even in behavioural changes, such as in repetitive movements, that may be otherwise viewed as physiological.^{44–45}

In conclusion, in the brains of patients with HE that we studied, abnormally high [¹¹C](R)-PK11195 binding in the frontal lobe was detected in those patients with the poorest performance on psychometric testing. This was independent of underlying aetiology of liver disease and most likely reflects glial activation with concomitant upregulation of PBBS. The observation is in keeping with the hypothesis that PBBS may have a key role in the pathogenesis of hepatic encephalopathy via neurosteroid synthesis. Given the fact that activated, but not resting, microglia (which represent the brain's intrinsic immune effector cells) are an important source of de novo (R)-PK11195 binding, our observation also supports more recent hypotheses that in HE, like in many other primarily neurological conditions, a local inflammatory mechanism within the brain itself may act synergistically with the toxic effects of a noxious agent, which in the case of HE would primarily be ammonia.^{14–15}

ACKNOWLEDGEMENTS

This study was supported by the UK Department of Health, the British Medical Research Council (G9900178), Deutsche Forschungsgemeinschaft, and Philips Medical Systems Inc (Cleveland, Ohio, USA). Dr Adrian Lim (Consultant Radiologist, Hammersmith Hospitals NHS Trust) reported the images of the brain. Dr Camilla Buckley and Dr Shahid Khan, Gastroenterology Unit, Hammersmith Hospital, London, helped with patient recruitment. We also thank Professor Graeme M Bydder (Magnetic Resonance Unit, Imperial College London, for his valuable insight and useful discussions on study design. AC was supported by a Fellowship from the European Community (BMH4/CT98/5100) in the Training and Mobility of Researchers Programme in Biomedicine. DF was supported by a fellowship from the European Association for the Study of the Liver and the St Mary's Hospital Trustees, London.

Authors' affiliations

A Cagnin, Department of Neurosciences, University of Padova, Padova, Italy, IRCCS-Ospedale San Camillo, Venezia, Italy, and MRC Cyclotron Unit, MRC Clinical Sciences Centre, Imperial College London, Hammersmith Hospital Campus, London, UK

S D Taylor-Robinson, **D M Forton**, Robert Steiner MR Unit, Imaging Sciences Department, MRC Clinical Sciences Centre, Imperial College London, Hammersmith Hospital Campus, London, UK, and Division of Medicine A, Imperial College London, Hammersmith Hospital Campus, London, UK

R B Banati, MRC Cyclotron Unit, MRC Clinical Sciences Centre, Imperial College London, Hammersmith Hospital Campus, London, UK, Department of Neuropathology, Imperial College London, Charing Cross Hospital Campus, London, UK, and School of Medical Radiation Sciences and Ramaciotti Centre for Brain Imaging, Brain-Mind Research Institute, University of Sydney, Lidcombe, New South Wales, Australia

Conflict of interest: None declared.

REFERENCES

- 1 Schenker S, Brady CE. Pathogenesis of hepatic encephalopathy. In: Conn HO, Bircner J, eds. *Hepatic encephalopathy: syndromes and therapies*. Bloomington: Medi-Ed Press, 1994:43–61.
- 2 Desjardins P, Bandeira P, Rao VLR, et al. Portacaval anastomosis causes selective alterations of peripheral-type benzodiazepine receptor expression in rat brain and peripheral tissues. *Neurochem Int* 1999;**35**:293–9.
- 3 Butterworth RF. The astrocytic ("peripheral-type") benzodiazepine receptor: role in the pathogenesis of portal systemic encephalopathy. *Neurochem Int* 2000;**36**:411–16.
- 4 Anholt RR, Pedersen PL, DeSouza EB, et al. The peripheral-type benzodiazepine receptor. Localisation to the mitochondrial outer membrane. *J Biol Chem* 1986;**261**:776–83.
- 5 Benavides J, Cornu P, Dennis T, et al. Imaging of human brain lesions with an omega 3 site radioligand. *Ann Neurol* 1988;**24**:708–12.
- 6 Gavish M, Bachman I, Shoukrun R, et al. Enigma of the peripheral benzodiazepine receptor. *Pharmacol Rev* 1999;**51**:629–50.
- 7 Shah F, Hume SP, Pike VW, et al. Synthesis of the enantiomers of [N-methyl-¹¹C]PK 11195 and comparison of their behaviours as radioligands for PK binding sites in rats. *Nucl Med Biol* 1994;**21**:573–81.
- 8 Kuhlmann AC, Guilarte TR. Cellular and subcellular localization of peripheral benzodiazepine receptors after trimethyltin neurotoxicity. *J Neurochem* 2000;**74**:1694–704.
- 9 Myers R, Manjil LG, Cullen BM, et al. Macrophage and astrocytes populations in relation to [³H] PK11195 binding in rat cerebral cortex following a local ischaemic lesion. *J Cereb Blood Flow Metab* 1991;**11**:314–22.
- 10 Banati RB, Myers R, Kreutzberg GW. PK ("peripheral benzodiazepine")-binding sites in the CNS indicate early and discrete brain lesions: microautoradiographic detection of [³H] PK11195 binding to activated microglia. *J Neurocytol* 1997;**26**:77–82.
- 11 Banati RB. Visualising microglial activation in vivo. *Glia* 2002;**40**:206–17.
- 12 Mankowski JL, Queen SE, Tarwater PJ, et al. Elevated peripheral benzodiazepine receptor expression in simian immunodeficiency virus encephalitis. *J Neurovirol* 2003;**9**:94–100.
- 13 Ahboucha S, Desjardins P, Chatauret N, et al. Normal coupling of brain benzodiazepine and neurosteroid modulatory sites on the GABA-A receptor complex in human hepatic encephalopathy. *Neurochem Int* 2003;**45**:551–6.
- 14 Shawcross DL, Davies NA, Williams R, et al. Systemic inflammatory response exacerbates the neuropsychological effects of induced hyperammonemia in cirrhosis. *J Hepatol* 2004;**40**:247–54.
- 15 Blei AT. Infection, inflammation and hepatic encephalopathy, synergism redefined. *J Hepatol* 2004;**40**:327–30.
- 16 Pugh RNH, Murray-Lyon IM, Dawson JL, et al. Transection of the oesophagus for bleeding oesophageal varices. *Br J Surg* 1972;**60**:646–9.
- 17 Conn HO, Leevy CM, Vlahcevic ZR, et al. Comparison of lactulose and neomycin in the treatment of chronic portal-systemic encephalopathy: a double blind trial. *Gastroenterology* 1977;**72**:573–83.
- 18 Conn HO. Trail making and number-connection tests in the assessment of mental state in portal systemic encephalopathy. *Am J Dig Dis* 1977;**22**:541–50.
- 19 Wechsler D. *Wechsler adult intelligence scale manual*. New York: Psychological Corporation, 1955.
- 20 Kendrick DC, Gibson AJ, Moyes IGA. The revised Kendrick battery: clinical studies. *Br J Soc Clin Psychol* 1979;**18**:329–39.
- 21 Taylor-Robinson SD, Sargentoni J, Marcus CD, et al. Regional variations in cerebral proton spectroscopy in patients with chronic hepatic encephalopathy. *Metab Brain Dis* 1994;**9**:347–59.
- 22 Taylor-Robinson SD, Sargentoni J, Oatridge A, et al. MR imaging and spectroscopy of the basal ganglia in chronic liver disease: Correlation of T₁-weighted contrast measurements with abnormalities in proton and phosphorus-31 MR spectra. *Metab Brain Dis* 1996;**11**:249–68.
- 23 Taylor-Robinson SD, Buckley C, Changani KK, et al. Cerebral phosphorus-31 and proton magnetic resonance spectroscopy in patients with subclinical hepatic encephalopathy. *Liver* 1999;**19**:389–98.
- 24 Cagnin A, Brooks DJ, Kennedy AM, et al. In-vivo measurement of activated microglia in dementia. *Lancet* 2001;**358**:461–7.
- 25 Grooten S, Spinks TJ, Sashin D, et al. Correction for scatter in 3D brain PET using a dual energy window method. *Phys Med Biol* 1996;**41**:2757–74.
- 26 Lammertsma AA, Hume SP. Simplified reference tissue model for PET receptor studies. *Neuroimage* 1996;**4**:153–8.
- 27 Gunn RN, Lammertsma AA, Cunningham VJ. Parametric imaging of ligand-receptor interactions using a reference tissue model and cluster analysis. In: Carson R, Daule M, Witherspoon P, et al. *Quantitative functional brain imaging with positron emission tomography*. San Diego: Academic Press, 1998:401–6.
- 28 Banati RB, Newcombe J, Gunn RN, et al. The peripheral benzodiazepine binding site in the brain in multiple sclerosis: quantitative in vivo imaging of microglia as a measure of disease activity. *Brain* 2000;**123**:2321–37.
- 29 Ashburner J, Haslam J, Taylor C, et al. A cluster analysis approach for the characterization of dynamic PET data. In: Myers R, Cunningham V, Bailey D, et al. *Quantification of brain function using PET*. San Diego: Academic Press, 1996:301–6.
- 30 Cagnin A, Myers R, Gunn RN, et al. In vivo visualization of activated glia by [¹¹C](R)-PK11195-PET following herpes encephalitis reveals projected neuronal damage beyond the primary focal lesion. *Brain* 2001;**124**:2014–27.
- 31 Studholme C, Hill DLG, Hawkes DJ. Automated three dimensional registration of magnetic resonance and positron emission tomography brain images by multiresolution optimisation of voxel similarity measures. *Med Phys* 1997;**24**:25–35.

- 32 **Snedecor GW**, Cochran WG. *Statistical methods*, 7th Edn., Iowa State: University Press 1980.
- 33 **Turkheimer FE**, Smith CB, Schmidt K. Estimation of the number of "true" null hypotheses in multivariate analysis of neuroimaging data. *Neuroimage* 2001;**13**:920–30.
- 34 **Taylor-Robinson SD**, Oatridge A, Hajnal JV, *et al.* MR Imaging of the basal ganglia in chronic liver disease: Correlation of T₁-weighted and magnetisation transfer contrast measurements with liver dysfunction and neuropsychiatric status. *Metab Brain Dis* 1995;**10**:175–88.
- 35 **Lavoie J**, Layrargues GP, Butterworth RF. Increased densities of peripheral-type benzodiazepine receptors in brain autopsy samples from cirrhotic patients with hepatic encephalopathy. *Hepatology* 1990;**11**:874–78.
- 36 **Spahr L**, Butterworth RF, Fontaine S, *et al.* Increased blood manganese in cirrhotic patients: relationship to pallidal magnetic resonance signal hyperintensity and neurological symptoms. *Hepatology* 1996;**24**:116–20.
- 37 **Kulisevsky J**, Pujol J, Junque C, *et al.* MRI pallidal hyperintensity and brain atrophy in cirrhotic patients: two different MRI patterns of clinical deterioration? *Neurology* 1993;**43**:2570–3.
- 38 **Lockwood AH**, Murphy BW, Donnelly KZ, *et al.* Positron-emission tomographic localization of abnormalities of brain metabolism in patients with minimal hepatic encephalopathy. *Hepatology* 1993;**18**:1061–8.
- 39 **Catafau AM**, Kulisevsky J, Berna L, *et al.* Relationship between cerebral perfusion in frontal-limbic-basal ganglia circuits and neuropsychologic impairment in patients with subclinical hepatic encephalopathy. *J Nucl Med* 2000;**41**:405–10.
- 40 **Benavides J**, Quarteronet D, Imbault F, *et al.* Labelling of "peripheral-type" benzodiazepine binding sites in the rat brain by using [³H]PK11195, an isoquinoline carboxamide derivative: kinetic studies and autoradiographic localization. *J Neurochem* 1983;**41**:1744–50.
- 41 **Casellas P**, Galiegue S, Basile AS. Peripheral benzodiazepine receptors and mitochondrial function. *Neurochem Int* 2002;**40**:475–86.
- 42 **Itzhak Y**, Roig-Cantisano A, Dombro RS, *et al.* Acute liver failure and hyperammonemia increase peripheral-type benzodiazepine receptor binding and pregnenolone synthesis in mouse brain. *Brain Res* 1995;**705**:345–8.
- 43 **Graeber MB**. Genetics of neuroinflammation in Alzheimer disease. *Neurogenetics* 1999;**2**:135–6.
- 44 **Gehrmann J**, Mies G, Bonnekoh P, *et al.* Microglial reaction in the rat cerebral cortex induced by cortical spreading depression. *Brain Pathol* 1993;**3**:11–17.
- 45 **Ehninger D**, Kempermann G. Regional effects of wheel running and environmental enrichment on cell genesis and microglial proliferation in the adult murine neocortex. *Cereb Cortex* 2003;**13**:845–51.

POETRY IN MOTION: A PATIENT'S VIEW OF COLECTOMY

Poet Laureate of Peebles

At the end of the bed their white coats Gleaming
 Standing there some hellish plot Scheming
 One would not know from their loose formed scrum
 That the subject discussed was namely my Bum
 The Surgeons voice was crisp and clear
 As his words crossed the void between mouth and Ear:
 "Your bowel is Rotten it must come out,
 Your rectum's redundant so we'll give you a spout
 And from this protuberance you'll fix a bag,
 You'll know when it's full when your trousers sag"
 "Make sure pouch has a nice snug fit,
 For if you don't you'll get a leg full of it.
 You'll no doubt find it a bit of a joke
 When the appliance falls off when having a poke"
 Not many laughs and very few smiles
 But at least the 'op' has got rid of my piles
 I now am missing my morning treat
 When armed with the Scotsman I'd heat the loo seat
 But now when around me people all flap
 I'm quietly standing – having a crap!!

R A B Anderson

Correspondence to: Dr E Craythorne Glasgow Royal Infirmary, emma_craythorne@blueyonder.co.uk