- 4 McGivern A, Wynter CV, Whitehall VL, *et al.* Promoter hypermethylation frequency and BRAF mutations distinguish hereditary non-polyposis colon cancer from sporadic MSI-H colon cancer. *Fam Cancer* 2004;**3**:101–7.
- 5 Samowitz WS, Sweeney C, Herrick J, *et al.* Poor survival associated with the BRAF V600E mutation in microsatellite-stable colon cancers. *Cancer Res* 2005;**65**:6063–9.
- 6 Malkhosyan SR, Yamamoto H, Piao Z, *et al.* Late onset and high incidence of colon cancer of the mutator phenotype with hypermethylated hMLH1 gene in women. *Gastroenterology* 2000;**119**:598.
- 7 Elsaleh H, Powell B, McCaul K, *et al.* P53 alteration and microsatellite instability have predictive value for survival benefit from chemotherapy in stage III colorectal carcinoma. *Clin Cancer Res* 2001;**7**:1343–9.
- 8 Watanabe T, Wu TT, Catalano PJ, *et al.* Molecular predictors of survival after adjuvant chemotherapy for colon cancer. *N Engl J Med* 2001;**344**:1196–206.
- 9 Ribic CM, Sargent DJ, Moore MJ, *et al.* Tumor microsatellite-instability status as a predictor of benefit from fluorouracil-based adjuvant chemotherapy for colon cancer. *N Engl J Med* 2003;**349**:247–57.
- 10 Carethers JM, Smith EJ, Behling CA, *et al.* Use of 5-fluorouracil and survival in patients with microsatellite-unstable colorectal cancer. *Gastroenterology* 2004;**126**:394–401.

[–215G>A; IVS3+2T>C] mutation in the *SPINK1* gene causes exon 3 skipping and loss of the trypsin binding site

Previous studies have shown an association between chronic pancreatitis (CP) and mutations, especially the N34S mutation, in the serine protease inhibitor Kazal type 1 (*SPINK1*) gene.^{1,2} The human *SPINK1* gene is approximately 7.5 kb long and consists of four exons.³ The gene product consists of 79 amino acids, including a 23 amino acid signal peptide. In exon 3, *SPINK1* possesses a reactive site that serves as a specific target substrate for trypsin.⁴ It has been suggested that *SPINK1* mutations might result in altered interaction between *SPINK1* and trypsin,

Figure 1 [–215G>A; IVS3+2] mutation produced a truncated transcript. Total RNA was isolated from the biopsy specimen of the stomach, and the entire coding region of the serine protease inhibitor Kazal type 1 (*SPINK1*) gene was amplified by reverse transcription polymerase chain reaction, followed by 2% agarose gel electrophoresis. Sm, size marker (100 base pair ladders), WT, healthy control. Patient A with alcoholic chronic pancreatitis (CP) was homozygous for the [–215G>A; IVS3+2T>C] mutation. His daughter B and patient C with idiopathic CP were heterozygous. In subjects carrying the [–215G>A; IVS3+2T>C] mutation, two bands were observed: a fragment corresponding to a normal (“F1”) and a truncated (“F2”) band.

thus affecting the protease/antiprotease balance within the pancreas.^{1,2} But the underlying molecular mechanisms remain unclear. Splicing defects are estimated to account for approximately 10–15% of disease causing mutations in humans.⁵ Changes in the splicing patterns and in levels of normal transcripts lead to phenotypic differences. The prevalence of splicing mutations in the *SPINK1* gene is unknown. Most reported mutations have only been described at the DNA level and have not been studied at the mRNA level, mainly due to unavailability of *SPINK1* mRNA from patients.

We have recently shown that the [–215G>A; IVS3+2T>C] mutation is associated with familial and idiopathic CP in Japan.^{6,7} Because the IVS3+2T>C mutation affects the consensus splicing donor site,⁸ we hypothesized that this mutation leads to alternative splicing, resulting in decreased *SPINK1* function. To overcome the difficulties in obtaining human pancreas samples, *SPINK1* mRNA was harvested from the stomach, where *SPINK1* is also abundantly expressed,⁹ of (1) a 70 year old male A with alcoholic CP carrying the homozygous [–215G>A; IVS3+2T>C] mutation, (2) his 40 year old daughter B, a heterozygote who had no abdominal complaints to date, and (3) a 54 year old female C with idiopathic CP, also heterozygous for the [–215G>A; IVS3+2T>C] mutation. Total RNA was isolated from biopsy specimen of the stomach. The entire coding region of the *SPINK1* gene was amplified by reverse transcription-polymerase chain reaction (PCR) and sequenced. Electrophoresis of the reverse transcription PCR products from subjects carrying the [–215G>A; IVS3+2T>C] mutation revealed two bands: a fragment corresponding to a normal (“F1”) and a truncated (“F2”) band (fig 1). Sequencing of the truncated fragment revealed complete deletion of exon 3. This mutated protein was predicted to consist of 63 amino acids: deletion of amino acid sequence from residues 30–64 and shifting of the reading frame at amino acid 65.

To our knowledge, this is the first study showing the splicing problem in the *SPINK1* gene at the mRNA level. Northern blot analysis revealed that the size of the *SPINK1* transcript was identical both in the pancreas and stomach,⁹ suggesting that exon 3 skipping is also likely to occur in the pancreas. It is logical to assume that skipping of exon 3 would result in functional loss of *SPINK1*, thus affecting the protease/antiprotease balance within the pancreas. Of note, the daughter of patient A carrying the heterozygous [–215G>A; IVS3+2T>C] mutation has not yet developed CP. Because this mutation has not been found in healthy controls,⁷ it is of interest to see whether she will develop CP in the future. Recently, Le Marechal and colleagues¹⁰ reported the IVS2+1G>A mutation in a CP patient carrying the P55S mutation in France. The IVS2+1G>A mutation affects the consensus splicing donor site of intron 2, implying a role of another splicing variation. Further studies using larger numbers of patients and different types of mutations will establish the role of splicing mutations in *SPINK1* related CP.

Acknowledgements

This work was supported in part by a grant-in-aid from the Japan Society for the Promotion of Science (to AM and to TS).

K Kume*, A Masamune*, K Kikuta, T Shimosegawa

Division of Gastroenterology, Tohoku University Graduate School of Medicine, Sendai, Japan

Correspondence to: Dr A Masamune, Division of Gastroenterology, Tohoku University Graduate School of Medicine, 1-1 Seiryō-machi, Aoba-ku, Sendai 980-8574 Japan; amasamune@int3.med.tohoku.ac.jp

doi: 10.1136/gut.2006.095752

*K Kume and A Masamune contributed equally to this work.

Conflict of interest: None declared

References

- 1 Witt H, Luck W, Hennies HC, *et al.* Mutations in the gene encoding the serine protease inhibitor, Kazal type 1, are associated with chronic pancreatitis. *Nat Genet* 2000;**25**:213–16.
- 2 Pfitzer RH, Barmada MM, Brunskill AP, *et al.* *SPINK1*/PSTI polymorphisms act as disease modifiers in familial and idiopathic chronic pancreatitis. *Gastroenterology* 2000;**119**:615–23.
- 3 Horii A, Kobayashi T, Tomita N, *et al.* Primary structure of human pancreatic secretory trypsin inhibitor (PSTI) gene. *Biochem Biophys Res Commun* 1987;**149**:635–41.
- 4 Bartelt DC, Shapanka R, Greene LJ. The primary structure of the human pancreatic secretory trypsin inhibitor: amino acid sequence of the reduced S-aminoethylated protein. *Arch Biochem Biophys* 1977;**179**:189–99.
- 5 Stenson PD, Ball EV, Mort M, *et al.* Human gene mutation database (HGMD): 2003 update. *Hum Mutat* 2003;**21**:577–81.
- 6 Kaneko K, Nagasaki Y, Furukawa T, *et al.* Analysis of the human pancreatic secretory trypsin inhibitor (PSTI) gene mutations in Japanese patients with chronic pancreatitis. *J Hum Genet* 2001;**46**:293–7.
- 7 Kume K, Masamune A, Mizutani H, *et al.* Mutations in the serine protease inhibitor Kazal type 1 (*SPINK1*) gene in Japanese patients with pancreatitis. *Pancreatol* 2005;**5**:354–60.
- 8 Faustino NA, Cooper TA. Pre-mRNA splicing and human disease. *Genes Dev* 2003;**17**:419–37.
- 9 Marchbank T, Chinery R, Hanby AM, *et al.* Distribution and expression of pancreatic secretory trypsin inhibitor and its possible role in epithelial restitution. *Am J Pathol* 1996;**148**:715–22.
- 10 Le Marechal C, Chen JM, Le Gall C, *et al.* Two novel severe mutations in the pancreatic secretory trypsin inhibitor gene (*SPINK1*) cause familial and/or hereditary pancreatitis. *Hum Mutat* 2004;**23**:205.

Neoadjuvant chemoradiation treatment impairs accuracy of MRI staging in rectal carcinoma

Neoadjuvant chemoradiotherapy (nCRT) is considered one of the treatment modalities of advanced rectal cancer (pT3/T4 or pN+) with the intention of downsizing and downstaging the tumour. Tumour restaging may be useful for planning the operation but tissue alteration after nCRT may disturb the accuracy of the imaging procedures.

Between July 2004 and August 2005, we analysed 28 consecutive patients (18 males, 10 females, ~63 years) with adenocarcinoma of the middle and distal third of the rectum. High spatial resolution magnet resonance imaging (MRI) with intraluminal contrast and endorectal ultrasonography (EUS) (Olympus EU-M30S, 12 MHz) were performed before and after nCRT as part of their

Figure 1 Comparison of magnetic resonance imaging and histological results in T staging of tumours.

preoperative staging. Radiotherapy was delivered by daily fractions of 1.8 Gy up to a total dose of 50.4 Gy. During days 1–5 and days 29–33 of radiotherapy, 5-fluorouracil was given at a dose of 1000 mg/m²/day (maximum 1800 mg). Surgery was performed after reassessment-staging six weeks later. There were seven abdominoperineal resections and 21 anterior resections, with a covering stoma in 17 patients.

Pathological staging was undertaken according to the TNM classification. Sections were examined meticulously to identify any residual foci of adenocarcinoma. Regression of tumour size was described in 88.9% of cases (fig 1). Improvement of T or N classification was 58.3% for EUS and 55.6% for MRI, respectively. Complete remission was seen by the radiologist in four cases, confirmed histologically in seven.

T stage and N stage were predicted correctly by MRI in 57.1% and 67.9% of cases, respectively. Nodal involvement was found in 27.8%. Overestimation of tumour size and nodal involvement in MRI was seen in 37.7% and 17.9% of cases, respectively.

Depth of extramural invasion, nodal involvement, and circumferential resection margin involvement are prognostic markers in many studies.¹ Total mesorectal excision decreased local recurrence rates below 10%,² achieving further improvement by long course nCRT. Downsizing of the tumour is reported in up to 86% of patients, with a complete response rate of 9–58%; this was 25% in our study.³ The discrepancy between clinical and pathological complete response is enormous, ranging from 25% to 77%.⁴ Curative R0 resection after nCRT is possible in 83–90% of patients, which was also confirmed by our results. Three and five year survival rates are up to 82% and 71%, respectively.^{5–7} The high frequency of distant metastasis (50–60%) during follow up remains a major problem of treatment.

Accuracy of EUS and MRI is reported to be 75–94% for tumour penetration and 72–83% for nodal metastases.^{1–8} Both methods have their limitations. EUS tends to overestimate tumour depth but is highly accurate in the assessment of early tumours. MRI is commonly used in staging of pelvic malignancies because of its fine resolution but chemoradiation may decrease its accuracy. Thickening of the rectal wall after radiation by marked fibrosis, peritumoral infiltration of inflammatory cells, and vascular proliferation may contribute to overestimation of stage assessment. Sauer *et al*

estimated the “neoadjuvant chemoradiotherapy overtreatment rate” (of pT1 and pT2 tumours) caused by MRI overstaging to be 15–20%.^{8–9} In the study of Kuo *et al*, the overall accuracy rate in T staging with MRI after neoadjuvant chemoradiotherapy was only 47%, and was associated with overstaging in 47% and understaging in 6% of 36 patients.³ Our study reflected the same problems after nCRT.

Lymph nodes of >0.5 cm in diameter are considered malignant but size criteria are not very accurate. Overall accuracy rate for N staging with MRI was 64%, and was associated with overestimation of nodal involvement in 18% of cases.¹⁰

In conclusion, MRI after nCRT is correlated with impaired accuracy. MRI overestimation of residual tumour should be taken into consideration by the surgeon. Preoperative MRI after nCRT is not recommended for routine clinical use.

J Jonas, R Bähr

Department of General, Visceral and Thoracic Surgery, Municipal Hospital Karlsruhe, Karlsruhe, Germany

Correspondence to: Dr J Jonas, Department of General, Visceral and Thoracic Surgery, Municipal Hospital Karlsruhe, Moltkestr 90, 76133 Karlsruhe, Germany; joerg.jonas@klinikum-karlsruhe.com

doi: 10.1136/gut.2006.095034

Conflict of interest: None declared.

References

- 1 Brown G, Davies S, Williams GT, *et al*. Effectiveness of preoperative staging in rectal cancer: digital rectal examination, endoluminal ultrasound or magnetic resonance imaging? *Br J Cancer* 2004;91:23–9.
- 2 Bruch HP, Schwandner O. Die totale mesorektale Exzision beim Rektumkarzinom—das Dissektionsprinzip ohne kontrollierte Daten. *Viszeralchirurgie* 2002;37:6–11.
- 3 Kuo LJ, Chern MC, Tsou MH, *et al*. Interpretation of magnetic resonance imaging for locally advanced rectal carcinoma after preoperative chemoradiation therapy. *Dis Colon Rectum* 2005;48:23–8.
- 4 Pucciarelli S, Capirci C, Emanuele U, *et al*. Relationship between pathologic T-stage and nodal metastasis after preoperative chemoradiotherapy for locally advanced rectal cancer. *Ann Surg Oncol* 2005;12:111–16.
- 5 Göhl J, Merkel S, Rödel C, *et al*. Can neoadjuvant radiochemotherapy improve the results of multivisceral resections in advanced rectal carcinoma (cT4). *Colorectal Dis* 2003;5:436–41.
- 6 Wheeler JMD, Dodds E, Warren BF, *et al*. Preoperative chemoradiotherapy and total mesorectal excision surgery for locally advanced rectal cancer: Correlation with rectal cancer regression grade. *Dis Colon Rectum* 2004;47:2025–31.
- 7 Wittekind C, Tannapfel A. Regressionsgrading des präoperativ-radiochemotherapierten Rektumkarzinoms. *Pathologie* 2003;24:61–5.
- 8 Sauer R, Fiekau R, Wittekind C, *et al*. Adjuvant versus neoadjuvant radiochemotherapy for locally advanced rectal cancer. *Strahlenther Onkol* 2001;4:173–81.
- 9 Sauer R, Becker H, Hohenberger W, *et al*. Preoperative versus postoperative chemoradiotherapy for rectal cancer. *N Engl J Med* 2004;351:1731–40.
- 10 Koh DM, Brown G, Temple L, *et al*. Distribution of mesorectal lymph nodes in rectal cancer: in vivo MR imaging compared with histopathological examination. Initial observations. *Eur Radiol* 2005;15:1650–7.

BOOK REVIEW

Gastrointestinal Cancer

Edited by J A Ajanai, S A Curley, N A Janjan, P M Lynch. Berlin: Springer, 2005, £61.50, pp 394. ISBN 0387220909.

This book is part of a cancer series from the MD Anderson on the multidisciplinary management of gastrointestinal cancers, as practised in their institution. The management of gastrointestinal cancers is a rapidly evolving and advancing field, with outcomes improving markedly over the past five years. This is particularly true for colorectal cancers (CRC) and gastrointestinal stromal tumours but there have also been small but significant advances in gastro-oesophageal and pancreatic cancers. There is a less clear impact in primary liver tumours, although there is consistency in phase III data for chemoembolisation of hepatocellular cancers. Finally, the outcome for neuroendocrine tumours has remained unchanged. Where there have been significant advances, the new targeted biological therapies have finally come to clinical fruition and need to be appropriately integrated into routine clinical practice.

The complexity of integrated gastrointestinal multidisciplinary advances make this a timely, concise, and compact addition to those practising in this area. As other books in the series tackle specific cancers, the task of encompassing the management of essentially eight or more disease specific areas, linked by the digestive system anatomically, is difficult in such a small book, and so naturally there are biases within particular topics. The authors have decided to structure the book differently from other similar specialist texts by initially grouping and attempting to encompass all gastrointestinal cancers in the first six chapters on staging, pathology, imaging, and a specific chapter on laparoscopic approaches. This works for some topics, but as the book is not designed as a reference text, it may disengage readers who are lower gastrointestinal specialists as opposed to upper gastrointestinal specialists, and vice versa. The pathology chapter, for example, is really only of interest to those involved in the care of CRC. Similarly, the key points at the end of the initial six chapters presumably reflect the disease specific bias of the authors.

The contributions arise from a wide range of specialties, many of whom are experienced leaders in their fields. Some chapters, such as the ones on CRC screening, familial aspects, genetics, and the chapter on gastrointestinal stromal tumours are concise yet comprehensive and very well written. The others represent discussions on selected areas of management. However, this does of necessity lead to some omissions and biases. For example, there is no clear discussion on neoadjuvant chemotherapy for downstaging colorectal liver metastases but yet considerable detail on hepatic artery infusion. Discussion on hepatocellular cancer and transarterial chemoembolisation implies there is no randomised evidence of benefit for a subset. The chapter on pancreatic cancer focuses on locally advanced disease and chemoradiation (on which there is a paucity of positive phase III data), but the discussion on systemic chemotherapy is in the palliative care section at the end of the book. There is a specific chapter on bowel management in patients with cancer which is an excellent