PostScript

LETTERS

Long term suppressive therapy with pegylated interferon for chronic hepatitis C associated membranoproliferative glomerulonephritis

Renal disease secondary to chronic hepatitis C virus (HCV) infection is infrequent but often life threatening and difficult to treat. Management is directed towards eradication of the virus. Combination therapy with interferon α or pegylated interferon α and ribavirin has been used successfully to induce recovery from membranoproliferative glomerulonephritis (MPGN), cryoglobulinaemia, and vasculitis, if viral clearance is achieved.¹⁻⁸ Unfortunately, symptoms recur in the absence of a sustained virological response. Long term suppressive maintenance therapy may be an alternative, but has not been reported previously.

History

A 46 year old man with HCV genotype 4 was referred to our institution for evaluation of chronic HCV infection (CHC) and nephrotic syndrome. Initial investigations revealed proteinuria (+++) and haematuria (+++) with dysmorphic red cells on urinalysis. Haemoglobin level (Hb) was 136 g/L, alanine aminotransferase (ALT) 243 U/L, albumin 18 g/L, creatinine 0.12 mmol/L (0.06–0.13 mmol/L), creatinine clearance 1.47 ml/s, and urinary protein excretion 7.6 g/day. Cryoglobulin levels were repeatedly <1%. Renal biopsy was consistent with MPGN. Liver biopsy demonstrated chronic active hepatitis with a Scheuer fibrosis stage of 2. Treatment with interferon α 3 million units

three times a week and ribavirin 1200 mg daily was commenced in November 1998 (fig 1). After seven weeks, ALT had declined to 14 U/L and HCV RNA was not detected (<100 copies/ml). Progressive severe symptomatic anaemia (Hb 74 g/L) however necessitated cessation of ribavirin.

In May 1999, while on interferon monotherapy, the HCV RNA test returned positive and ALT levels increased to 84 U/L. Ribavirin was reinstituted at 600 mg daily (maximum tolerated dose) and combination therapy was continued to August 2000 and then ceased. Over the ensuing 15 months, the patient developed significant renal impairment (creatinine 0.39 mmol/L) and proteinuria which peaked at 34 g/day. HCV RNA testing was positive.

In November 2001, the patient was commenced on pegylated interferon α -2b 1.5 μ g/kg/week. Ribavirin was not reinstituted due to poor tolerability and prior lack of efficacy. After six months, liver tests returned to normal, albumin had risen to 36 g/L, and creatinine had declined to 0.13 mmol/L. Furthermore, proteinuria declined to 4.5 g/day and the HCV RNA was not detected. After 12 months of suppressive therapy, ALT, aspartate aminotransferase (AST), albumin, and creatinine were normal; proteinuria had improved to 0.54 g/L and HCV-RNA was negative. After 21 months of treatment, the pegylated interferon dose was reduced to 75% of the initial dose. Within weeks, the patient became HCV PCR positive. Full dose therapy was reinstituted. After eight weeks, HCV PCR was again negative and ALT, AST, albumin, and creatinine were normal. Suppressive therapy for 44 months to February 2006 has been well tolerated. HCV polymerase chain reaction remains negative and liver tests, serum albumin, and creatinine are within normal limits.

Discussion

This is the first report of long term successful suppressive antiviral therapy with pegylated interferon for CHC. After failure of 15 months of combination therapy complicated by symptomatic anaemia, the patient was maintained on suppressive doses of pegylated interferon for more than four years. This has resulted in a sustained clinical remission, undetectable HCV RNA, and resolution of the nephrotic syndrome. An attempt at dose reduction was followed promptly by a return to serum HCV RNA positivity, suggesting that therapy was unable to eradicate low level viraemia. Long term pegylated interferon use has been relatively well tolerated and was considered more cost effective than the alternative of chronic renal failure with its associated symptoms and need for dialysis.

In conclusion, we report the successful long term use of suppressive antiviral therapy for HCV associated MPGN. Viral suppression is a suitable and practical alternative for life threatening extrahepatic manifestations of HCV infection, if viral clearance cannot be achieved.

I H Y Cua, V Kwan, M Henriquez

Storr Liver Unit, Westmead Millennium Institute, Department of Gastroenterology and Hepatology, University of Sydney at Westmead Hospital, Westmead, New South Wales, Australia

J Kench

Department of Tissue Pathology, Institute of Clinical Pathology and Medical Research, University of Sydney at Westmead Hospital, Westmead, New South Wales, Australia

J George

Storr Liver Unit, Westmead Millennium Institute, Department of Gastroenterology and Hepatology, University of Sydney at Westmead Hospital, Westmead, New South Wales, Australia

Correspondence to: Professor J George, Department of Medicine, Westmead Hospital, Westmead, NSW 2145, Australia; jacob.george@wmi.usyd.edu.au

doi: 10.1136/gut.2006.094508

Conflict of interest: None declared.

References

- 1 Alric L, Plaisier E, Thebault S, et al. Influence of antiviral therapy in hepatitis C virus-associated cryoglobulinemic MPGN. *Am J Kidney Dis* 2004;43:617–23.
- 2 Garini G, Allegri L, Carnevali L, et al. Interferon-alpha in combination with ribavirin as initial treatment for hepatitis C virus-associated cryoglobulinemic membranoproliferative glomerulonephritis. *Am J Kidney Dis* 2001;38:E35.
- 3 Jefferson JA, Johnson RJ. Treatment of hepatitis C-associated glomerular disease. *Semin Nephrol* 2000;20:286–92.
- 4 Rossi P, Bertani T, Baio P, et al. Hepatitis C virus-related cryoglobulinemic glomerulonephritis: long-term remission after antiviral therapy. *Kidney Int* 2003;63:2236–41.
- 5 Sabry AA, Sobh MA, Sheaaah HA, et al. Effect of combination therapy (ribavirin and interferon) in HCV-related glomerulopathy. *Nephrol Dial Transplant* 2002;17:1924–30.
- 6 Sarac E, Bastacky S, Johnson JP. Response to high-dose interferon-alpha after failure of

Figure 1 Effect of antiviral therapy on 24 hour protein excretion and serum alanine aminotransferase (ALT) levels. HCV, hepatitis C virus; MU tiw, million units, three times a week; RBV, ribavirin; ULN, upper limit of normal.

standard therapy in MPGN associated with hepatitis C virus infection. *Am J Kidney Dis* 1997;**30**:113–15.

- 7 Bruchfeld A, Lindahl K, Stahle L, et al. Interferon and ribavirin treatment in patients with hepatitis C-associated renal disease and renal insufficiency. *Nephrol Dial Transplant* 2003;**18**:1573–80.
- 8 Orlent H, Mathot RA, Van Bommel EF, et al. Peginterferon and dose-titrated ribavirin for hepatitis C-associated nephrotic membranoproliferative glomerulonephritis type 1. *Dig Dis Sci* 2005;**50**:1804–6.

Diagnosis of lymphogranuloma venereum from biopsy samples

The Health Protection Agency Centre for Infections launched an alert in October 2004 to improve the awareness, diagnosis, and control of lymphogranuloma venereum (LGV), a sexually transmitted chlamydial infection, following a series of outbreaks in Western Europe.¹ To date (9/3/2006), 334 cases of LGV have been diagnosed in 334 men. The case definition for a confirmed case of LGV is the presence of *C trachomatis* specific DNA, using two nucleic acid amplification tests (NAATs) with different primers, of serovars L1, L2, or L3, determined by genotyping (http://www.hpa.org.uk/infections/topics_az/hiv_and_sti/LGV/lgv.htm). All cases of LGV to date in the UK have been in men who have sex with men and typically present with proctitis and/or inguinal lymphadenopathy. Some of the men in the UK diagnosed with LGV reported long duration of symptoms presenting to gastroenterologists and having been wrongly diagnosed with inflammatory bowel disease before referral to genitourinary medicine (GUM).

Initially, the Sexually Transmitted Bacteria Reference Laboratory (STBRL) only accepted swabs, residual or unprocessed NAAT specimens, and enzyme immunoassay samples, which we had previously validated as being suitable for testing. In January 2005 we were approached about accepting a histology specimen from a male patient who in retrospect the clinician suspected as having LGV. Since this date we have processed seven histological specimens, which we now report.

Genomic DNA was extracted from formalin fixed paraffin embedded biopsies using a QIAamp DNA mini kit (Qiagen, Crawley, Sussex). The paraffin was removed by incubating at 56°C and then we proceeded with the manufacturer's recommended tissue protocol without any modifications. *C trachomatis*

Figure 1 Rectal biopsy of patient No 5. The lymph node architecture is preserved but severely distorted. There is a florid active and chronic inflammatory process within the node and extending into the surrounding soft tissues. This consists of epithelioid granulomata showing severe central necrosis which is geographic in outline. There is a severe acute inflammatory component with microabscess formation. There is also follicular hyperplasia.

was confirmed to be present in the extracted DNA by two different real time polymerase chain reaction methods, each having different gene targets. These targets are different to other commercially available NAATs. The genotype of the *C trachomatis* was then determined by the method of Lan and colleagues.² Since September 2005, all samples positive for *C trachomatis* were tested for LGV using the real time polymerase chain reaction method of Morre and colleagues.³

Seven biopsies were processed from six patients attending GUM clinics in the UK (table 1).

Of seven biopsy samples submitted to STBRL in the past nine months for LGV investigation, we have successfully been able to extract DNA from the biopsies and diagnose three cases of LGV in two patients (Nos 3, 4 and 5), two that were not LGV associated (Nos 1 and 7), and two samples were negative for *C trachomatis* (Nos 2 and 6) (table 1). A rectal biopsy from patient No 5 is shown in fig 1. It is interesting that two patients (Nos 1 and 7) presented with clinical symptoms characteristic of LGV that were determined to be infected with chlamydia, but of a non-LGV associated serovar. Further clinical research is indicated to consider whether these chlamydial infections were causing the severe inflammation of proctitis

and inguinal lymphadenopathy, or if these were incidental findings.

A diagnosis of LGV should be taken into consideration by gastroenterologists and histologists in patients presenting with proctitis or inguinal lymphadenopathy, particularly in men who have sex with men. Examination of biopsy tissue may be useful for diagnosing LGV in patients where cases are being re-examined due to the increased awareness of LGV in the UK, or where a swab for *C trachomatis* testing was not obtained. The algorithm for accepted samples has now been altered to include histological specimens.

Acknowledgments

We would like to thank all of the clinicians for submitting the biopsy samples, for providing information on the patients, and for reviewing this report: Dr Duncan Churchill, Brighton GUM; Dr Zubayr Sulaiman Gloucester GUM; Dr Colm O'Mahony Chester GUM; Dr Ben Goorney, Sexual Health Clinic, Salford; Dr Sarah Schoeman, Leeds GUM; Jamie Hardie, Newham GUM.

I M C Martin, S A Alexander, C A Ison
Sexually Transmitted Bacteria Reference Laboratory,
Health Protection Agency Centre for Infections,
London, UK

N Macdonald
HIV and STI Section, Health Protection Agency Centre
for Infections, London, UK, and Department of
Infectious Disease Epidemiology, Imperial College
Faculty of Medicine, London, UK

K McCarthy
Department of Pathology, Cheltenham General
Hospital, Cheltenham, UK

H Ward
HIV and STI Section, Health Protection Agency Centre
for Infections, London, UK, and Department of
Infectious Disease Epidemiology, Imperial College
Faculty of Medicine, London, UK

Correspondence to: Dr I Martin, Sexually Transmitted
Bacteria Reference Laboratory, Health Protection
Agency Centre for Infections, 61 Colindale Avenue,
London NW9 5HT, UK; iona.martin@hpa.org.uk

doi: 10.1136/gut.2006.096370

Conflict of interest: None declared.

References

- 1 Macdonald N, Ison CA, Martin I, et al. Initial results of enhanced surveillance for lymphogranuloma venereum (LGV) in England.

Table 1 Biopsy results from six patients attending genitourinary medicine clinics in the UK

Biopsy No	Site of biopsy	<i>C trachomatis</i> result	<i>C trachomatis</i> genotyping result	LGV specific real time result	Symptoms	Histological findings
1	Rectum	Positive	Not LGV associated	Not available	Proctitis, fibrotic stricture of rectum	Chronic inflammation, ulceration of mucosa
2	Rectum	Negative	–	Negative	Proctitis, bloody diarrhoea, weight loss,	Severe inflammation, goblet cell depletion, distortion crypt architecture
3, 4	Lymph nodes of left groin	Positive	L2 (LGV associated)	Positive	Inguinal lymphadenopathy	Lymph node necrosis, inflammatory infiltrate
5	Rectum	Positive	Not amplified	Positive	Colo-proctitis	Inflammatory infiltrate with early abscess formation
6	Left inguinal lymph node	Negative	–	Negative	Inguinal lymphadenopathy	Perivascular inflammation
7	Right groin	Positive	Not LGV associated	Not available	Inguinal lymphadenopathy	Suppurative granuloma reaction

Euro Surveillance 2005, volume 10. <http://www.eurosurveillance.org/ew/2005/050127.asp#5> (accessed 8 May 2006).

- 2 **Lan J**, Ossewaarde JM, Walboomers JMM, *et al*. Improved PCR sensitivity for direct genotyping of *Chlamydia trachomatis* serovars by using a nested PCR. *J Clin Micro* 1994;**32**:528–30.
- 3 **Morre SA**, Spaargaren J, Fennema JS, *et al*. Molecular diagnosis of lymphogranuloma venereum. PCR-based restriction fragment length polymorphism and real-time PCR. *J Clin Microbiol* 2005;**43**:5412–13.

Efficacy of ecabet sodium enema on steroid resistant or steroid dependent ulcerative colitis

Ecabet sodium (ES), a 12-sulfodehydroabietic acid monosodium salt derived from an ingredient in pine resin, has been shown to topically enhance several gastric mucosal defensive factors.¹ ES is widely used for the treatment of gastritis and gastric ulcer in Japan. A recent open label study demonstrated that ES also has therapeutic effects in active ulcerative colitis (UC).² In that study, the authors demonstrated that six of seven UC patients with proctosigmoiditis achieved complete remission by treatment with ES enema. However, successive clinical trials of the efficacy of ES enema in UC patients have not been reported. Here we demonstrate the efficacy of ES enema in steroid resistant or steroid dependent UC patients.

Treatment with ES enema was performed after obtaining informed consent from steroid dependant, steroid resistant, or steroid contraindicated UC patients who had inflamed lesions mainly in the rectum and sigmoid colon except for one patient with ileal pouchitis after total proctocolectomy. This clinical trial was approved by the ethics committee of Akita University School of Medicine. Patient profiles are shown in table 1. ES enema was prepared with 1 g of ES and 20–50 ml of tepid water, and one or two enemas per day were instilled into the rectum. Disease activity was assessed before and after treatment using the ulcerative colitis-disease activity index (UC-DAI).³ In case No 1, the endoscopic point was not included in the score. Doses of drugs for UC, including sulfasalazine, 5-aminosalicylic acid, and prednisolone were not changed or increased during treatment with ES enema.

One patient (case No 6) was withdrawn because of a sense of abdominal fullness. Two of five patients (case Nos 1 and 2) significantly responded to the treatment and achieved almost clinical remission in three and seven weeks, respectively. Endoscopic findings in the rectum of case No 2 before and after treatment are shown in fig 1A and 1B. One patient with pouchitis (case No 4) dramatically responded to treatment after two weeks. Two patients (case Nos 3 and 5) with deep ulcers in the rectum and sigmoid colon also responded to the treatment but did not achieve clinical remission. Endoscopic findings of case No 3 before and after the treatment are shown in fig 1C and 1D. Mean disease activity index declined by 54.5% after ES enema treatment compared with that before treatment ($p = 0.0289$).

Our results suggest that ES enema may have therapeutic actions in some steroid resistant or steroid dependent UC patients. ES enema also had therapeutic effects in the patient with ileal pouchitis. However, the effect of ES enema in two cases with deep ulcer was slight and clinical remission was

Table 1 Patients profiles

Case	Sex	Age (y)	Extent of lesions	Clinical course	Intractability
1	F	45	Left sided colitis	Relapsing remitting	PS dependent
2	M	44	Left sided colitis	Relapsing remitting	Irritation to PS
3	F	41	Left sided colitis	Chronic continuous	PS resistant
4	F	33	Pouchitis	Relapsing remitting (post colectomy)	PS induced bone necrosis
5	M	23	Left sided colitis	Relapsing remitting	PS resistant
6	F	32	Left sided colitis	Relapsing remitting	PS resistant

PS, prednisolone.

not achieved. The detailed functional mechanism of ES enema for improvement in inflammation in colonic mucosa was not clearly elucidated but induction of several defensive factors and an antioxidative stress function⁴ are suggested as being involved. Further randomised case controlled trials with larger number are needed to assess the efficacy of ES enema in patients with UC.

M Iizuka, H Ito, S Konno, T Shirasaka, Y Horie, K Shindo, S Watanabe

Department of Internal Medicine, Akita University School of Medicine, Akita, Japan

Correspondence to: Dr M Iizuka, Department of Internal Medicine, Akita University School of Medicine, 1-1-1 Hondo, Akita 010-8543, Japan; maizuka@doc.med.akita-u.ac.jp

doi: 10.1136/gut.2006.099440

Conflict of interest: None declared.

References

- 1 **Kinoshita M**, Kume E, Tamaki H. Roles of prostaglandins, nitric oxide and the capsaic insensitive sensory nerves in gastroprotection produced by ecabet sodium. *J Pharmacol Exp Ther* 1995;**275**:494–501.
- 2 **Kono T**, Nomura M, Kasai S, *et al*. Effect of ecabet sodium enema on mildly to moderately active ulcerative proctosigmoiditis: An open-label study. *Am J Gastroenterol* 2001;**96**:793–7.
- 3 **Sutherland LR**, Martin F, Greer S, *et al*. 5-Aminosalicylic acid enema in the treatment of distal ulcerative colitis, proctosigmoiditis, and proctitis. *Gastroenterology* 1987;**92**:1894–8.
- 4 **Sasaki K**, Iizuka M, Konno S, *et al*. Ecabet sodium prevents the delay of wound repair in intestinal epithelial cells induced by hydrogen peroxide. *J Gastroenterol* 2005;**40**:474–82.

Figure 1 Endoscopic findings of the rectum in case Nos 2 case 3 before and after treatment with ecabet sodium (ES) enema. (A) Case No 2, before treatment; whitish coated shallow erosions are seen diffusely. (B) Case No 2, after treatment; lesions have improved dramatically. (C) Case No 3, before treatment; deep ulcers can be seen diffusely. (D) Case No 3, after treatment; deep ulcers are partially improved.

EUS could detect ascites missed by CT scan

I read with great interest the article by Lee and colleagues (*Gut* 2005;54:1541–5) regarding the accuracy of endoscopic ultrasonography (EUS) in diagnosing ascites and predicting peritoneal metastases in patients with gastric cancer. I would like to thank the authors for quoting our study.¹ Lee and colleagues commented that the sensitivity of detection of ascites was lower in our study and that this might be due to the use of catheter probe. I would however like to point out that such a comparison was unfair as the two studies were fundamentally different in two ways.

(1) The patient populations of the two studies were different. Our study excluded all patients with evidence of ascites on physical examination or computed tomography (CT) scan. We believe that there is no need for an additional EUS to confirm the presence of ascites in such patients. Moreover, such patients should have paracentesis for cytological examination rather than a locoregional staging investigation like EUS.

During the study period of our paper (September 1995 to January 2002), 89 patients had evidence of ascites on CT scan or physical examination. There would be no difficulty in detecting ascites in these patients by EUS. If we included these patients, the detection rate by EUS would have been 25.5% ($36+89/402+89 = 125/491$). The overall incidence of ascites was 29.5% ($56+89/402+89 = 145/491$). The “adjusted” sensitivity would be 84.8% ($34+89/56+89 = 123/145$), not much lower than the 87.1% reported by Lee *et al.*

(2) CT scan was performed for all patients in our study. Of the 89 patients who were excluded from the study, 69 had evidence of ascites on CT scan. The sensitivity of CT scan for detection of ascites was 47.6% ($69/145$), higher than the 16.1% sensitivity (combined US and CT scan) reported by Lee *et al.* We suspect the lower sensitivity may be due to the predominant use of US scan (231 patients) rather than CT scan (99 patients) in their study. A direct comparison however is not possible as the study populations of the two studies may be different.

In summary, I would like to applaud Lee *et al.* for their systematic study. The comment that the sensitivity of detection of ascites was lower in our study was unfair.

K-M Chu

Correspondence to: Professor K-M Chu, Department of Surgery, The University of Hong Kong, Queen Mary Hospital, Pokfulam, Hong Kong, China; chukm@hku.hk

Conflict of interest: None declared.

Reference

- 1 Chu KM, Kwok KF, Law S, *et al.* A prospective evaluation of catheter probe EUS for the detection of ascites in patients with gastric carcinoma. *Gastrointestinal Endoscopy* 2004;59:471–4.

Author's reply

We would like to thank Dr Chu for his interest in our paper and for supplementing some of the data missing in his previous publication.¹ We agree that both studies had fundamental difference in study design and

we would like to point out the major difference for discussion.

One of the major differences lies in the definition of presence of ascites. Chu *et al.* used positive laparoscopic detection of ascites as the gold standard when comparing with their endoscopic ultrasonography (EUS) findings. However, in our study, we combined all investigation results (ultrasound (US), computed tomography (CT), EUS, and operation) to determine the status of ascites. We believe that a small amount of fluid, such as several millilitres as detected by EUS, might not be visibly appreciated during operation.

Another major difference is the type of EUS used in the two studies. We used an echoendoscope (7.5–12 MHz) with a higher penetration depth, which allowed scanning through thick tumours. In contrast, Chu *et al.* used a miniprobe (20 MHz) in their study, which actually had a limited depth of penetration. This may explain why the overall incidence of ascites was higher in our study (37.2%) compared with Chu's series (29.5%, the supplemented data combining CT, physical examination, and operative finding). In our study, EUS was more sensitive (87.1%) than the operative findings (40.9%) in detecting ascites. Therefore, we do not agree with Chu's conclusion that “laparoscopy and laparotomy remain the reference standard for the detection of ascites” as “ascites was missed by EUS in nearly 40% of patients”.¹ We suspect if an echoendoscope was used in Chu's study, the sensitivity of EUS would have increased and CT decreased, and the projected results would then have come close to ours.

In fact, CT scan is not our routine in the preoperative assessment of patients with gastric cancer, especially in those early gastric cancers, or if EUS showed no local invasion. There is no evidence in the literature showing that CT scan is better than US in the detection of ascites. CT scan could not be used as the gold standard as, according to Chu's data, CT scan missed 20 patients with ascites which was indeed detectable on physical examination.

We are however agreed on the same conclusion that “EUS is useful for the detection of ascites in patients with gastric carcinoma” and “The presence of ascites was significantly associated with peritoneal seeding”.¹

Y T Lee, E K W Ng

Institute of Digestive Disease, Chinese University of Hong Kong, Hong Kong

Correspondence to: Dr Y T Lee, Institute of Digestive Disease, Chinese University of Hong Kong, Shatin NT, Hong Kong; leeytong@cuhk.edu.hk

Conflict of interest: None declared.

Reference

- 1 Chu KM, Kwok KF, Law S, *et al.* A prospective evaluation of catheter probe EUS for the detection of ascites in patients with gastric carcinoma. *Gastrointestinal Endosc* 2004;59:471–4.

Anti-TNF- α therapy for orofacial granulomatosis: proceed with caution

A 36 year old female patient was reviewed at the gastrointestinal clinic. She had, since the age of 15 years, been affected by troublesome

orofacial granulomatosis (OFG) manifest as lower lip swelling together with a midline fissure. In the past she had received numerous therapies, including intralesional and systemic corticosteroid (short term benefit only), cinnamon and a benzoate free diet (lack of compliance), azathioprine (intolerance), and topical tacrolimus (ineffective).

In 2001, due to increasing distress about her appearance compounded by her forthcoming wedding, we decided to treat her with an infliximab infusion at 5 mg/kg. Within seven days there was a noticeable improvement, followed by complete healing of her labial fissure six weeks later, just prior to her wedding. Two weeks after this she found out she was pregnant. She subsequently gave birth to a healthy baby boy but failed to attend the clinic and was lost to follow up for four years.

On re-referral to the clinic in 2005, she was in the midst of a course of oral prednisolone (prescribed by her general practitioner) as her OFG had once again become problematic. In view of her excellent previous response to anti-tumour necrosis factor α (TNF- α) therapy coupled with the significant risks of an infusion reaction if rechallenged with infliximab (long drug “holiday” with no concomitant immunosuppression), we elected to treat her with subcutaneous adalimumab, 80 mg initially and then 40 mg fortnightly. After five weeks of treatment there was both a subjective and objective improvement, with partial healing of the midline fissure (figs 1, 2). At eight weeks the patient noted some left sided facial pain and swelling just below the corner of her mouth. She attended her dentist who excluded any periodontal sepsis. Three days later she was admitted to our unit with fever sweats and worsening facial pain and swelling (fig 3). Clinically she had a perioral cellulitis with bilateral perioral swelling and erythema, together with pyrexia and raised inflammatory indices. She received

Figure 1 Pre-adalimumab treatment; swollen lower lip with deep midline fissure.

Figure 2 At five weeks, after three adalimumab injections; marked improvement in midline fissure.

Figure 3 At eight weeks, after four adalimumab injections; although the midline fissure continued to heal, there was now a florid bilateral perioral cellulitis and the patient was systemically unwell.

intravenous benzylpenicillin and flucloxacillin to which there was minimal response but there was a rapid resolution of the cellulitis with intravenous piperacillin. Her blood cultures were negative. Adalimumab therapy was terminated immediately.

OFG is a chronic inflammatory disorder of the orofacial tissues characterised by non-caseating granulomas on biopsy.¹ Numerous Crohn's therapies have been used to treat this condition, although due to the relative rarity of OFG, none has been subjected to randomised controlled trials. Thus physicians have to base their treatment decisions on small case series. Anti-TNF- α therapy has been used to treat OFG, with success reported with both thalidomide and infliximab.^{2,3} Adalimumab is a recently developed fully human IgG1 monoclonal antibody to TNF- α and preliminary data have shown this drug to have similar efficacy to infliximab in those Crohn's patients intolerant to⁴ or in whom response has become attenuated⁵ with infliximab.

It has become commonplace for gastroenterologists to actively exclude sepsis when considering infliximab therapy for inflammatory bowel disease, as will be the case for adalimumab if and when it is fully licensed. This is clearly difficult in OFG, a disease characterised by facial pain, swelling, erythema, and mucosal breaks. In addition, the oropharyngeal mucosa, the presumed portal of bacterial entry in this case, is colonised by a wide variety of organisms in health, thus swabbing this region prior to anti-TNF therapy will almost certainly give positive results, but is unlikely to assist in the decision to give or withhold therapy. Furthermore, patients will almost certainly learn to self administer this medication and without proper warnings it is conceivable that patients could continue to take this medicine in the context of worsening sepsis.

This case highlights that while anti-TNF- α therapy may have a therapeutic role in OFG, extreme caution and close monitoring must be undertaken in those patients who receive it.

D R Gaya, S Aitken, J Fennell, J Satsangi, A G Shand

Gastrointestinal Unit, Molecular Medicine Centre, University of Edinburgh, Western General Hospital, Edinburgh, UK

Correspondence to: Dr D R Gaya, Gastrointestinal Unit, Molecular Medicine Centre, University of Edinburgh, Western General Hospital, Edinburgh EH4 2XU, UK; danielgaya@aol.com

doi: 10.1136/gut.2006.099143

Conflict of interest: None declared.

References

- 1 Weisenfeld D, Ferguson MM, Mitchell DN, *et al*. Oro-facial granulomatosis—a clinical and pathological analysis. *Q J Med* 1985;**54**:101–13.
- 2 Hegarty A, Hodgson T, Porter S. Thalidomide for the treatment of recalcitrant oral Crohn's disease and orofacial granulomatosis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2003;**95**:576–85.
- 3 Mahadevan U, Sandborn WJ. Infliximab for the treatment of orofacial Crohn's disease. *Inflamm Bowel Dis* 2001;**7**:38–42.
- 4 Youdim A, Vasilakias EA, Targan SR, *et al*. A pilot study of adalimumab in infliximab-allergic patients. *Inflamm Bowel Dis* 2004;**10**:333–8.
- 5 Papadakis KA, Shayne OA, Vasilakias EA, *et al*. Safety and efficacy of adalimumab (D2E7) in Crohn's disease patients with an attenuated response to infliximab. *Am J Gastroenterol* 2005;**100**:75–9.

Adverse events in clinical trials with azathioprine and mesalamine for prevention of postoperative recurrence of Crohn's disease

We read with great interest the study by Ardizzone and colleagues (*Gut* 2006;**55**:47–53) and the excellent review of Sands (*Gut* 2006;**55**:437–41) commenting on the efficacy and side effects of azathioprine (AZA) in the therapy of ulcerative colitis. Ardizzone *et al* observed in their investigator blinded study, which included patients with steroid dependent ulcerative colitis, more mild to moderate adverse events in azathioprine than in mesalamine (5-ASA) treated patients (26% *v* 6%; *p* = 0.046). However, only two of 36 patients on AZA were withdrawn from the study because of adverse events. We would like to comment on the side effects of AZA, which we observed in a double blind, double dummy, randomised, prospective, multicentre study on the efficacy and safety of AZA (2.0–2.5 mg/kg/day) and 5-ASA (4 g/day) for prevention of postoperative endoscopic recurrence in Crohn's disease.

Seventy nine patients (AZA, 42; 5-ASA, 37) were randomised within two weeks after surgery. TPMT genotyping was performed at baseline in order to exclude subjects with homozygous TPMT deficiency. However, the study was stopped prematurely because an interim analysis revealed that the hypothesis of superiority of AZA versus 5-ASA could not be tested with the planned sample size. In 37 patients (AZA, 18; 5-ASA, 19) who completed the study according to the protocol (treatment for one year), the primary study end point (treatment failure: severe endoscopic relapse, withdrawal due to clinical relapse or to adverse drug reaction) was evaluated.

Treatment failure was found to be equally high in each group (AZA, 9 of 18; 5-ASA, 9 of 19; *p* = 1.00, two sided Fisher's exact test). Six of 18 patients on AZA and two of 19 patients on 5-ASA therapy were withdrawn because of adverse drug reactions (33% *v* 11%; *p* = 0.12, two sided Fisher's exact test); reasons were leucopenia/anaemia (AZA, 1; 5-ASA, 1), elevated liver enzymes, arthralgia/myalgia, vomiting, abdominal pain, macroscopic fecal excretion of study medication (AZA, 1 each), and pancreatitis (5-ASA, 1). Clinical or severe endoscopic relapse was observed in three of 18 patients on AZA therapy and in seven of 19 patients on 5-ASA therapy (17% *v* 37%; *p* = 0.27, two sided Fisher's exact test). Considering all 79 patients, adverse events were reported in

approximately 70% of patients in each group (AZA, 29 of 42; 5-ASA, 26 of 37). Furthermore, in three of 42 "non-completers" an intolerable adverse event led to withdrawal (AZA, ileus; 5-ASA, cholecystitis, ankylosing spondylitis). Two further trials investigating the efficacy and side effects of AZA to prevent postoperative relapse of Crohn's disease have been published recently.^{1,2}

In an open label study by Ardizzone and colleagues,¹ adverse events were observed more frequently (39% *v* 25%) in patients receiving AZA (2 mg/kg/day) than in those receiving 5-ASA (3 g/day). Fifteen of 69 patients in the AZA group and six of 69 patients in the 5-ASA group were withdrawn because of adverse events (22% *v* 9%; *p* = 0.04); reasons for withdrawal were leucopenia/thrombocytopenia (AZA, 7; 5-ASA, 0), elevated liver enzymes (AZA, 4; 5-ASA, 1), pancreatitis (AZA, 3; 5-ASA, 0), epigastric intolerance (AZT, 1; 5-ASA, 2), and increased serum creatinine (AZT, 0; 5-ASA, 3). In a double blind placebo controlled trial by Hanauer and colleagues,² nine of 47 (19%) patients receiving a relatively low dose of 6-mercaptopurine (6-MP 50 mg/day), six of 44 (14%) patients receiving 5-ASA (3 g/day), and four of 40 (10%) patients on placebo were withdrawn from the study because of adverse events, respectively; reasons for withdrawal were diarrhoea (6-MP, 2, 5-ASA, 2), leucopenia (6-MP, 2; 5-ASA, 0), alopecia (6-MP, 2; 5-ASA, 0), elevated liver enzymes (6-MP, 0; 5-ASA, 1), flatulence, gastrointestinal bleeding, phlebitis (6-MP, 1 each), and allergic reaction, bowel obstruction, and arthralgia (5-ASA, 1 each).

In summary, we could not provide evidence for the superiority of AZA over 5-ASA in our prospective clinical trial. In contrast with the trials described above, we observed a higher rate of adverse drug reactions leading to withdrawal from the study in the AZA group. Placebo controlled trials are needed urgently to address the question of best postoperative immunosuppressive management.³ However, our observations indicate the difficulties that may arise in future trials for reaching an adequate statistical power to provide a valid answer to this question.

H Herfarth

Department of Internal Medicine, University of Regensburg, Regensburg, Germany

C Tjaden

Department of Surgery, University of Heidelberg, Heidelberg, Germany

M Lukas

IV Interni Klinika, Charles University, Praha, Czech Republic

F Obermeier

Department of Internal Medicine, University of Regensburg, Regensburg, Germany

K Dilger, R Müller

Dr Falk Pharma GmbH, Freiburg, Germany

J Schölmerich

Department of Internal Medicine, University of Regensburg, Regensburg, Germany

A Z T-1 Study Group

Correspondence to: Professor H Herfarth, Division of Gastroenterology and Hepatology, CB#7080 Bioinformatics Bldg, University of North Carolina, Chapel Hill, 27599 North Carolina, USA; hherf@med.unc.edu

Conflict of interest: None declared.

References

- 1 **Ardizzone S**, Maconi G, Sampietro GM, *et al*. Azathioprine and mesalamine for prevention of relapse after conservative surgery for Crohn's disease. *Gastroenterology* 2004;**127**:730–40.
- 2 **Hanauer SB**, Korelitz BI, Rutgeerts P, *et al*. Postoperative maintenance of Crohn's disease remission with 6-mercaptopurine, mesalamine, or placebo: a 2-year trial. *Gastroenterology* 2004;**127**:723–9.
- 3 **Sandborn WJ**, Feagan BG. The efficacy of azathioprine and 6-mercaptopurine for the prevention of postoperative recurrence in patients with Crohn's disease remains uncertain. *Gastroenterology* 2004;**127**:990–3.

Impaired portal circulation resulting from L-arginine deficiency in patients with lysinuric protein intolerance

Lysinuric protein intolerance (LPI) is a rare autosomal recessive defect of the dibasic amino acid transporter, which is caused by mutations in the *SLC7A7* gene and results in L-arginine deficiency.¹ L-arginine is the only known substrate for nitric oxide (NO) production,² and thus the availability of L-arginine is crucial for regulation of NO production.³ NO is an endothelium derived vascular relaxation factor that plays an important role in vascular endothelial function.^{4–6} However, the effects of L-arginine deficiency on portal circulation in humans have not been studied.

Portal circulation was evaluated in seven Japanese LPI patients (six males). Seven healthy subjects (six males) without liver disease served as controls. The study was approved by the ethics committee of our university. Mean age of the LPI patients was 18.9 (10.7) years (range 5–32) and that of the controls 15.6 (7.3) years (range 7–26). Mean body weight of the patients was 38.0 (11.4) kg (range 14.6–50.3) and that of the controls 42.1 (13.6) kg (range 19.0–58.0). There was no significant difference in age or body weight between the two groups. Also, there was no significant difference in body mass index between the two groups (LPI patients 18.7 (1.9) kg/m² (range 16.4–21.2); controls 17.4 (2.0) kg/m² (range 13.2–18.9)).

Six patients were homozygotes of a common mutation in the *SLC7A7* gene (R410X) and one was a compound heterozygote with two mutations (S238F and R410X).^{7,8} In all LPI patients, plasma levels of L-arginine (38.5 (8.1) nmol/ml (range 22.9–47.6)) were lower than the lower limit of the normal range (54–130 nmol/ml), and plasma levels of L-lysine and L-ornithine were also less than the normal range, except for one patient.

Portal blood flow volume (PBFV) and portal vein diameter (PVD) were determined by ultrasonography.^{9,10} Serum levels of NO derivatives and PBFV before and after administration of L-arginine or an NO donor, isosorbide dinitrate (ISDN), were evaluated. Fasting serum levels of NO derivatives in patients (67.3 (8.9) μ mol/l) were significantly lower than those in controls (120.3 (19.2) μ mol/l) ($p < 0.01$) (fig 1A). PBFV in patients (411.3 (173.1) ml/min) was significantly lower than in controls (837.5 (214.2) ml/min) ($p < 0.01$) (fig 1B). PVD in patients (6.1 (1.4) mm) was also significantly lower than in controls (8.5 (1.3) mm) ($p < 0.01$) (fig 1C). Serum NO levels increased from baseline in the three patients given ISDN and in the three patients given L-arginine: from 71.8 (8.9) to 103.5

(6.2) μ mol/l in patients receiving ISDN, and from 64.5 (10.3) to 95.6 (9.1) μ mol/l in patients receiving L-arginine. Serum NO levels after administration of ISDN or L-arginine to patients were almost restored to those in controls. PBFV increased from baseline in the three patients given ISDN and in the three patients given L-arginine: from

301.3 (96.8) to 687.6 (111.8) ml/min in patients receiving ISDN and from 521.3 (170.9) to 1119.3 (192.6) ml/min in patients receiving L-arginine. PBFV after administration of ISDN or L-arginine was restored to close to that in controls (fig 1D, E).

It is still not known whether L-arginine administration can improve portal circulation

Figure 1 Serum levels of nitric oxide (NO) derivatives (nitrite and nitrate) were measured with an NO₂/NO₃ Assay Kit-Fil (Wako Pure Chemical Industries, Osaka, Japan). Isosorbide dinitrate (5 mg/5 kg up to a maximum of 40 mg) was administered to three patients transdermally for three hours, and L-arginine hydrochloride (0.5 g/kg) was infused to three other patients intravenously for one hour. Statistical analysis was performed using the Mann-Whitney U test or Wilcoxon signed rank test. (A) Fasting concentrations of serum NO derivatives in patients with lysinuric protein intolerance (LPI) and controls. Baseline data for portal vein flow in LPI patients and controls. Baseline portal blood flow volume (PBFV) (B) and baseline portal vein diameter (PVD) (C). Changes in PBFV in LPI patients before and after isosorbide dinitrate (D) or L-arginine (E) administration. The shaded area represents the mean (SD) of the PBFV in controls. Three LPI patients were given isosorbide dinitrate (D) and three were given L-arginine (E).

by restoring endothelial cell function of the portal vein in patients. However, the present study suggests that L-arginine deficiency in LPI can cause impairment of portal circulation via a decrease in NO production.

T Takeda, H Watanabe, T Saito, K Saito, H Takeda, H Togashi

Department of Gastroenterology, Faculty of Medicine, Yamagata University, Yamagata, Japan

J Fujii

Department of Biochemistry, Faculty of Medicine, Yamagata University, Yamagata, Japan

Y Takasago

Morioka Children's Hospital, Morioka, Japan

S Kawata

Department of Gastroenterology, Faculty of Medicine, Yamagata University, Yamagata, Japan

Correspondence to: Professor S Kawata, Department of Gastroenterology, Faculty of Medicine, Yamagata University, 2-2-2 Iida-nishi, Yamagata 990-9585, Japan; kawata@med.id.yamagata-u.ac.jp

doi: 10.1136/gut.2005.086603

Conflict of interest: None declared.

References

- 1 Simell O. Lysinuric protein intolerance and other aminoacidurias. In: Scriver CR, Beaudet AL, Sly WS, et al, eds. *The metabolic and molecular bases of inherited disease*, vol 3, 7th edn. New York: McGraw-Hill, 1995:3603–27.
- 2 Sakuma I, Stuehr DJ, Gross SS, et al. Identification of arginine as a precursor of endothelium-derived relaxing factor. *Proc Natl Acad Sci U S A* 1988;**85**:8664–7.
- 3 Kilbourn RG, Belloni P. Endothelial cell production of nitrogen oxides in response to interferon γ in combination with tumor necrosis factor, interleukin-1, and endotoxin. *J Natl Cancer Inst* 1990;**82**:772–6.
- 4 Palmer RMJ, Ferrige AG, Moncada S. Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature* 1987;**327**:524–6.
- 5 Rubanyi GM, Romero JC, Vanhoutte PM. Flow-induced release of endothelium-derived relaxing factor. *Am J Physiol* 1986;**250**:H1145–9.
- 6 Kamada Y, Nagaretani H, Tamura S, et al. Vascular endothelial dysfunction resulting from L-arginine deficiency in a patients with lysinuric protein intolerance. *J Clin Invest* 2001;**108**:717–24.
- 7 Noguchi A, Shoji Y, Koizumi A, et al. SLC7A7 genomic structure and novel variants in three Japanese lysinuric protein intolerance families. *Human Mutat* 2000;**15**:357–72.
- 8 Koizumi A, Shoji Y, Nozaki J, et al. A cluster of lysinuric protein intolerance (LPI) patients in a northern part of Iwate, Japan due to a founder effect. *Human Mutat* 2000;**16**:270–1.
- 9 Taourel P, Blanc P, Dauzat M, et al. Doppler study of mesenteric, hepatic, and portal measurements and the severity of portal hypertension and hepatic failure. *Hepatology* 1998;**28**:932–6.
- 10 Piscaglia F, Zironi G, Gaiani S, et al. Systemic and splanchnic hemodynamic changes after liver transplantation for cirrhosis: A long-term prospective study. *Hepatology* 1999;**30**:58–64.

Association study of *TNFSF15* polymorphisms in Japanese patients with inflammatory bowel disease

Tumour necrosis factor superfamily (TNFSF) 15 is a novel member of the TNFSF and its mRNA and protein expression is upregulated in inflammatory bowel disease (IBD), particularly in Crohn's disease (CD).^{1,2} Recently, Yamazaki *et al* performed a large scale case control study using single nucleotide polymorphism (SNP) markers and reported that polymorphisms in *TNFSF15* conferred susceptibility to CD in both Japanese and UK populations.³ They also suggested a potential association between a Caucasian ulcerative colitis (UC) cohort and *TNFSF15*, but this association was not studied in Japanese patients. To investigate this possible association between *TNFSF15* and Japanese UC, and to replicate this association with CD in Japanese, we performed a case control association study in Japanese patients with CD and UC.

We selected six SNPs—*tnfsf15*₂₆, *tnfsf15*₂₈, *tnfsf15*₃₁, *tnfsf15*₃₃, *tnfsf15*₃₅, *tnfsf15*₃₆—which were reported to show a strong association ($p < 10^{-10}$), and genotyped these six SNPs in 286 patients with CD, 263 patients with UC, and 277 healthy controls (HCs) by polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) analysis. All patients and HCs were Japanese, and none had a family history of IBD.

We replicated all six SNPs that were significantly associated with the Japanese CD cohort (table 1). On the other hand, in contrast with Caucasians,³ none of the SNPs were associated with the Japanese UC cohort. Risk allele frequencies of *tnfsf15*₃₅, *tnfsf15*₃₆, and *tnfsf15*₃₁ were higher in the CD group with anal lesions than in those without, but the associations were statistically weak ($p = 0.019$, 0.019 , and 0.037 , respectively). No significant differences were found in allele frequencies between the CD subgroups classified by age at diagnosis, location of disease, existence of fistula, stenosis, need for steroid therapy, and past history of surgical treatment, and also the UC subgroups classified by age at diagnosis, extent of disease, need for intensive intravenous steroid therapy, and need for surgical treatment.

In this study, we confirmed the findings of a previous report concerning a significant association between *TNFSF15* and CD. On the other hand, no evidence for an association with Japanese UC was observed, although a potential association with Caucasian UC was reported. It is generally accepted that UC and CD may share some susceptibility genes. Ethnic differences in genetic susceptibility may be explained by differences in the haplotypic background. Some *TNFSF15* polymorphisms identified in the Japanese were monomorphic or nearly monomorphic in the Caucasian population.³ Thus it seems likely that population specific patterns of haplotypes may contribute to differences in UC susceptibility.

Although a previous report described that *tnfsf15*₂₈ showed the lowest p value and highest odds ratio among the SNPs in *TNFSF15*,³ our results showed that p values and odds ratios of *tnfsf15*₃₆ and *tnfsf15*₃₅ were similar to those of *tnfsf15*₂₈. Thus to identify the pathogenic SNP, a functional study is clearly needed. We analysed the transcription factor binding sites in the promoter region of *TNFSF15* by TFSEARCH⁴ and found that GATA-1, 2, and 3 possibly bind the *tnfsf15*₃₅-T allele while the GATA binding *cis* element is absent in the *tnfsf15*₃₅-C allele (risk associated). It is well known that GATA-3 promotes a Th2 mediated immunological state and suppresses expression of Th1 mediated cytokines.^{5,6} These findings have raised the possibility that GATA-3 may not bind the promoter region with the *tnfsf15*₃₅-C risk allele resulting in lack of suppression of *TNFSF15* expression. Consequently, over-expressed *TNFSF15* promotes a Th1 mediated immunological state and initiates or exacerbates the severity of CD. Although we do not have experimental evidence for this hypothesis, we intend to elucidate the functional significance in the future.

Y Kakuta, Y Kinouchi, K Negoro, S Takahashi, T Shimosegawa

Division of Gastroenterology, Tohoku University Graduate School of Medicine, Sendai, Japan

Correspondence to: Dr Y Kakuta, Division of Gastroenterology, Tohoku University Graduate School of Medicine, 1-1 Seiryō, Aoba, Sendai, 980-8574, Japan; ykakuta@mail.tains.tohoku.ac.jp

doi: 10.1136/gut.2006.100297

Conflict of interest: None declared.

References

- 1 Migone TS, Zhang J, Luo X, et al. TL1A is a TNF-like ligand for DR3 and TR6/DcR3 and functions as a T cell costimulator. *Immunity* 2002;**16**:479–92.

Table 1 Allele frequencies of *TNFSF15*

SNP	dbSNP	Location	Allele1	Allele2	Frequency of allele2 (%) / OR / p value		
					HCS (n=277)	CD (n=286)	UC (n=263)
<i>tnfsf15</i> ₃₆	rs7848647	5'-flanking region	G	A	39.0	25.7 (OR = 1.85, $p = 1.84 \times 10^{-6}$)	36.3 (NS)
<i>tnfsf15</i> ₃₅	rs6478109	5'-flanking region	C	T	39.0	25.7 (OR = 1.85, $p = 1.84 \times 10^{-6}$)	36.3 (NS)
<i>tnfsf15</i> ₃₃	rs6478108	Intron 1	A	G	51.8	37.6 (OR = 1.78, $p = 1.60 \times 10^{-6}$)	52.3 (NS)
<i>tnfsf15</i> ₃₁	rs4979462	Intron 1	A	G	39.5	26.2 (OR = 1.84, $p = 1.98 \times 10^{-6}$)	36.5 (NS)
<i>tnfsf15</i> ₂₈	—	Intron 3	C	T	53.1	38.8 (OR = 1.78, $p = 1.58 \times 10^{-6}$)	52.3 (NS)
<i>tnfsf15</i> ₂₆	rs3810936	Exon 4 (Val201Val)	G	A	40.3	26.9 (OR = 1.83, $p = 2.15 \times 10^{-6}$)	36.3 (NS)

SNP, single nucleotide polymorphism; HCs, healthy controls; CD, Crohn's disease; UC, ulcerative colitis; NS, no significant. OR, odds ratio.

- 2 **Bamias G**, Martin C iii, Marini M, et al. Expression, localization, and functional activity of TL1A, a novel Th1-polarizing cytokine in inflammatory bowel disease. *J Immunol* 2003;**171**:4868–74.
- 3 **Yamazaki K**, McGovern D, Ragoussis J, et al. Single nucleotide polymorphisms in TNFSF15 confer susceptibility to Crohn's disease. *Hum Mol Genet* 2005;**14**:3499–506.
- 4 **Heinemeyer T**, Wingender E, Reuter I, et al. Databases on transcriptional regulation: TRANSFAC, TRRD and COMPEL. *Nucleic Acids Res* 1998;**26**:362–7.
- 5 **Zheng W**, Flavell RA. The transcription factor GATA-3 is necessary and sufficient for Th2 cytokine gene expression in CD4 T cells. *Cell* 1997;**89**:587–96.
- 6 **Ferber IA**, Lee HJ, Zonin F, et al. GATA-3 significantly downregulates IFN-gamma production from developing Th1 cells in addition to inducing IL-4 and IL-5 levels. *Clin Immunol* 1999;**91**:134–44.

NOTICE

The XXXI Pan American Congress of Digestive Diseases: Global challenges in gastroenterology from the end of the world

This conference will be held in Santiago, Chile, on 11–14 November 2008.

EDITOR'S QUIZ: GI SNAPSHOT

Answer

From question on page 1422

An abdominal operation revealed a hard whitish tumour. The tumour involved that part of the small intestine (fig 2). On pathological examination it was diagnosed as a metastasis from a pleomorphic carcinoma of the lung (fig 3). Metastasis of lung carcinoma to the small intestine has been reported to be in the range 2.6–10.7% in autopsy studies.^{1–2} Some cases of small intestine metastasis showed various symptoms, such as obstruction, malabsorption, haemorrhage, and perforation. Berger *et al* reported that 0.5% of patients operated on for lung carcinoma developed symptomatic small intestine metastases.³ In this case, the shape of the small intestine metastasis was unusual. Cancer cells were substituted for normal cells all round the intestinal wall. Intestinal fluid passed through inside the tumour so that, although he had

slight abdominal pain after meals, he did not have obstruction. Fortunately, it did not perforate.

Careful examination for intra-abdominal lesions is needed after resection of primary lung carcinoma. If acute abdomen occurs in patients with a known history of lung carcinoma, gastrointestinal metastasis must be considered.

doi: 10.1136/gut.2005.090035

References

- 1 **McNeill PM**, Wagman LD, Neifeild JP. Small bowel metastases from carcinoma of the lung. *Cancer* 1987;**59**:1486–9.
- 2 **Antler AS**, Ough Y, Pitchumoni CS, et al. Gastrointestinal metastases from malignant tumor of the lung. *Cancer* 1982;**49**:170–2.
- 3 **Berger A**, Cellier C, Daniel C, et al. Small bowel metastases from primary carcinoma of the lung: clinical findings and outcome. *Am J Gastroenterol* 1999;**94**:1884–7.

Figure 2 The tumour. The small intestine was segmented by the tumour but intestinal juice passed through the lumen of the tumour.

Figure 3 Pathological findings. On pathological examination, the structures of both the primary tumour (A) and metastatic tumour (B) were made of spindle cells and giant cell carcinoma. The similarity of the two tumours strongly supports the idea that the abdominal mass is a secondary.