COVER ILLUSTRATION

....deceived with ornament

In Shakespeare's *Merchant of Venice*, Portia is the clever heroine with beauty and wit who fulfils her role as a mistress of disguise. Her namesake spider has similar qualities and is the undoubtedly the queen of the family of jumping spiders—Salticidae.

Portia fimbriata is a spider's spider. The family Salticidae numbers 5000 known species, and all are aggressive carnivores, even of other spiders. *Portia*, though, goes a step beyond that, as she prefers to hunt jumping spiders; as a result she has acquired special adaptations.

Jumping spiders are positively charming creatures, and you will know that to be true if you have ever watched one closely. These are common spiders and range from approximately 3–17 mm in length; they will watch you closely as you approach them. They have four pairs of eyes, with the large anterior median (AM) set the most obvious. These circular eyes provide an "attentive child" appearance because they are fixed and are relatively large based on body size, but they are tiny on an absolute scale. These placid eyes belie the organised complexity and evolutionary genius that lies beneath the carapace.

The AM eyes are Galilean telescopes with a corneal lens fixed to the carapace, and a second "lens" at the end of a small tube immediately in front of the retina. This second lens deserves a closer inspection. It consists of a steeply sloped foveal-like pit. There is an index of refraction gradient between the amorphous fluid in the tube (analogous to vitreous) and the wall of the foveal-like pit as this tissue contains the retina. This creates a minus lens and a Galilean telescope much like the convexiculate fovea of many raptors. A creature this small cannot have much retinal area, so to maximise the number of photoreceptors struck by photons, the retina is tiered. This means that there are four layers of retinal cells for light to traverse. Each layer of retina extracts information from the light passing through the rhabdom into the next tier until it strikes the fourth and final layer. The angle of separation from the centre

of one rhabdom to another perpendicular to incident light is approximately 1.7 μm , creating a mosaic slightly more than three times the wavelength of light.

This visual system is indeed unique and begins with the aforementioned corneal lens connected to an elongated tube within the cephalothorax (first segment of the spider's body). There are six muscles to move the tube. Since the external cornea/lens is fixed to the carapace, it creates an image that is fixed at one point within the tube. This compact telephoto lens system combined with the tiered retina achieves excellent acuity, but only a very tiny field of vision. So, to increase the field of acute vision, this optical marvel moves the tube housing the retina, with the six attached muscles per eye, by using a variety of mostly conjugate eye movements, including pendular and rotary movements in a form of scanning eye. *Portia* scans its world much like painting a wall with a fine brush although the retina is not linear, but shaped more like a boomerang (extraordinary seminal work done by Land M, *J Exp Biol* 1969;51:443–70 and 471–93). The other pairs of eyes do not scan and are principally used as motion detectors to find other animals for the AM eyes to decipher (Harland DP, Jackson RR. In: FR Prete, ed, *Complex Worlds from Simpler Nervous Systems*. Cambridge, MA: MIT Press, 2004:5–40).

With the AM eyes, jumping spiders have the finest discrimination of all arthropods, and probably all invertebrates, as they are visual hunters; most other spiders use the tools of silk. But *Portia* goes a step beyond that, for she hunts other Salticids—other sharp eyed

predators with vision as good as hers. To do so, she must have more than just good vision. She must have stealth, patience, crypsis, and cunning.

Portia does. This magnificent spider has several techniques that allow it to stalk other Salticidae and larger spiders. As it stalks its prey, it may alert or alarm the intended victim who may turn to see the predator. If *Portia* is exposed and is not in range, it will freeze and since it is cryptically coloured, it will look like a piece of detritus. *Portia* will keep its palps hidden, as these and its legs apparently are key to identification by other spiders. Or, if it is hunting an orb weaver, it has an aggressive camouflage technique to lure the spider by plucking the web as if it were a caught insect. This plucking will lure the poor sighted orb weaver to its death. Or, if necessary, she will stalk with extreme patience using wind gusts to ply the silk fibres (Jackson RR, Pollard SD, *Annual Review of Entomology* 1996;41:287–308).

Portia is heuristic and learns to detour if the prey cannot be easily approached. For example, if her prey can be approached only from one angle, say the edge of a leaf, *Portia* can do this even when the prey will be obscured by the detour. This may not seem difficult, but *Portia* must have a plan and remember where the prey was located since the detour may block visualisation of its prey. Following the detour, once *Portia* gets close enough, perhaps 3–4 body lengths, it leaps, bites, and injects a venom that rapidly will kill the prey. Surprisingly, *Portia* can and does kill larger and much more venomous spiders.

The bard would have been proud.
"So may the outward shows be least themselves;

The world is still deceived with ornament."
Bassanio 3.2 *Merchant of Venice*

I R Schwab,
University of California Davis, 4860 Y Street,
Suite 2400, Sacramento, CA 95817, USA;
irschwab@ucdavis.edu

R Jackson,
University of Canterbury, Christchurch, New
Zealand

Cover photographs by Robert Jackson, PhD.
Thanks to Duane Harland and Michael Land
for their comments on the essay.