unselected consecutive series of 223 secondary and tertiary periocular BCC referrals to a UK regional oculoplastic service. Our case mix should therefore be representative of that in other similar UK units and our results are therefore more relevant to them than those of the high risk biased Australian Mohs series and our results can be used as an audit benchmark. Although we have used conventional non-Mohs treatment, we achieved a very low recurrence rate.

We fully support that full histological margin control, as in Mohs micrographic surgery, is the best treatment option in recurrent, high risk cases. Our study shows that careful non-Mohs BCC excision remains a valuable, safe and cost effective option. Our technique offered the patients with primary non-infiltrative BCC excellent cure rates and functional and cosmetic outcomes with no recurrence over the 5 year follow up.

S Hamada

Corneo Plastic Unit, Queen Victoria Hospital, Holtye Road, East Grinstead, RH19 3D2, UK

V T Thaller

Royal Eye Infirmary, Apsley Road, Plymouth PL4 6PL, UK

Correspondence to: Mr V T Thaller, Royal Eye Infirmary, Apsley Road, Plymouth PL4 6PL, UK; vladimir.thaller@phnt.swest.nhs.uk

doi: 10.1136/bjo.2006.095968

Accepted for publication 4 April 2006

Reference

- 1 Hamada S, Kersey T, Thaller VT. Eyelid basal cell carcinoma: non-Mohs excision, repair and outcome. *Br J Ophthalmol* 2005;**89**:992–4.

NAION and treatment of erectile dysfunction: reply from Pfizer

McGwin *et al*¹ suggest that treatment of erectile dysfunction with sildenafil (Viagra) or tadalafil (Cialis), two phosphodiesterase inhibitors (PDE5-I), may increase the odds of non-arteritic anterior ischaemic optic neuropathy (NAION) in men with a history of myocardial infarction (MI) or hypertension (HTN). We believe, however, that this study is fraught with significant limitations (as the authors acknowledge) that preclude drawing any conclusions about this relation.

Case-control studies are susceptible to several biases that must be carefully considered and controlled for in the study design, implementation, and analysis.² For example, accurate and complete ascertainment of exposure is critical in a retrospective case-control study because both disease and exposure occur before study initiation. The authors note that the interviewers were *not* blinded to the case or control status of the patient, making it possible that the interviewer may, even unconsciously, probe cases differently from controls for exposure to a PDE5-I (interviewer bias). Another obstacle is that patients were not consistently interviewed at the time of the NAION event (or index date for controls) about their PDE5-I usage. Hence, NAION patients may have been more likely to remember drug usage (recall bias). Furthermore, exposure misclassification may have occurred as timing, dose, and duration of drug use relative to event onset were not captured (exposure bias). This information is crucial for drugs used as

needed such as PDE5-I, and particularly for short half life drugs like sildenafil.

Perhaps, the most troublesome weakness of the study was the limited sample size and differential participation rates of cases and controls, likely resulting in selection bias that distorts the conclusion. The authors note that almost one fifth of the cases and one third of the controls refused to participate. The baseline cardiovascular characteristics, while not significantly different (with the exception of MI) between cases and controls, were consistently more prevalent in the NAION group. This finding is not surprising given that these cardiovascular conditions, especially in combination, are also risk factors for NAION.³ Thus, the MI and HTN subgroup analyses presented in table 3 should be interpreted with scepticism.

Exacerbating the inherent problems are subgroup analyses that had no a priori hypothesis. The dangers of unplanned subgroup analyses in research are well documented.⁴ Compounding matters is the sparse number of patients, reflected in the exceptionally wide confidence intervals (table 3). The robustness of such extremely small cell numbers must also be questioned, as the observed statistical significance for patients with MI can be eliminated if only one or two patients are switched to an alternative category. The authors also did not provide individual patient totals by exposure group with and without MI, rendering it impossible to replicate their results. Furthermore, there appear to be errors in the numbers/percentages and crude odds ratios presented in table 2.

In summary, the methodological limitations call into serious question the authors' conclusions. For men with a history of MI or HTN, therefore, this study does not provide any valid evidence that the use of Viagra or Cialis may increase the risk of NAION.

R E Sobel

Pfizer Inc, Worldwide Development, 150 E 42nd Street, MS#150-3-72, New York, NY 10017, USA

J C Cappelleri

Pfizer Inc, Global Research and Development, Groton, CT, USA

Correspondence to: Rachel E Sobel, Pfizer Inc, Worldwide Development, 150 E 42nd Street, MS#150-3-72, New York, NY 10017, USA; rachel.sobel@pfizer.com

doi: 10.1136/bjo.2006.094375

Accepted for publication 9 March 2006

References

- 1 McGwin G, Vaphiades MS, Hall TA, *et al*. Non-arteritic anterior ischaemic optic neuropathy and the treatment of erectile dysfunction. *Br J Ophthalmol* 2006;**90**:154–7.
- 2 Rothman KJ, Greenland S. *Modern epidemiology*. Philadelphia: Lippincott-Raven, 1998.
- 3 Hatzichristou D. Phosphodiesterase 5 inhibitors and nonarteritic anterior ischemic optic neuropathy (NAION): coincidence or causality? *J Sex Med* 2005;**2**:751–8.
- 4 Assman SF, Pocock SJ, Enos LE, *et al*. Subgroup analysis and other (mis)uses of baseline data in clinical trials. *Lancet* 2000;**355**:1064–9.

Surgical characteristics of the retinal inner limiting membrane

In a recent issue of the *BJO* Steven *et al* describe secondary retinal holes after removal of the inner limiting membrane (ILM).¹ We

consider this a key retinal publication and therefore wish to strengthen the importance of the authors' findings for modern surgery of the vitreoretinal interface.

Interestingly, the paper included only patients with macular puckers and cystoid macular oedema, but not macular holes. In cases of macular holes only a small sized central part of the ILM will usually be peeled. In contrast, macular puckers and cystoid oedemas will typically affect a larger region of the posterior pole. Thus a more extensive fraction of the ILM will be stripped if surgery is performed in such cases.

More than 30 years ago Foos² described two anatomical features of the ILM that might explain why current peeling of the peripheral ILM may be unsafe. Firstly, the ILM is thickest at the posterior pole and will become continuously thinner in the periphery. In the initiation of the ILM removal this may increase the risk for direct retinal trauma. Secondly, the Müller glia are more strongly attached to the ILM away from the macula. This may lead to indirect surgical trauma of the non-neuronal parts of the retina during the peeling process.

In conclusion, the more eccentric from the macula, the thinner and the more adherent to the retina is the ILM. As described by Steven *et al*, removal of these peripheral ILM fractions may lead to retinal hole formation.¹ Direct and/or indirect surgical trauma may be responsible for such injury, which may not be related to toxic effect of dyes used. One should therefore avoid stripping the peripheral ILM outside the macular region to prevent inadvertent retinal damage.

T Theelen, M A D Tilanus

Radboud University Nijmegen Medical Centre, Nijmegen, Netherlands

Correspondence to: Dr Thomas Theelen, Radboud University Nijmegen Medical Centre, Philips van Leydenlaan 15, Nijmegen, Netherlands; t.theelen@ohk.umcn.nl

doi: 10.1136/bjo.2006.095950

Accepted for publication 4 April 2006

References

- 1 Steven P, Laqua H, Wong D, *et al*. Secondary paracentral retinal holes following internal limiting membrane removal. *Br J Ophthalmol* 2006;**90**:293–5.
- 2 Foos RY. Vitreoretinal juncture; topographical variations. *Invest Ophthalmol* 1972;**11**:801–8.

Lutein and zeaxanthin dietary intake and age related macular degeneration

We read with interest the letter published by Vu *et al*¹ which investigated the risk of age related macular degeneration (AMD) and its association with the dietary carotenoids, lutein and zeaxanthin (LZ), stratified by linoleic acid intake. Vu *et al* reported a marked increase in the risk of both early and late AMD among people who consumed greater than the median intake of linoleic acid and higher dietary intakes of LZ.

We have a number of concerns in relation to the authors' letter and their conclusions. The letter used cross sectional data based on photographic macular assessments of 71.9% of their sample of 2448 people, who attended follow up examinations. The authors also included 212 people who did not have