vaccination and corneal transplant rejection. If patients have symptoms of corneal transplant rejection, including blurred vision, injected eye and discomfort, they should contact their general practitioner or ophthalmologist immediately. Rejection can take place up to 2 months after influenza vaccination and is potentially reversible. However each rejection episode leads to the inexorable loss of endothelial cells and potentially irreversible graft failure. Owing to the increased publicity of potential influenza pandemics and the increased uptake of vaccination, we think that it is important that this causal relation, between influenza vaccination and corneal transplant rejection, be recognised and publicised.

M S Wertheim

Bristol Eye Hospital, Lower Maudlin Street,
Bristol BS1 2LX, UK

M Keel

Pembroke Road Surgery, 111 Pembroke Road,
Bristol BS8 3EU, UK

S D Cook, D M Tole

Bristol Eye Hospital, Lower Maudlin Street,
Bristol BS1 2LX, UK

Correspondence to: Mr Michael S Wertheim, Bristol Eye Hospital, Lower Maudlin Street, Bristol BS1 2LX, UK; drwertheim@hotmail.com

doi: 10.1136/bjo.2006.093187

Accepted for publication 9 March 2006

References

- 1 **UK Transplant.** *Transplant activity in the UK 2004–2005.* Bristol: UK Transplant, 2005:36–40.
- 2 **The Collaborative Corneal Transplantation Studies Research Group.** The collaborative corneal transplantation studies (CCTS). Effectiveness of histocompatibility matching in high-risk corneal transplantation. *Arch Ophthalmol* 1992;11:1392–403.
- 3 **Streilein JW.** New thoughts on the immunology of corneal transplantation. *Eye* 2003;17:943–8.
- 4 **Duchini A, Goss JA, Karpen S, et al.** Vaccinations for adult solid-organ transplant recipients: current recommendations and protocols. *Clin Microbiol Rev* 2003;16:357–64.
- 5 **Department of Health.** *The influenza immunisation programme.* London: DoH, 2005.
- 6 **Solomon A, Frucht-Pery J.** Bilateral simultaneous corneal graft rejection after influenza vaccination. *Am J Ophthalmol* 1996;121:708–9.
- 7 **Steinemann TL, Koffler BH, Jennings CD.** Corneal allograft rejection following immunization. *Am J Ophthalmol* 1988;106:575–8.

MAILBOX

Eyelid basal cell carcinoma

We read with interest the paper by Hamada *et al.*¹ which draws a number of conclusions from a 5 year follow up study of 69 periocular basal cell carcinomas (BCCs) treated by conventional surgery and, in particular, suggests that there is no place for Mohs micrographic surgery (MMS) in patients with periocular BCCs. MMS is the serial saucerisation excision with mapped horizontal tissue sections examining 100% of the surgical margins to produce histological evidence of tumour negative margins. Unfortunately, the data included in the paper are incomplete and if such conclusions are to be considered, then further clarification is required.

Risk of BCC recurrence relates directly to the nature of the tumours treated.² The

principal risk factors for recurrence include previous treatment, large tumour size, and an infiltrative or micronodular histological growth pattern. No information is given on the first two factors and the histological subtype was non-specified in approximately 45% of cases. We calculate from the data provided that the authors experienced a 19% 5 year recurrence rate in patients with a histologically infiltrative BCC.

If most of the “non-specified” tumours in Hamada’s series were small nodular tumours, as the paper implies, then Hamada’s series also differs significantly from other larger series in that it represents a group of patients with an inherently better prognosis. Other comments hint at this, in that 76% of BCCs were on the lower eyelid and 72% were amenable to direct closure. If the majority of the tumours in this series were not in a high risk group then only one recurrence would be anticipated. In contrast, the Australian Mohs’ database series³ reported on a much higher incidence of high risk tumours (50% were infiltrative, morphoeic, basosquamous, or superficial), of which only 54% were on the lower eyelid while 41% affected the medial canthus, a site with a proved higher risk of recurrence. Despite this, they reported a 0% recurrence rate for primary BCCs of all histological subtypes treated by MMS surgery.

Hamada *et al* also conclude that 4 mm margins are justified for well defined nodular tumours, on the basis of the 5 year recurrence rates and the fact that most eyelids can still be directly repaired after such excisions. It is of interest, however, that 16% of the patients reported had incomplete tumour excision at the first attempt, although we do not know what margins were used for this group. As conventional pathological sectioning examines less than 1% of the margins,⁴ it is likely that the actual incomplete excision rate was higher.

Hamada *et al* argue that MMS is not necessary for periocular tumours on the basis that it is difficult to obtain true MMS sections. Unless either periosteum or anterior orbital septum are breached by the tumour, then there is no evidence for the former statement. What about cost benefit? MMS histology costs are greater than routine sections. However, when the other advantages of MMS are taken into account MMS is cost effective in comparison with traditional surgical excision.⁵

The current best evidence shows that recurrence rates are lower with MMS than with any other technique, with MMS for recurrent or high risk tumours showing the greatest advantage over conventional surgery. Furthermore, Hamada *et al* acknowledge that the tissue sparing quality of MMS is an important issue in that 36% of patients will develop a second BCC within 5 years.

Although we believe that MMS is the treatment of choice for optimal cure rates in periocular BCC, we would agree with Hamada *et al* that it is currently impractical for all tumours because of the patchy availability of the service in the United Kingdom. However we believe the current evidence shows that MMS remains the optimal treatment for all high risk tumours based on treatment status, site, size, and histological subgroup and do not feel that the data presented warrant a different conclusion.

E A Barnes, A J Dickinson

Department of Ophthalmology, Royal Victoria Infirmary, Newcastle upon Tyne NE1 4LP, UK

J A A Langtry, C M Lawrence

Department of Dermatology, Royal Victoria Infirmary, Newcastle upon Tyne NE1 4LP, UK

Correspondence to: Mr Eric A Barnes, Royal Victoria Infirmary, Claremont Wing, Victoria Road, Newcastle upon Tyne NE1 4LP, UK; eric.barnes@nuth.nhs.uk

doi: 10.1136/bjo.2006.092429

Accepted for publication 9 March 2006

Competing interests: none to declare.

References

- 1 **Hamada S, Kersey T, Thaller VT.** Eyelid basal cell carcinoma: non-Mohs excision, repair and outcome. *Br J Ophthalmol* 2005;89:992–4.
- 2 **Lawrence CM.** Mohs surgery—a critical review. *Br J Plast Surg* 1993;46:599–606.
- 3 **Malhotra R, Huilgol SC, Huynh NT, et al.** The Australian Mohs Database, part II: Periocular basal cell carcinoma outcome at 5 year follow up. *Ophthalmol* 2004;111:631–6.
- 4 **Abide JM, Nahai F, Benner RG.** The meaning of surgical margins. *Plast Reconstr Surg* 1984;73:492–6.
- 5 **Cook J, Zitelli JA.** Mohs micrographic surgery: a cost analysis. *J Am Acad Dermatol* 1998;39:698–703.

Authors’ reply

We thank Barnes *et al* for their interest in our study and for affording us the opportunity to correct some misinterpretations that others may also have made regarding our paper.¹

We agree entirely that histological examination of the complete excision surface as suggested by Mohs would be the ideal one should aim at and that it would be superior to conventional bread loaf histological sectioning. We dispute that this is realistically possible in inhomogeneous and exceedingly mobile eyelid tissues unless they are fixed before excision as in the chemosurgery that Mohs originally described.

Trying to cut a complete 2 mm thick saucer of tissue through fresh skin, orbicularis, tarsal plate, orbital septum, and fat and then transferring it without distorting the relative tissue orientation is a challenge that few are equal to. It might have been helpful if Barnes *et al* had shared their practical experience of the technique with us in their letter.

That the risk of basal cell carcinoma (BCC) recurrence relates to the tumour type is not in dispute and is indeed borne out in our results. But it is helpful of Barnes *et al* to remind other readers of this fact. Histological subtype was not specified in the greater portion of our series as our histopathologists only pass comment when BCCs are infiltrative or unusual in other respects. We apologise for not stating this explicitly as it has clearly led to some misunderstanding.

However, there is no justification for Barnes *et al*’s assumption that the “non-specified” tumours were small because 72% were managed by direct closure. We have clearly failed to get over a most important point. Direct closure under tension is not only possible in much larger lid defects than conventionally taught but also gives far superior functional and cosmetic results. As such, we urge others to use this method of repair more widely.

That 76% of our tumours occurred on the lower lid should not come as a surprise to anyone as it is a well recognised fact. It does not suggest that our series is in any way biased.

That our series differs from the selected series in the Australian Mohs database is self evident. We described our experience with an