symptoms. The bilateral visual acuity was 20/20 and no remarkable scotoma was observed by microperimetry. There were several Lisch nodules bilaterally. No remarkable fundus abnormalities were observed by ophthalmoscopic and biomicroscopic fundus examinations or fluorescein angiography. The transverse retinal images obtained using 3-D OCT showed small, round, isolated, high intensity lesions near the optic disc (fig 1A); the longitudinal OCT images showed that the lesions were in the retinal nerve fibre layer (fig 1B). Areas of multiple, bright patches were observed non-invasively with SLO infrared imaging (fig 2A) using the direct confocal mode. Two regions of bright and dark patches using the indirect mode (fig 2B) were observed. Only the dark patches corresponded to the hypofluorescent area observed by indocyanine green angiography (fig 2C).

Comment

We detected small, round, isolated high intensity retinal lesions on the transverse images using 3-D OCT. We also evaluated the retinal layer in which the lesions were located by longitudinal imaging and found that the lesions were in the retinal nerve fibre layer. The retinal lesions were undetected by conventional means—that is, funduscopy or fluorescein angiography, because they were too small and had no neovascularisation. Therefore, these occult lesions could hardly be detected by longitudinal imaging without the availability of transverse images. Astrocytic hamartomas had been reported as a rear retinal manifestation of neurofibromatosis.³ Because astrocytes are distributed mainly within 5–6 mm of the optic disc, astrocytic hamartomas have been reported close to the disc and in the nerve fibre layer or ganglion cell layer. Therefore, the retinal lesions that could be detected using 3D OCT might be astrocytic hamartomas. Our findings suggested that retinal lesions might not be as rare as previously thought but rather are undetectable by conventional means.

Figure 1 (A) Transverse OCT image and (B) longitudinal OCT image. The transverse OCT image shows the area of the lesion (arrow), and the longitudinal OCT image shows that the lesion is in the retinal nerve fibre layer (arrowhead).

In the present case, multiple, bright patches were observed with infrared imaging using the SLO direct confocal mode, the same as have been reported. Because no remarkable abnormalities were observed by fluorescein angiography, the lesions would not be caused by retinal or choriocapillaris circulation failure. Furthermore, a bright reflex using the SLO with a confocal aperture indicated the presence of highly reflective tissue. Yasunari *et al*⁵ reported this to be choroidal neurofibromas or choroidal nodules because of histological observations. These bright patches showed two types of lesions using the indirect mode. When using the indirect mode, the ring aperture blocks the reflex from the focal plane and allows imaging of deep structures. The finding that only the dark patches corresponded to the hypofluorescent area might be because the choroidal region close to the retina blocks fluorescence from the underlying choroidal vessels and the deeper region does not. Therefore, these differences might indicate the depth of the lesions in the choroid. SLO infrared imaging using the indirect mode demonstrates choroidal lesions and may provide information about their depth.

3-D OCT and SLO facilitate non-invasive analysis of retinal and choroidal pathologies in eyes with neurofibromatosis type 1. The advanced technology enables us to observe structures that were previously undetectable using conventional methods. Retinal and choroidal lesions in eyes with neurofibromatous 1 might not be as rare as previously thought.

Figure 2 (A) Confocal SLO infrared image, (B) indirect SLO infrared image, and (C) indocyanine green angiography. (A) Several bright patches, (B) bright patch (arrowhead), and dark patch (arrow). (C) The area of hypofluorescence (arrow) is only observed corresponding to the dark patch in the indirect SLO infrared image.

S Ishiko, A Yoshida, Y Kato, H Kagokawa
Department of Ophthalmology, Asahikawa Medical College, Asahikawa, Japan

Correspondence to: Satoshi Ishiko, MD, Department of Ophthalmology, Asahikawa Medical College, 2-1 Midorigaoka Higashi, Asahikawa, 078-8510 Japan; ishiko@asahikawa-med.ac.jp

doi: 10.1136/bjo.2006.092718

Accepted for publication 28 March 2006

References

- 1 To KW, Rabinowitz SM, Friedman AH, *et al*. Neurofibromatosis and neural crest neoplasms: primary acquired melanosis and malignant melanoma of the conjunctiva. *Surv Ophthalmol* 1989;**33**:373–9.
- 2 Warwar RE, Bullock JD, Shields JA, *et al*. Coexistence of 3 tumors of neural crest origin: neurofibroma, meningioma, and uveal malignant melanoma. *Arch Ophthalmol* 1998;**116**:1241–3.
- 3 Destro M, D'Amico DJ, Gragoudas ES, *et al*. Retinal manifestations of neurofibromatosis. *Arch Ophthalmol* 1991;**109**:662–6.
- 4 Huson S, Jones D, Beck L. Ophthalmic manifestations of neurofibromatosis. *Br J Ophthalmol* 1987;**71**:235–8.
- 5 Yasunari T, Shiraki K, Hattori H, *et al*. Frequency of choroidal abnormalities in neurofibromatosis type 1. *Lancet* 2000;**356**:988–92.
- 6 Puliafito CA, Hee MR, Lin CP, *et al*. Imaging of macular diseases with optical coherence tomography. *Ophthalmology* 1995;**102**:217–29.
- 7 Podoleanu AG, Dobre GM, Cucu RG, *et al*. Combined multipolar optical coherence tomography and confocal scanning ophthalmoscopy. *J Biomed Opt* 2004;**9**:86–93.
- 8 Yannuzzi LA, Ober MD, Slakter JS, *et al*. Ophthalmic fundus imaging: today and beyond. *Am J Ophthalmol* 2004;**137**:511–24.
- 9 Minami Y, Ishiko S, Takai Y, *et al*. Retinal changes in juvenile X linked retinoschisis using three dimensional optical coherence tomography. *Br J Ophthalmol* 2005;**89**:1663–4.

Disseminated keratoconjunctival epitheliopathy in a patient with systemic lupus erythematosus

Conjunctivocorneal involvement in systemic lupus erythematosus (SLE) is quite infrequent.^{1–3} We report the impression cytological (IC), immunohistochemical (IHC), and transmission electron microscopic (TEM) features of a case with unusual keratoconjunctival epitheliopathy refractory to medical treatment.

Case report

A 76 year old woman was referred with a long history of bilateral foreign body sensation. She had been on topical antibiotics, artificial tears, corticosteroids, hyaluronate, and aciclovir without improvement. At initial examination, biomicroscopy revealed diffuse, translucent, disseminated peripheral corneal intraepithelial coin lesions with a cystic appearance (fig 1A, B), also in the bulbar conjunctiva over the limbus (fig 1C). Fluorescein staining revealed increased permeability with negative staining. Schirmer's test was normal in both eyes. The patient had systemic and dermatological features diagnosed as SLE and had been taking prednisolone 7.5 mg/day for 1 year. Conjunctival cultures were negative for infectious processes. A thorough systemic evaluation for other systemic diseases and serum antibodies for herpes simplex virus (HSV-1, HSV-2), varicella zoster virus (VZV), and Epstein-Barr virus (EBV) were negative.

Figure 1 Biomicroscopic features of ocular surface in a patient with SLE. (A) Fluorescein staining revealed disseminated, coin-like epithelial lesions predominantly in the peripheral cornea and bulbar conjunctiva at the initial examination. (B) Retroillumination observation revealed intraepithelial cystic appearances of the lesions. (C) The lesions observed in both the cornea and bulbar conjunctiva cross over the limbus with no apparent differences in morphology. (D) Coin lesions changing to a different location over time. (E) The epitheliopathy attained a vortex phenotype following therapeutic contact lens wear.

Figure 2 Histopathological features of conjunctival specimens obtained from a patient with SLE. (A) IC revealing inflammatory cells over and in between the epithelium. Magnification, $\times 200$. (B) Haematoxylin and eosin staining also showed conjunctival epithelial thickening, cystic spaces, and numerous mononuclear cells near the basement membrane and in the stroma. Magnification, $\times 400$. (C) Immunohistochemistry showed IgG depositions beneath basal epithelial layers and perivascularly in the stroma. Epithelial thickening, cystic spaces and numerous mononuclear cells near the basement membrane and in the stroma were also shown. Magnification, $\times 400$. (D) Immunohistochemistry also showed depositions of C3 beneath basal epithelial layers and perivascularly in the stroma. Magnification, $\times 400$. (E) Representative isotype control specimen showing lack of stain for IgG deposition. Magnification, $\times 400$. (F) Conjunctival transmission electron microscopy from a coin lesion site showed decreased cellular cohesion, necrotic cells, and disorganised basement membrane. Magnification, $\times 2500$.

The patient was managed with non-preserved artificial tears, topical corticosteroids and therapeutic lenses with temporary resolution, but on recurrence of the lesions these changed their distribution and shape (fig 1D) attaining vortex morphology (fig 1E). IC revealed inflammatory cells over and in between the epithelium (fig 2A). Haematoxylin and eosin staining also showed conjunctival epithelial thickening, cystic spaces, and numerous mononuclear cells near the basement membrane and in the

stroma (fig 2B). IHC of the conjunctival limbal biopsy showed positive staining for immunoglobulin (IgG, IgA) or complement (C3 and C4) deposition beneath basal epithelial layers and perivascularly (figs 2C, D). Isotype control specimens did not reveal any staining for all markers (fig 2E). Conjunctival TEM from a coin lesion showed decreased cellular cohesion, necrotic cells, and disorganised basement membrane (fig 2F). The epitheliopathy persisted for the next year without complete resolution to date.

Comment

The unusual epitheliopathy of this patient has several different characteristics including (1) diffuse keratoconjunctival involvement, (2) persistence with migratory changes in shape and localisation, (3) association with conjunctival inflammation without involvement of the anterior chamber or posterior segment, (4) unresponsiveness to conventional medical treatments including topical corticosteroids. While histopathological features accorded in part with previous reports,^{4,5} a PubMed review revealed no report describing migrating coin lesions combining to form a vortex keratopathy in SLE. Although corneal staining is common in secondary Sjögren's syndrome,⁶ Schirmer's test was normal in our case. Simultaneous conjunctivocorneal lesions suggested infectious, toxic, or immunological aetiologies including viral infections and Thygeson's superior punctate epithelitis. However, repeated laboratory tests were negative for infectious processes, and topical corticosteroids were not effective. It is also unlikely that viral infection of the epithelium could stay active for many years.

Although the exact pathogenesis is unclear, immunological and inflammatory reactions related to SLE may explain the interesting epitheliopathy. Indeed, IC and biopsies revealed an inflammatory process that may have resulted in weakening of cellular cohesion, increased cellular death, and epithelial turnover manifesting with epithelial thickening, cystic spaces corresponding to necrotic cells or areas of increased cellular exfoliation viewed as coin lesions. TEM showed decreased intercellular cohesion and disorganisation of basement membrane, which in part may be responsible for the epitheliopathy. In summary, we reported new clinical conjunctivocorneal features of an unusual disseminated epitheliopathy in SLE which was associated with histopathological evidence of inflammatory and immunological reactions.

S Den, M Dogru

Department of Ophthalmology, Tokyo Dental College, 5-11-13 Sugano, Ichikawa, Chiba, 272-8513, Japan

M Dogru, N Kato

Department of Ophthalmology, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, Tokyo, 160-8582, Japan

Y Tanaka

Clinical Laboratory, Tokyo Dental College, 5-11-13 Sugano, Ichikawa, Chiba, 272-8513, Japan

S Shimmura

Department of Ophthalmology, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku and Department of Ophthalmology, Tokyo Dental College, Tokyo, 160-8582, Japan

J Shimazaki

Department of Ophthalmology, Tokyo Dental College, 5-11-13 Sugano, Ichikawa, Chiba, 272-8513, and Department of Ophthalmology, Keio University School of Medicine, Japan

Correspondence to: Seika Den, MD, Department of Ophthalmology, Tokyo Dental College, 5-11-13 Sugano, Ichikawa, Chiba, 272-8513, Japan; seikatdc@u01.gate01.com

doi: 10.1136/bjo.2006.090498

Accepted for publication 18 March 2006

The authors have no proprietary interests in any products mentioned in this manuscript.

References

- 1 Gold DH, Morris DA, Henkind P. Ocular findings in systemic lupus erythematosus. *Br J Ophthalmol* 1972;**56**:800-4.
- 2 Huey C, Jakobiec FA, Iwamoto T, et al. Discoid lupus erythematosus of the eyelids. *Ophthalmology* 1983;**90**:1389-98.
- 3 Borruat FX. Neuro-ophthalmologic manifestations of rheumatologic and associated disorders. *Curr Opin Ophthalmol* 1996;**7**:10-18.
- 4 Heiligenhaus A, Dutt JE, Foster CS. Histology and immunopathology of systemic lupus erythematosus affecting the conjunctiva. *Eye* 1996;**10**(Pt 4):425-32.
- 5 Karpik AG, Schwartz MM, Dickey LE, et al. Ocular immune reactants in patients dying with systemic lupus erythematosus. *Clin Immunol Immunopathol* 1985;**35**:295-312.
- 6 Hochberg MC, Boyd RE, Ahearn JM, et al. Systemic lupus erythematosus: a review of clinical-laboratory features and immunogenetic markers in 150 patients with emphasis on demographic subsets. *Medicine (Baltimore)* 1985;**64**:285-95.

A denileukin diftotox (Ontak) associated retinopathy?

Two patients with haematological malignancies presented with visual symptoms during therapy with denileukin diftotox. Both experienced loss of central and colour vision, coarse pigmentary macular changes, and decreased electroretinographic amplitudes. These cases raise the possibility of a novel drug associated retinopathy.

Case 1

A 58 year old man with chronic lymphocytic leukaemia was enrolled in a clinical trial of denileukin diftotox (Ontak, Ligand Pharmaceuticals) receiving 10 standard cycles of therapy with good response.¹ Eight months later his disease progressed and he received four further cycles. During this treatment he noted blurred vision, photopsias and difficulty adjusting to bright lights. Acuity was 6/12 bilaterally and coarse macular pigment changes were present (fig 1). Automated fields showed paracentral scotomas and electroretinograms (ERG) revealed delayed and diminished photopic wave forms with preserved scotopic responses. Brain magnetic resonance imaging (MRI) and cerebrospinal fluid (CSF) were normal.

After 1 month of visual symptoms high serum levels of retinal α -enolase antibodies were detected. Antibodies to recoverin were absent. Denileukin diftotox was stopped, he received plasmapheresis, pooled human immunoglobulin, and high dose prednisolone. There was no visual improvement though anti-enolase testing became negative. Nine months later he was diagnosed with metastatic adenocarcinoma of unknown primary from which he died 2 months later.

Case 2

A 62 year old man with relapsed primary cutaneous T cell lymphoma (CTCL) was enrolled into a trial of denileukin diftotox and received four cycles.² His lymphoma responded to treatment; however, he noted gradual blurring of vision. Visual acuity was 6/18 bilaterally and coarse macular pigmentary changes were present (fig 2). Brain MRI and CSF were normal. Electrophysiology revealed reduced photopic and scotopic amplitudes. Retinal antibodies against 30 kDa, 35 kDa, and 46 kDa (enolase) retinal proteins were detected. He died 2 months later from liver failure.

Figure 1 Colour fundus photographs and mid-phase fluorescein angiogram reveals marked macular pigmentary change in patient 1.

Comment

Denileukin diftotox is a fusion protein of portions of diphtheria toxin and interleukin 2. It targets IL-2 receptors on malignant cells and induces cell death. It is approved for the treatment of CTCL and trials in other malignancies continue. Both patients experienced visual symptoms concurrent with their denileukin diftotox therapy and were receiving no other chemotherapy agents. They had strikingly similar macular changes, ERG loss, and anti-enolase in their sera.

There exists a theoretical link between denileukin diftotox and autoimmune retinal disease. A subset of regulatory T cells crucial in suppressing autoimmune disease also express

the IL-2 receptor subunit known as CD25.³ In a proportion of mice depleted of this regulatory subset of T cells, loss of self tolerance to retinal antigens has been reported.⁴

We considered other possible causes of these retinal changes including malignant infiltration and paraneoplastic retinopathy. Infiltration was excluded on the clinical fundus appearance, and the absence of progressive disease in CSF. Both fit the phenotype of cone loss that Weleber found in seven of 12 patients with presumed anti-enolase retinopathy,⁵ but our patients' macular changes are distinctive. In the three reported cases of paraneoplastic retinopathy associated with haematological malignancy,⁶⁻⁸

Figure 2 Red free photographs and mid-phase fluorescein angiogram of patient 2.