We tested two red snapper fish of the type he normally consumed and found tissue mercury concentrations of 0.47 and 0.66 µg/g. An average serving of fish is about 150 g. This suggests his average weekly mercury intake to be between 848 and 1017 µg. This is much higher than US FDA guidelines, which recommend that an average person should consume no more than 42 µg of mercury/week.

Electroretinography (ERG) carried out on our patient (table 1) indicates a bilateral optic neuropathy or bilateral cerebral visual pathway damage as the main cause of the reduced acuity. We also found diminished colour sensitivity, especially along the protan axis, as previously reported. These findings are reported as being "consistent with damage to the optic nerves, and possibly the occipital cortex, as has been reported in mercury poisoning". We believe that these results, combined with the patient's visual field abnormalities, suggest the primary cause of his visual loss to be bilateral optic neuropathies secondary to chronic mercury poisoning.

He consulted a dietitian for dietary advice and now avoids any likely source of mercury in his diet. In humans, about half the body burden of mercury can be eliminated in 70 days if no further mercury is ingested. We continue to monitor his progress in the neuro-ophthalmology clinic.

Discussion

Chronic mercury exposure in a large population was first studied following development of unexplained neurological symptoms in people living at Minamata Bay, Japan, in 1960 (giving it the name Minamata disease).

The most common source of mercury in our diet (in the form of methylmercury) is from the consumption of certain fish caught in known polluted waters. Tissue mercury concentrations increase in fish that have a longer lifespan, and larger predatory game fish (such as swordfish, shark and king mackerel) are likely to have the highest mercury concentrations.¹

Exposure to methylmercury has been implicated in many conditions, including cardiovascular disease, peripheral neuropathies, psychiatric disorders and visual disturbance.² A "safe" blood mercury concentration for an adult male is uncertain. Our patient's blood mercury concentration was 68 nmol/l (13.9 µg/l). The US Environmental Protection Agency recommends keeping blood mercury concentration <5 µg/l. They found the mean blood mercury concentration among the US general population to be 1.3 µg/l.³ A study conducted on a group of women from a Caribbean island (Vieques) showed a median hair mercury concentration three times that of the US average.⁴

ERG carried out on our patient showed several abnormalities previously reported by Ventura *et al*⁵ in patients with a history of mercury intoxication. We used the International Society for Clinical Electrophysiology of Vision protocol for multifocal ERG and pattern ERG as in the published study. Their paper noted that central retinal function was markedly depressed and in our patient, multifocal visual evoked potential showed almost abolition of the central responses in both eyes. Multifocal ERG in our patient also showed

mild reduced amplitudes and delayed responses.

Neurological recovery after removal of the source of mercury is variable. Colour vision impairment as a result of occupational mercury exposure was noted to be reversible, with a return to normal after 12 months free from exposure.⁶ Other symptoms such as peripheral neuropathies have been noted to persist with little improvement, even up to 30 years later, after prolonged occupational exposure to inorganic mercury.⁷

In these cases, an early diagnosis is important, as continued exposure could cause further mercury accumulation and more serious neurological damage. We are uncertain as to the likelihood of visual improvement and continue to monitor his progress.

M Saldana, C E Collins, R Gale, O Backhouse
Leeds General Infirmary, Leeds, UK

Correspondence to: C E Collins, Leeds General Infirmary, 19 Devonshire Avenue, Roundhay, Leeds LS8 1AU, UK;
collinscjan@doctors.org.uk

doi: 10.1136/bjo.2006.094821

Competing interests: None declared.

References

- 1 US Environmental Protection Agency. What you need to know about mercury in fish and shellfish. EPA-823-R-04-005. March 2004.
- 2 Uchino M, Eto K, Kumamoto T, *et al*. Neurologic features of chronic Minamata disease (organic mercury poisoning) certified at autopsy. *Intern Med* 1995;**34**(8):744-7.
- 3 Weil M, Bressler J, Parsons P, *et al*. Blood mercury levels and neurobehavioral function. *JAMA*, 2005 **293**:1875-82.
- 4 Ortiz-Roque C, Lopez-Rivera. Mercury contamination in reproductive age women in a Caribbean island: Vieques. *J Epidemiol Commun Health* 2004;**58**:756-7.
- 5 Ventura DF, Costa MT, Costa MF, *et al*. Multifocal and full field electroretinogram changes associated with colour-vision loss in mercury vapour exposure. *Visual Neurosci* 2004;**21**:421-9.
- 6 A, Gobba F. Reversible colour vision loss in occupational exposure to metallic mercury. *Environ Res*, 1998 May, **77**:173-7.
- 7 Letz R, Gerr F, Cragle D, *et al*. Residual neurologic deficits 30 years after occupational exposure to elemental mercury. *Neurotoxicology* 2000;**21**:459-74.

Optical coherence tomography pseudo-macular hole appearance after photodynamic therapy

The case reports of two patients with myopia and one patient with age-related macular degeneration are presented. The patients were treated with photodynamic therapy for choroidal neovascularisation and have full-thickness retinal thinning in the area of treatment that mimics the appearance of a macular hole on optical coherence tomography (OCT). Ophthalmologists should recognise that this appearance does not represent a true macular hole when deciding how to manage these patients.

OCT distinguishes full-thickness macular holes (FTMH) from other entities such as lamellar holes.^{1,2} We present three patients treated with photodynamic therapy (PDT) for choroidal neovascular membranes (CNVM) who had full-thickness retinal thinning that mimicked an FTMH on OCT.

who had full-thickness retinal thinning that mimicked an FTMH on OCT.

Case reports

A 58-year-old woman with moderate myopia (-5.00 D) presented after four PDT sessions (spot diameters: 4.4-4.7 mm) for CNVM in the left eye; visual acuity was 20/63. Examination showed subfoveal fibrosis with adjacent atrophy, whereas OCT showed full-thickness thinning overlying an area of increased reflectivity consistent with treated CNVM (fig 1A).

A 53-year-old woman with high myopia (-17.50 D) was treated with PDT twice (spot diameters: 2.9 and 3.1 mm) for CNVM in the right eye. Vision improved from 20/200 to 20/50. Macular examination disclosed atrophy and subfoveal fibrosis, whereas OCT showed full-thickness retinal thinning overlying hyper-reflectivity at the RPE level corresponding to the treated CNVM (fig 1B).

An 89-year-old woman with age-related macular degeneration (AMD) had PDT four times (spot diameters: 4.6-5.3 mm), for CNVM in the right eye. Vision started at 20/100 and stabilised at 20/320 after treatment. Examination showed subretinal fibrosis and atrophy with minimal subretinal haemorrhage, whereas OCT showed almost full-thickness thinning (fig 1C).

For all the three patients, autofluorescence studies with the Heidelberg retina angiograph (Heidelberg Engineering, Germany) showed no autofluorescence signal,³ and fluorescein angiography showed no discrete areas of hyperfluorescence corresponding to the areas of thinning on OCT.

Comment

These three patients had on their OCT the appearance of a macular hole after PDT for CNVM secondary to myopia or AMD. Although macular holes in high myopia may be difficult to visualise on examination,⁴ we do not believe the patients had true macular holes because examination, fluorescein angiography and autofluorescence did not support this diagnosis. In addition, the OCTs lacked findings often seen with FTMH, such as adjacent intraretinal cystic changes, rounding of the retinal edges or subretinal fluid. Simultaneous CNVMs and FTMHs have previously been reported, but these FTMHs had the typical associated findings.^{5,6}

The cause of this "pseudo-macular hole" appearance is unclear. Retinal thinning on OCT has been described after PDT for CNVM.⁷ This FTMH-like configuration has not been reported, but may represent a severe form of post-PDT thinning. However, the "pseudo-macular hole" diameters were smaller than the spot diameters used. Parafoveal retinal thinning and geographic atrophy with retinal thinning have been seen on OCT in myopia⁸ and AMD,⁹ respectively, but not to the degree seen in our patients. The fovea does not appear to be selectively affected; foveal sparing allowed the first two patients to retain relatively good vision.

OCT findings must be interpreted in the context of examination and ancillary test results and not in a vacuum, as this could lead to an erroneous diagnosis of FTMH and recommendation of surgery. Clinicians should be aware that full-thickness thinning mimicking an FTMH may present on OCT in patients with AMD or myopia and a history of PDT.

Figure 1 (A) Optical coherence tomography (OCT) showing full-thickness retinal thinning overlying the hyper-reflectivity corresponding to the treated choroidal neovascular membrane (CNVM). (B) OCT showing full-thickness thinning over increased reflectivity in the retinal pigment epithelium (RPE) correlating with the treated CNVM. (C) OCT showing nearly full-thickness thinning overlying the irregular thickening in the RPE corresponding to the treated CNVM.

L Sobrin, J W Miller

Department of Ophthalmology, Harvard Medical School, Massachusetts Eye and Ear Infirmary, Boston, MA, USA

Correspondence to: J W Miller, Massachusetts Eye and Ear Infirmary, 243 Charles Street, Boston, MA 02114, USA;
Joan_Miller@meei.harvard.edu

doi: 10.1136/bjo.2006.095356

Accepted 10 June 2006

Competing interests: None declared.

References

- 1 Hee MR, Puliafito CA, Wong C, *et al.* Optical coherence tomography of macular holes. *Ophthalmology* 1995;**102**:748–56.
- 2 Jaffe GJ, Caprioli J. Optical coherence tomography to detect and manage retinal disease and glaucoma. *Am J Ophthalmol* 2004;**137**:156–9.
- 3 Ciardella AP, Lee GC, Langton K, *et al.* Autofluorescence as a novel approach to diagnosing macular holes. *Am J Ophthalmol* 2004;**137**:956–9.
- 4 Coppe AM, Ripandelli G, Parisi V, *et al.* Prevalence of asymptomatic macular holes in highly myopic eyes. *Ophthalmology* 2005;**112**:2103–9.
- 5 Recchia FM, Williams GA. Visual improvement following sequential photodynamic therapy and vitrectomy for coexisting macular hole and choroidal neovascularization. *Retina* 2003;**23**:110–2.
- 6 Elsing SH, Postel EA, Gill MK, *et al.* Management of eyes with both idiopathic macular hole and choroidal neovascularization. *Retina* 2001;**21**:613–18.
- 7 Rogers AH, Martidis A, Greenberg PB, *et al.* Optical coherence tomography findings following photodynamic therapy of choroidal neovascularization. *Am J Ophthalmol* 2002;**134**:566–76.
- 8 Lim MC, Hoh ST, Foster PJ, *et al.* Use of optical coherence tomography to assess variations in macular retinal thickness in myopia. *Invest Ophthalmol Vis Sci* 2005;**46**:974–8.
- 9 Mavroufides EC, Villate N, Rosenfeld PJ, *et al.* Age-related macular degeneration. In: Schuman JS, Puliafito CA, Fujimoto JG, eds. *Optical coherence tomography of ocular diseases*. 2nd edn. Thorofare, NJ: Slack, 2004:257–9.

Central retinal vein occlusion in Wegener's granulomatosis without retinal vasculitis

The classic type of Wegener's granulomatosis is characterised by necrotising granulomatous lesions of the upper and lower respiratory tracts, generalised focal necrotising vasculitis and glomerulonephritis, whereas the limited type has no renal involvement.¹ Ocular manifestations occur in 30–50% of

the patients,¹ with retinal involvement less frequent and varying from 1 to 13% in the literature.^{2–3} We report a case of Wegener's granulomatosis presenting with a central retinal vein occlusion (RVO) without clinical evidence of intraocular inflammation or retinal vasculitis.

Case report

A 22-year-old white man presented with a 1-week history of intermittent obscuration of

Figure 1 Fundus photography and fluorescein angiography of the patient with Wegener's granulomatosis. (A) Right eye fundus showing a hyperaemic optic disc, peripapillary retinal haemorrhage in the superotemporal sector with cotton wool spots along the edge, and dilatation and tortuosity of retinal veins. Fluorescein angiography showing (B) delayed and prolonged filling of the retinal vasculature, (C) blocked fluorescence as a result of the retinal haemorrhages and (D) mild vessel staining in a few areas in the late phases.