Comment

The pathogenesis of CSCR is not yet fully understood. It is known, however, that it affects mainly men from 20 to 45 years of age, with type-A personality, and is often triggered by emotional stress.

Our patient had indeed a type-A personality and was under professional stress. Nevertheless, we were puzzled to see CSCR in a 45-year-old woman who was not hypertensive, pregnant or under steroid treatment. Detailed questioning, however, disclosed testosterone intake. No relapse of CSCR was observed following cessation of testosterone treatment. We assume therefore that the testosterone treatment in this case may have contributed to the disease. We even postulate that a relatively high level of testosterone may be a risk factor for CSCR in general, as androgen receptors have been found in human retinal pigment epithelial cells.¹ This hypothesis would also explain the preponderance of males CSCR as well as the with age range of patients by age dependence of testosterone level.² Further, it is known that subjects with type-A personality on average have higher testosterone levels,³ which may be further increased under emotional stress.⁴

Most diseases result from the interplay of many factors. For CSCR, testosterone seems to be one of these acting factors.

M C Grieshaber

Department of Ophthalmology, University Hospital, Basle, Switzerland

J-J Staub

Division of Endocrinology, Department of Internal Medicine, University Hospital, Basle, Switzerland

J Flammer

Department of Ophthalmology, University Hospital, Basle, Switzerland

Correspondence to: Dr M C Grieshaber, Department of Ophthalmology, University Hospital Basle, Mittlere Strasse 91, PO Box, CH-4031 Basle, Switzerland; mgrieshaber@uhbs.ch

doi: 10.1136/bjo.2006.098277

Accepted 1 July 2006

Competing interests: None.

References

- 1 Rocha EM, Wickham LA, da Silveira LA, *et al.* Identification of androgen receptor protein and 5 α -reductase mRNA in human ocular tissues. *Br J Ophthalmol* 2000;**84**:76–84.
- 2 Zmuda JM, Cauley JA, Kriska A, *et al.* Longitudinal relation between endogenous testosterone and cardiovascular disease risk factors in middle-aged men. A 13-year follow-up of former Multiple Risk Factor Intervention Trial participants. *Am J Epidemiol* 1997;**146**:609–17.
- 3 Zumoff B, Rosenfeld, RS, Friedman M, *et al.* Elevated daytime urinary excretion of testosterone glucuronide in men with the type A behaviour pattern. *Psychosom Med* 1984;**46**:223–5.
- 4 King JA, Rosal MC, Ma Y, *et al.* Association of stress, hostility and plasma testosterone levels. *Neuro Endocrinol Lett* 2005;**26**:355–60.

High-speed, ultra-high-resolution optical coherence tomography of acute macular neuroretinopathy

Acute macular neuroretinopathy (AMN) is a rare, macular disorder of unknown aetiology. Patients with AMN are typically young women

who present with paracentral scotomata in one or both eyes corresponding to red wedge-shaped parafoveal lesions. The retinal location of the lesion in patients with AMN is not clear. High-speed, ultra-high-resolution optical coherence tomography (hsUHR-OCT) is an investigational research prototype instrument capable of producing cross-sectional images of the retina; it supports an axial resolution of about 3.5 μ m compared with about 10 μ m in Stratus OCT (Dublin, California, USA)¹ which enables enhanced imaging of intraretinal morphology including photoreceptor inner segments, outer segments and the external limiting membrane.² We report a patient with AMN who underwent imaging with hsUHR-OCT suggesting that the lesion in AMN is located in the outer retina.

Case report

A 51-year-old woman presented with a 10-day history of a sudden onset of a grey, oval paracentral scotoma in her right eye. Her medical history was notable for systemic hypertension; her drugs included trivoral and

ramipril. Best-corrected visual acuity was 20/25 in both eyes. She was able to precisely demarcate the paracentral scotoma on an Amsler grid. Funduscopy of the right eye showed a focal, reddish petaloid lesion superior to fixation. Fluorescein angiography and images on the Stratus OCT were unremarkable. hsUHR-OCT images showed focal depression of the external limiting membrane, inner/outer photoreceptor segment (IS/OS) junction, photoreceptors and retinal pigment epithelium. Changes in the photoreceptor outer segments (fig 1A) in the region of the petaloid lesion were also noted. The inner retina appeared normal.

After 3 months, visual acuity was 20/25 in the right eye, and the patient reported a slight reduction of the scotoma. Funduscopy showed resolution of the petaloid lesion. hsUHR-OCT showed realignment of the outer photoreceptor layer and the IS/OS junction (fig 1B).

Comments

The pathogenesis of AMN remains unknown, although an acute inflammatory process or

Figure 1 Three-dimensional, high-speed ultra-high-resolution optical coherence tomography scan of the macula. A 180-image raster scan (512 axial scans per image) provides a 6 $\times$ 6-mm pattern. The cross-sectional images are precisely registered to a fundus image created from an anterior view of all 180 images. (A) Perimacular petaloid lesion (arrowheads), corresponding to a focal reduction in the photoreceptor outer segments (arrow). (B) Marked resolution of both the petaloid lesion and outer segment morphology at 3-month follow-up. ELM, external limiting membrane; IS/OS, inner/outer segment photoreceptor junction; PR IS, photoreceptor inner segments; PR OS, photoreceptor outer segments; RPE, retinal pigment epithelium.

vascular disease associated with hypertension have both been proposed as mechanisms.^{3,4} Acute macular neuroretinopathy has also been associated with oral contraceptive use,⁵ eclampsia⁶ and heavy caffeine consumption.⁷ The resulting scotomata can persist for several years.⁵ Proven treatment is not available.

An early report suggested that the AMN lesion was located in the superficial layers of the retina.³ A later report using OCT suggested that the lesion was located in the outer retina.⁸ Studies using early receptor potential⁹ and a multifocal electroretinogram¹⁰ pointed to photoreceptor involvement.

hsUHR-OCT showed that the focal lesion in patients with AMN occurs in the outer retina, possibly at the level of the photoreceptor outer segments with distortion of the IS/OS junction. The lesion may also represent a disturbance at the outer nuclear layer, resulting in the downward depression and disruption of the underlying external limiting membrane, photoreceptors and retinal pigment epithelium. hs UHR-OCT shows that the structural changes in the outer retina in this patient with AMN may be reversible despite the persistence of visual symptoms.

B K Monson

New England Eye Center, Tufts-New England Medical Center, Boston, Massachusetts, USA

P B Greenberg

Division of Ophthalmology, Brown Medical School/Rhode Island Hospital, Providence, Rhode Island, USA

E Greenberg

New England Eye Center, Tufts-New England Medical Center, Boston, Massachusetts, USA

J G Fujimoto

Department of Electrical Engineering and Computer Science and Research Laboratory of Electronics, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

V J Srinivasan

Department of Electrical Engineering and Computer Science and Research Laboratory of Electronics, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

J S Duker

New England Eye Center, Tufts-New England Medical Center, Tufts University, Boston, USA

Correspondence to: B K Monson, New England Eye Center, Tufts-New England Medical Center, 260 Tremont Street Biewend Building, 9–11th Floor, Boston, MA 02111, USA; brayan.monson@tufts.edu

doi: 10.1136/bjo.2006.098871

Accepted 10 June 2006

Competing interests: JGF received royalties from intellectual property licensed by MIT to Carl Zeiss Meditec.

References

- Drexler W, Morgner U, Ghanta RK, et al. Ultrahigh-resolution optical coherence tomography. *Nat Med* 2001;7:502–7.
- Ko TH, Fujimoto JG, Schuman JS, et al. Comparison of ultrahigh- and standard-resolution optical coherence tomography for imaging macular pathology. *Ophthalmology* 2005;112:1922.
- Bos PJM, Deutman AF. Acute macular neuroretinopathy. *Am J Ophthalmol* 1975;80:573–84.
- O'Brien DM, Farmer SG, Kalina RE, et al. Acute macular neuroretinopathy following intravenous sympathomimetics. *Retina* 1989;9:281–6.
- Desai UR, Sudhamathi K, Natrajan S. Intravenous epinephrine and acute macular neuroretinopathy. *Arch Ophthalmol*, 111:1026–7.
- Kalina RE. Acute macular neuroretinopathy. In eds: Guyer DR, Yannuzzi LA, Chang S, et al. *Retina-Vitreous-Macula*. W B Saunders Co. 1999:593–6.
- Kerrison JB, Pollock SC, Biousse V, et al. Coffee and doughnut maculopathy: a cause of acute central ring scotomas. *Br J Ophthalmol* 2000;84:158–64.
- Feigl B, Haas A. Optical coherence tomography (OCT) in acute macular neuroretinopathy. *Acta Ophthalmol* 2000;78:714–16.
- Sieving PA, Fishman GA, Salzano T, et al. Acute macular neuroretinopathy: early receptor potentials change suggests photoreceptor pathology. *Br J Ophthalmol* 1984;68:229–34.
- Browning AC, Gupta R, Barber C, et al. The multifocal electroretinogram in acute macular neuroretinopathy. *Arch Ophthalmol* 2003;121:1506–7.
- Kiire CA, Dhillon B. The aetiology and associations of conjunctival intraepithelial carcinoma. *Br J Ophthalmol* 2006;90:109–13.
- Wabinga H, Parkin D, Wabwire-Mangen F, et al. Trends in cancer incidence in Kyadondo county, Uganda, 1960–1997. *Br J Cancer* 2000;82:1585–92.
- Ateenyi-Agaba C, Newton R. Squamous cell carcinoma of the conjunctiva: an HIV-associated cancer. *Bull R Soc Trop Med* 1999;7:3–4.
- Kestelyn P, Stevens A, Ndayambaje A, et al. HIV and conjunctival malignancies [letter]. *Lancet* 1990;336:51–2.
- Ateenyi-Agaba C. Conjunctival squamous-cell carcinoma associated with HIV infection in Kampala, Uganda. *Lancet* 1995;345:695–6.
- Waddell K, Lewallen S, Lucas S, et al. Carcinoma of the conjunctiva and HIV infection in Uganda and Malawi. *Br J Ophthalmol* 1996;80:503–38.
- Newton R, Ziegler J, Beral V, et al. A case control study of human immunodeficiency virus infection and cancer in adults and children residing in Kampala, Uganda. *Int J Cancer* 2001;92:622–7.
- Goedert JJ, Cote TR. Conjunctival malignant disease with AIDS in USA. *Lancet* 1995;ii:257–8.
- Waddell KM, Dowling RG, Lucas SB, et al. Corneoconjunctival carcinoma in Uganda. *Eye* 2006;20:893–9.
- Holkar S, Mudhar HS, Jain A, et al. Regression of invasive conjunctival squamous carcinoma in an HIV-positive patient on antiretroviral therapy. *Int J STD AIDS* 2005;16(12):782–3.
- Beral V, Newton R. Overview of the epidemiology of immunodeficiency associated cancers. *Monogr Natl Cancer Inst* 1998;23:1–6.
- Newton R. A review of the aetiology of squamous cell carcinoma of the conjunctiva. *Br J Cancer* 1996;74:1511–3.
- Newton R, Ziegler J, Ateenyi-Agaba C, et al. The epidemiology of conjunctival squamous cell carcinoma in Uganda. *Br J Cancer* 2002;87:301–8.
- Waddell K, Magyezi J, Bousarghin L, et al. Antibodies against human papillomavirus type 16 (HPV-16) and conjunctival squamous cell neoplasia in Uganda. *Br J Cancer* 2003;88:2002–3.
- Ateenyi-Agaba C, Weiderpass E, Smet A, et al. *Epidermodysplasia verruciformis*. human

for a median time of 32 months, using a simple surgical protocol applicable to resource-poor countries.⁹ They discuss treatment with topical mitomycin C and 5-fluorouracil, but these require prolonged follow-up and are not practical in much of Africa. They also call for a large controlled trial of interferon α 2b; only in Africa could a large enough series be found. Excision biopsy remains the treatment of choice for most patients, requiring only one visit, except for the rare diffuse tumours. We agree that a low threshold for advising excision of lesions is important, as tumours as small as 2–3 mm can be invasive; endophytic tumours are especially deceptive, sometimes looking like pingueculae.⁹

K M Waddell

Ruharo Eye Hospital and Uganda Eye Project, Mbarara, Uganda, Africa

R Newton

Epidemiology and Genetics Unit, Department of Health Sciences, University of York, York, UK

Correspondence to: Dr R Newton, Epidemiology and Genetics Unit, Department of Health Sciences, University of York, Seabrook Rowntree Building, Heslington, York YO10 5DD, UK; rob.newton@egu.york.ac.uk

doi: 10.1136/bjo.2006.100297

Accepted 27 June 2006

Competing interests: None.

References

- Kiire CA, Dhillon B. The aetiology and associations of conjunctival intraepithelial carcinoma. *Br J Ophthalmol* 2006;90:109–13.
- Wabinga H, Parkin D, Wabwire-Mangen F, et al. Trends in cancer incidence in Kyadondo county, Uganda, 1960–1997. *Br J Cancer* 2000;82:1585–92.
- Ateenyi-Agaba C, Newton R. Squamous cell carcinoma of the conjunctiva: an HIV-associated cancer. *Bull R Soc Trop Med* 1999;7:3–4.
- Kestelyn P, Stevens A, Ndayambaje A, et al. HIV and conjunctival malignancies [letter]. *Lancet* 1990;336:51–2.
- Ateenyi-Agaba C. Conjunctival squamous-cell carcinoma associated with HIV infection in Kampala, Uganda. *Lancet* 1995;345:695–6.
- Waddell K, Lewallen S, Lucas S, et al. Carcinoma of the conjunctiva and HIV infection in Uganda and Malawi. *Br J Ophthalmol* 1996;80:503–38.
- Newton R, Ziegler J, Beral V, et al. A case control study of human immunodeficiency virus infection and cancer in adults and children residing in Kampala, Uganda. *Int J Cancer* 2001;92:622–7.
- Goedert JJ, Cote TR. Conjunctival malignant disease with AIDS in USA. *Lancet* 1995;ii:257–8.
- Waddell KM, Dowling RG, Lucas SB, et al. Corneoconjunctival carcinoma in Uganda. *Eye* 2006;20:893–9.
- Holkar S, Mudhar HS, Jain A, et al. Regression of invasive conjunctival squamous carcinoma in an HIV-positive patient on antiretroviral therapy. *Int J STD AIDS* 2005;16(12):782–3.
- Beral V, Newton R. Overview of the epidemiology of immunodeficiency associated cancers. *Monogr Natl Cancer Inst* 1998;23:1–6.
- Newton R. A review of the aetiology of squamous cell carcinoma of the conjunctiva. *Br J Cancer* 1996;74:1511–3.
- Newton R, Ziegler J, Ateenyi-Agaba C, et al. The epidemiology of conjunctival squamous cell carcinoma in Uganda. *Br J Cancer* 2002;87:301–8.
- Waddell K, Magyezi J, Bousarghin L, et al. Antibodies against human papillomavirus type 16 (HPV-16) and conjunctival squamous cell neoplasia in Uganda. *Br J Cancer* 2003;88:2002–3.
- Ateenyi-Agaba C, Weiderpass E, Smet A, et al. *Epidermodysplasia verruciformis*. human