However, Vogt Koyanagi Harada syndrome follows a relapsing remitting course, often with systemic symptoms despite treatment with immune suppression, unlike sympathetic ophthalmia, which is a potentially curable disease if managed properly in acute stages. An absence of recurrence of disease process on stoppage of therapy, as well as the clinical history of our patient, makes the diagnosis of sympathetic ophthalmia more likely.

Nadeem Ahmad

Ophthalmology Department, Royal Hallamshire Hospital, Sheffield, UK

Terrence K Soong, Sachin Salvi, Paul A Rudle, Ian G Rennie
Royal Hallamshire Hospital, Sheffield, UK

Correspondence to: Mr N Ahmad, Ophthalmology Department, A Floor, Royal Hallamshire Hospital, Glossop Road, Sheffield S10 2JF, UK; ophthdoc@tesco.net

Informed consent was obtained for publication of the person's details in this report.

doi: 10.1136/bjo.2006.102384

Accepted 7 July 2006

Competing interests: None.

References

- Kilmartin DJ, Dick AD, Forrester JV. Prospective surveillance of sympathetic ophthalmia in the UK and Republic of Ireland. *Br J Ophthalmol* 2000;**84**:259–63.
- Margo CE, Pautler SE. Granulomatous uveitis after treatment of a choroidal melanoma with proton-beam irradiation. *Retina* 1990;**10**:140–3.
- Fries PD, Char DH, Crawford JB, et al. Sympathetic ophthalmia complicating helium ion irradiation of a choroidal melanoma. *Arch Ophthalmol* 1987;**105**:1561–4.
- El-Asrar AM, Al-Obeidan SA. Sympathetic ophthalmia after complicated cataract surgery and intraocular lens implantation. *Eur J Ophthalmol* 2001;**11**:193–6.
- Sabates R. Choroiditis compatible with the histopathologic diagnosis of sympathetic ophthalmia following cyclocryotherapy of neovascular glaucoma. *Ophthalmic Surg* 1988;**19**:176–82.
- Lam S, Tessler HH, Lam BL, et al. High incidence of sympathetic ophthalmia after contact and noncontact neodymium:YAG cyclotherapy. *Ophthalmology* 1992;**99**:1818–22.
- Kumar N, Chang A, Beaumont P. Sympathetic ophthalmia following ciliary body laser cyclophotocoagulation for rubeotic glaucoma. *Clin Exp Ophthalmol* 2004;**32**:196–8.
- Bechrakis NE, Muller-Stolzenburg NW, Helbig H, et al. Sympathetic ophthalmia following laser cyclocoagulation. *Arch Ophthalmol* 1994;**112**:80–4.
- Gass JDM. Sympathetic ophthalmia following vitrectomy. *Am J Ophthalmol* 1982;**93**:552–8.
- Pollack AL, McDonald HR, Ai E, et al. Sympathetic ophthalmia associated with pars plana vitrectomy without antecedent penetrating trauma. *Retina* 2001;**21**:146–54.

Frequency of non-arteritic anterior ischaemic optic neuropathy in adult Chinese: The Beijing Eye Study

Non-arteritic anterior ischaemic optic neuropathy (NAION) is an acute optic neuropathy occurring predominantly in small optic nerve heads in elderly patients.^{1–3} It has been debated which factors, other than old age and a small optic disc size, predispose to the disease. Data

on the prevalence of NAION in the general population has so far been scarce.^{4,5} They have been completely missing for the Chinese. Therefore, the aim of this study was to evaluate the prevalence of optic nerve damage with the features of a preceding NAION in the Beijing Eye Study.

The Beijing Eye Study is a population-based cohort study in Northern China.⁶ The Medical Ethics Committee of the Beijing Tongren Hospital approved the study protocol and all participants gave their informed consent, according to the Declaration of Helsinki. Of 5324 individuals aged ≥ 40 years residing in the study area, 4439 individuals (2505 women) participated in the eye examination (response rate, 83.4%). This study included 4027 (90.7%) subjects for whom readable optic disc photographs were available. The mean (standard deviation (SD)) age was 55.2 (10) (median, 55; range, 40–101) years. The participants underwent an ophthalmic examination including photographs of the optic disc and macula (Fundus camera CR6-45NM, Canon, Lake Success, New York, USA) and frequency doubling perimetry (screening program C-20-1; Zeiss-Humphrey, Dublin, California, USA) as described in detail previously.⁶ For all eyes with visual acuity < 0.6 and for all eyes with any visual field loss, the photographs of the macula and optic disc were assessed twice by a panel that included several ophthalmologists (YW, LX, JBJ). The diagnostic criteria for NAION were a small optic disc with a cup/disc diameter ratio of ≤ 0.4 , segmental pallor, segmental loss of the retinal nerve fibre layer and segmental visual field loss.^{1–3,7}

Of the 8876 eyes included in the study, only one eye fulfilled the definition of NAION. The age of the male subject with suspected previous NAION was 71 years, his vision was 0.30, and he showed visual field defects in the inferior hemisphere. His optic disc, with a size of 2.0 mm², did not have cupping and exhibited a pale neuroretinal rim in its superior half. The visibility of the retinal nerve fibre layer was markedly reduced in the temporal superior region, and the temporal superior artery was markedly thinner than the temporal inferior artery. In a mathematical sense, the calculated mean (SD) prevalence of NAION in the whole population was 0.02 (1.6%) (95% confidence interval -0.02 to -0.07), or about 1 in 4500 subjects.

Comment

The prevalence rate of 1 in about 4500 Chinese with an age of ≥ 40 years may be at the lower end of the frequency range reported and calculated from previous studies on Caucasian populations in which about 2.3–10.3 patients per 100 000 inhabitants > 50 years were affected with NAION in the US per year.^{4,5} Despite its marked statistical limitation due to the low prevalence rate of NAION, this study may, therefore, support the finding from previous investigations that whites may be affected by NAION more commonly than population groups of another ethnic background such as Chinese.^{4,5} One may consider that the inter-ethnic differences in optic disc size, with the smallest optic discs found in Caucasians, medium-sized optic discs in Asians and largest discs in Afro-Americans, might be responsible for the interethnic differences in the frequency of NAION, which occurs predominantly in small optic nerve heads. The major limitation of this study is the low prevalence rate of NAION so that the prevalence rate for NAION as calculated in the present investigation may have to be

confirmed in even larger population-based studies with > 10 000 subjects included. One may conclude that NAION-associated optic nerve damage is present in 1 out of about 4500 adult Chinese. The figure will be around 100 000 patients with NAION-induced optic nerve damage in the whole of China.

Yaxing Wang, Liang Xu

Beijing Institute of Ophthalmology, Beijing Tongren Hospital, Capital University of Medical Science, Beijing, China

Just B Jonas

Department of Ophthalmology, Faculty of Clinical Medicine, Mannheim, University of Heidelberg, Mannheim, Germany

Correspondence to: Professor J B Jonas, Beijing Institute of Ophthalmology, Beijing Fongren Hospital, Capital University of Medical Science, 17 Hougou Street, Chong Wen Men, 100005 Beijing, China; jost.jonas@augen.ma.uni-heidelberg.de

doi: 10.1136/bjo.2006.103788

Accepted 31 July 2006

Competing interests: None declared.

References

- Hayreh SS, Joos KM, Podhajsky PA, et al. Systemic diseases associated with non-arteritic anterior ischemic optic neuropathy. *Am J Ophthalmol* 1994;**118**:766–80.
- Hayreh SS. Acute ischemic disorders of the optic nerve: pathogenesis, clinical manifestations and management. *Ophthalmol Clin North Am* 1996;**9**:407–42.
- Beck RW, Savino PJ, Repka MX, et al. Optic disc structure in anterior ischemic optic neuropathy. *Ophthalmology* 1984;**91**:1334–7.
- Johns LN, Arnold AC. Incidence of nonarteritic anterior ischemic optic neuritis (AION, population based study). *J Neuroophthalmol* 1994;**14**:38–49.
- Hattenhauer MG, Leavitt JA, Hodge DO, et al. Incidence of nonarteritic anterior ischemic optic neuropathy. *Am J Ophthalmol* 1997;**123**:103–7.
- Xu L, Cui T, Yang H, et al. Prevalence of visual impairment among adults in China. The Beijing Eye Study. *Am J Ophthalmol* 2006;**141**:591–3.
- Jonas JB, Budde WM, Panda-Jonas S. Ophthalmoscopic evaluation of the optic nerve head. *Surv Ophthalmol* 1999;**43**:293–320.

Frequency of spontaneous pulsations of the central retinal vein

The central retinal vein is the only structure in the body which can be examined non-invasively, runs through the cerebrospinal fluid space, and has a shape that depends on the relationship between its internal pressure and the pressure in the space surrounding it. Estimation of central retinal vein pressure is, therefore, helpful in the assessment of cerebrospinal fluid pressure—that is, the intracranial pressure.^{1–4} Central retinal vein pressure may be assessed by determining the external pressure at which the central retinal vein starts to pulsate. This method of assessment is similar to Riva-Rocci's method of indirect measurement of arterial blood pressure. For the central retinal vein, the external pressure is the intraocular pressure. The purpose of the present study was to find out the proportion percentage of subjects in whom the central retinal vein shows spontaneous pulsations, indicating that the pressure in the vein is lower than the intraocular pressure. The assessment of spontaneous retinal venous

pulsations may be very useful for the ophthalmic assessment whether the intracranial pressure is normal, or if spontaneous vein pulsation cannot be detected, whether the intracranial pressure is likely to be elevated.¹⁻⁴

Case report

Our clinical observational case series included 199 patients (384 eyes; mean (standard deviation (SD)) age, 63.9 (12.3) years; range, 7–90 years) attending the hospital because of cataract ($n = 66$), age-related macular degeneration ($n = 37$) or other diseases ($n = 96$) not affecting the optic nerve or primarily the inner retina. Intraocular pressure ranged between 10 and 21 mm Hg. The mean (SD) refractive error was -0.15 (3.04) D (range, -15.5 to $+6.50$ D). Using tropicamide and phenylephrine 5% eye drops, the pupil was dilated. The optic disc was examined using a slit lamp and a hand-held 78-D ophthalmoscopic lens for at least 30 s. Any small pulsatile synchronous movement of the central retinal vein or its major branches inside the optic disc was noted as spontaneous pulsations. The non-contact examination was part of a routine ophthalmoscopic examination of the fundus of the patients. The medical ethics committee approved the study protocol, fulfilling the criteria of the Declaration of Helsinki.

Spontaneous central retinal vein pulsations were detected in 343 (89.3%; 95% confidence interval (CI), 7.6 to 13.8) eyes of 186 (93.5%; 95% CI, 3.1 to 10) patients. The presence of a spontaneous central retinal vein pulsation was statistically independent of age ($p = 0.57$; 95% CI, 0.98 to 1.04) and gender ($p = 0.78$; odds ratio (OR): 1.19; 95% CI, 0.39 to 3.68). Frequency of detected spontaneous central retinal vein pulsations was significantly associated with hyperopic refractive error ($p = 0.017$; 95% CI, 1.08 to 2.19).

Comment

Confirming the findings of previous studies,^{5,6} the results of the present study (relatively high frequency of ophthalmoscopically detected spontaneous pulsations of the central retinal vein), suggest that about 10% of otherwise normal people do not have ophthalmoscopically detectable spontaneous central retinal vein pulsations. Our results as well as the previous findings may be important for several reasons: in a patient with an increased intracranial pressure, the retrobulbar cerebrospinal fluid pressure rises, eventually equalling the intraocular pressure. When this occurs, there is no longer a fluctuating intravascular pressure gradient between the intraocular retinal veins and the retrolaminar retinal vein. The venous blood flow becomes constant, and spontaneous central retinal vein pulsations cease.⁴ Based on the results of previous studies and the present investigation a lack of a spontaneous central retinal vein pulsation in a patient with a presumed increase in intracranial pressure may suggest with a probability of 10% a normal finding and with a probability of about 90% an abnormal situation. To cite an example, by Levine searched for spontaneous retinal venous pulsations in 218 subjects in a previous investigation.⁶ Spontaneous venous pulsations were present in 87.6% of 146 unselected subjects 20–90 years of age, and were absent in 100% of 33 patients with raised intracranial pressure without papilloedema and in 10 patients with papilloedema. Lumbar puncture in nine patients with raised intracranial pressure established the upper level at which spontaneous pulsations

disappear at 190 mm H₂O, and no pressure above 180 mm H₂O was found in 29 patients with venous pulsations present before lumbar puncture. Some normal subjects with no pulsations showed definite pulsations on subsequent examinations. Levine's findings show that presence of spontaneous venous pulsations is a reliable indicator of an intracranial pressure below 180–190 mm H₂O. They suggest, like the results of the present investigation, that pulsations may be absent in normal subjects and absence of pulsations is, therefore, not a reliable guide to raised intracranial pressure.

In conclusion, ophthalmoscopic assessment of spontaneous pulsation of the central retinal vein is useful for the estimation of increased intracranial pressure; in a similar manner, it may be useful for the diagnosis of glaucomatous optic neuropathy and retinal vein occlusions, in which increased central retinal vein pressure has been reported.⁷

Bjoern Harder, Jost B Jonas

Department of Ophthalmology, University of Heidelberg, Mannheim, Germany

Correspondence to: Dr J B Jonas, Universitäts-Augenklinik, Theodor-Kutzer-Ufer 1–3, 68167 Mannheim, Germany; Jost.Jonas@augen.ma.uni-heidelberg.de

doi: 10.1136/bjo.2006.103341

Accepted 24 July 2006

Competing interests: None.

References

- 1 Draeger J, Rumberger E, Hechler B. Intracranial pressure in microgravity conditions: non-invasive assessment by ophthalmodynamometry. *Aviat Space Environ Med* 1999;**70**:1227–9.
- 2 Firsching R, Schütze M, Matschmann M, et al. Venous ophthalmodynamometry: a noninvasive method for assessment of intracranial pressure. *J Neurosurg* 2000;**93**:33–6.
- 3 Jonas JB, Harder B. Ophthalmodynamometric estimation of cerebrospinal fluid pressure in pseudotumor cerebri. *Br J Ophthalmol* 2003;**87**:361–2.
- 4 Jacks AS, Miller NR. Spontaneous retinal venous pulsation: aetiology and significance. *J Neurol Neurosurg Psychiatry* 2003;**74**:7–9.
- 5 Lorentzen SE. Incidence of spontaneous venous pulsation in the retina. *Acta Ophthalmol* 1970;**48**:765–76.
- 6 Levin BE. The clinical significance of spontaneous pulsations of the retinal vein. *Arch Neurol* 1978;**35**:37–40.
- 7 Morgan WH, Hazelton ML, Azar SL, et al. Retinal venous pulsation in glaucoma and glaucoma suspects. *Ophthalmology* 2004;**111**:1489–94.

Late-stage inclusion body conjunctivitis: trachoma?

Chlamydia trachomatis is the most commonly found infectious agent in chronic conjunctivitis. Diagnosis is often delayed due to the oligosymptomatic course of the disease and the difficulty of obtaining enough infected cells in the conjunctival swab to confirm the diagnosis. These difficulties often lead to a prolonged course of the infection and possibly to scarring when infection occurs owing to chronic inflammation. We report on a 30-year-old Swiss patient having trachomatous entropion with trichiasis of the upper eyelid requiring corrective surgery, but with no history of travelling to trachoma endemic areas.

Figure 1 Cicatricial changes of the palpebral conjunctiva. Informed consent was obtained for publication of this figure.

Case report

A 34-year-old patient was seen in consultation for trichiasis involving both upper eyelids causing keratopathy. According to the patient, the development of entropion was preceded by a 1-year episode of recurrent conjunctivitis. Except for hay fever there was no other relevant ophthalmic or medical history, especially no history of possible infection elsewhere. On eversion of the eyelid, there were cicatricial changes and inflammation involving the entire palpebral conjunctiva with marked papillae (fig 1). Infection with *C trachomatis* was suspected and confirmed by a polymerase chain reaction. The patient was treated with a 2×100 mg doxycycline regimen for 2 weeks, with topical ofloxacin ointment treatment for 4 weeks. Owing to advanced entropion and trichiasis, a lid margin rotation (Trabut) procedure was performed (fig 2). The post-operative course was uneventful, with successful anatomical correction of the upper eyelids.

Comment

Chlamydiae are obligate intracellular Gram-negative eubacteria that lack cytochromes and the ability to synthesise their own ATP. *C trachomatis* has been a source of human disease especially in the eye and urogenital system for millennia, thriving in poor, crowded and unsanitary conditions. It has been known as a blinding disease for more than 3500 years.¹ Trachoma was rampant throughout Europe and North America in the 1800s, but disappeared in England before the Second World War. In the 1950s, trachoma also disappeared in Italy and the Soviet Union. This was mainly due to improved hygiene standards.

In the developing world, trachoma is still the leading infectious cause of blindness. The World Health Organisation estimates that 70

Figure 2 Lid margin rotation procedure. Informed consent was obtained for publication of this figure.