We select the letters for these pages from the rapid responses posted on bmj.com favouring those received within five days of publication of the article to which they refer. Letters are thus an early selection of rapid responses on a particular topic. Readers should consult the website for the full list of responses and any authors' replies, which usually arrive after our selection.

LETTERS

MARK MANSON/REX

STIGMATISING SEX WORKERS

BMA public health doctor's claims lack evidence

There is no epidemiological evidence to support the recent statement by Spencer Jones that 70% of all sexually transmitted infections (STIs) in Birmingham are circulating in a pool of prostitutes and their clients.¹ Surveillance data routinely collected by clinics for genitourinary medicine (GUM) and analysed by the Health Protection Agency do not include the proportion of STIs diagnosed in sex workers or their clients. Furthermore, data from the "Safe Project," a dedicated sexual health promotion service for Birmingham's sex workers, indicates that over the past year, the prevalence of STIs among the 208 sex workers attending this targeted service was low (chlamydia (1.9%), HIV (0), gonorrhoea (2.9%), and syphilis (1%)).

More recently a local enhanced surveillance programme introduced for syphilis indicated that, although commercial sex work or reported use of sex workers were identified among those found to be infected with syphilis, just 5% of the total number of syphilis diagnoses were attributable to this group.

Trends in HIV infections in Birmingham are similar to those seen nationally, and there is no evidence that contact with commercial sex workers is a significant risk factor in those newly diagnosed with HIV infection.^{2,3}

Overall the claims made by the chairman of the BMA public health committee seem to be without substance. As a public health doctor he has a duty not only to ensure that important public

health issues are discussed but also to check out the facts before going public.

Jacky Chambers director of public health
Heart of Birmingham Primary Care Trust,
Bartholomew House, Birmingham B16 9PA
Jacky.chambers@hobtpct.nhs.uk

Penny Goold consultant
Sharon Myring consultant
Department of Genitourinary Medicine, Whittall Street Clinic,
Birmingham B4 6DH

Competing interests: None declared.

- 1 Day M. BMA public health doctor is accused of stigmatising sex workers. *BMJ* 2007;334:767. (14 April.)
- 2 Health Protection Agency. *Sexually transmitted infections in residents of Birmingham and Solihull Jan-Dec 2005*. West Midlands: HPA, 2006.
- 3 The UK Collaborative Group for HIV and STI Surveillance. *A complex picture. HIV and other sexually transmitted infections in the United Kingdom: 2006*. London: Health Protection Agency Centre for Infections, 2006. www.hpa.org.uk/publications/2006/hiv_sti_2006/pdf/a_complex_picture_

For debate: Should prostitution be legalised and regulated?

The BMA's annual public health conference is an opportunity for doctors to air public health issues and debate them, to challenge each other and establish what is known and what is not known.¹ The specialty of public health takes evidence based medicine at least as seriously as other specialties, but from time to time it is important to free oneself from its fetters.

On 29 March a motion was put forward from the West Midlands region (that I drew up): "That this conference believes that the key to prevention of sexually transmitted disease is two-fold: (i) school based education that ensures that all children understand the risks associated with unprotected sexual intercourse before they become sexually active, and (ii) the legalisation and regulation of prostitution." The first part of the motion was accepted by the conference. The second part was taken "as reference": a polite way of saying, go back and work on this more.

The BMA press office issued a press release ahead of the conference to excite press interest.¹ I believe it included reference to part (ii) of the motion, which I would not have endorsed. A news agency reported that I had said that 70% of sexually transmitted infections (STIs) are in people working in the pay for sex sector,

when in fact I said that 70% of frequent fliers in STI clinics come from this sector.

My reason for submitting the motion seems to have served its primary purpose: to open up some debate. I urge colleagues to consider that it is as important to think about the "big picture" of what is going on in our society as well as what is going on in any specialty.

Certainly that is what we believe we have to do in public health. The question is, should prostitution be legalised and regulated? It is an honest question.

C J Spencer Jones chair
Committee for Public Health Medicine and Community Health (CPHMCH), BMA House, London WC1H 9JP
dalgie@bma.org.uk

Competing interests: None declared.

- 1 Day M. BMA public health doctor is accused of stigmatising sex workers. *BMJ* 2007;334:767. (14 April.)

Public health policy must be based on sound evidence

Discussions around regulatory frameworks in commercial sex require careful use of terminology. A widely circulated report of the BMA's annual public health conference seems to be advocating the legalisation and regulation of sex work in order to submit prostitutes to regular testing.^{1,2} This approach is not only unjustified, and an affront to human dignity and rights, but an inappropriate diversion of scarce NHS resources. Furthermore, it would be ineffective and dangerous.³ Coercion of sex workers merely drives them further underground and alienates them from the services they need, leading to a breakdown in sexual health practices, and an increase in sexually transmitted infections.⁴

The major health problems among sex workers are related to stigmatisation, to which this report contributes further.¹ Inaccurate and inflammatory statements such as these reported comments are likely to lead to increased levels of violence against them, as seen in Ipswich, and will even place outreach workers at risk.

It is unfortunate that Spencer Jones's statements are being attributed to the BMA, giving them an undeserved authority, and hence credibility. They do

not reflect BMA policy,⁵ and the BMA needs to clarify this with some urgency, look to its accountability, and return to evidence based practice.

Michael D E Goodyear assistant professor, Department of Medicine, Dalhousie University, Halifax, Nova Scotia, Canada B3H 2Y9 mgoodyear@dal.ca

Competing interests: None declared.

- 1 LifeStyleExtra. 'Legalise prostitution to save NHS cash' 29 March 2007. www.lse.co.uk/ShowStory.asp?story=MH2935662C&news_headline=legalise_prostitution_to_save_nhs_cash (accessed 5 April 2007).
- 2 Day M. BMA public health doctor is accused of stigmatising sex workers. *BMJ* 2007;334:767. (14 April.)
- 3 Ward H, Aral SO. Globalisation, the sex industry, and health. *Sex Transm Infect* 2006;82:345-7.
- 4 Jeal N, Salisbury C. Self-reported experiences of health services among female street-based prostitutes: a cross-sectional survey. *Br J Gen Pract* 2004;54:515-9.
- 5 BMA. Sexually transmitted infections: an update from the board of science. January 2007. www.bma.org/ap.nsf/content/stiup07.

INTENSIVE CARE

Help for patients and relatives

The DIPEX (Directory of Individual Patient Experiences) Research Group at the University of Oxford publishes a website of patients' experiences of health and illness. There is a module on intensive care (www.dipex.org/intensiveware) where you can read and watch interviews with patients of all ages talking about their experiences.¹

Lisa E Hinton graduate student
Department of Primary Care, University of Oxford,
Oxford OX3 7LF lisa.hinton@dphpc.ox.ac.uk

Competing interests: None declared.

- 1 Perrier E. Intensive care. *BMJ* 2007;334:796-7. (14 April.)

COPD

Drug management of COPD

In the second step of prescribing for long term management of chronic obstructive pulmonary disease, McIvor and Little recommend the addition of a long acting β_2 agonist (LABA) to a short acting β_2 agonist as needed, supplemented by stepwise addition of regular short or long acting anticholinergic inhalers.¹ I agree with this, and it is supported by evidence; tiotropium has advantages over a LABA. The National Institute for Health and Clinical Excellence (NICE) also recommends adding a LABA in its COPD guideline (No12, 2004).² At that time there was no evidence to show whether this was a useful combination, but now evidence shows the addition of a LABA to tiotropium has no additional benefit over tiotropium alone,³ so this is not really a suitable option.

The authors' next step is to add an inhaled corticosteroid or a combination of inhaled corticosteroid and a LABA as a single inhaler. If an inhaled corticosteroid is to be used in COPD then it should be in line with the NICE guideline. Evidence that a combination inhaler such as Seretide (fluticasone propionate and salmeterol) produces clinically important benefit over and above the separate components is lacking.⁴

If a patient on regular tiotropium and, as needed, a short acting β_2 agonist requires additional drug treatment then I suggest that the two options are to add an inhaled corticosteroid (as per NICE guidance) or add an oral mucolytic. To assess the effectiveness of drug therapy I suggest using five recommended questions (similar to using three questions in asthma).⁵

Peter D Burrill specialist pharmaceutical adviser for public health, Derbyshire County Primary Care Trust, Chesterfield S41 7PF
peter.burrill@derbyshirecountypct.nhs.uk

Competing interests: None declared.

- 1 McIvor A, Little P. Chronic obstructive pulmonary disease. *BMJ* 2007;334:798. (14 April.)
- 2 NICE. Chronic obstructive pulmonary disease. London: NICE, 2004. <http://guidance.nice.org.uk/CG12>
- 3 Aaron SD, Vandemheen KL, Fergusson D, Maltais F, Bourbeau J, Goldstein R, et al. Tiotropium in combination with placebo, salmeterol, or fluticasone-salmeterol for treatment of chronic obstructive pulmonary disease: a randomized trial. *Ann Intern Med* 2007;146:545-55.
- 4 Calverley PM, Anderson JA, Celli B, Ferguson GT, Jenkins C, Jones PW, et al. Salmeterol and fluticasone propionate and survival in chronic obstructive pulmonary disease. *N Engl J Med* 2007;356:775-89.
- 5 Jones PW. Health status measurement in chronic obstructive pulmonary disease. *Thorax* 2001; 56: 880-7.

COELIAC DISEASE

Decision tool needs to be developed for children

The decision tool for coeliac disease developed by Hopper et al is restricted to adults.¹ However, coeliac disease often presents in childhood with different symptoms and signs.^{2,3} It may therefore be inappropriate to use the clinical characteristics suggested by Hopper et al when dividing children into high and low risk groups with respect to coeliac disease.

The second part of the clinical decision tool of Hopper et al is testing with tissue transglutaminase autoantibodies.¹ Also here clinicians need to consider the implications of age. In a recent review by Rostom et al,⁴ the pooled sensitivity of human tissue transglutaminase IgA autoantibodies was lower in children (95.7%) than in adults (98.1%); more false negative cases can therefore be expected.

The clinical decision tool proposed by Hopper et al should be adapted for children, and thereafter tested prospectively in an unselected population.

Jonas F Ludvigsson consultant paediatrician
Department of Paediatrics, Örebro University Hospital,
701 85 Örebro, Sweden jonasludvigsson@yahoo.com

Anneli Ivarsson epidemiologist,
Department of Public Health and Clinical Medicine,
Umeå University, Sweden

Competing interests: None declared.

- 1 Hopper AD, Cross SS, Hurlstone DP, McAlindon ME, Lobo AJ, Hadjivassiliou M, et al. Pre-endoscopy serological testing for coeliac disease: evaluation of a clinical decision tool. *BMJ* 2007;334:729-32. (7 April.)
- 2 Fasano A. Clinical presentation of coeliac disease in the pediatric population. *Gastroenterology* 2005;128(4 suppl 1):S68-73.
- 3 Ludvigsson JF, Ansved P, Falth-Magnusson K, Hammersjö JA, Johansson C, Edvardsson S, et al. Symptoms and signs have changed in Swedish children with coeliac disease. *J Pediatr Gastroenterol Nutr* 2004;38(2):181-6.
- 4 Rostom A, Dube C, Cranney A, Saloojee N, Sy R, Garrity C, et al. The diagnostic accuracy of serologic tests for coeliac disease: a systematic review. *Gastroenterology* 2005;128(4 suppl 1):S38-46.

SHIFT WORK

The view from Denmark

As a junior doctor I worked all the different work rotas imaginable, both full time and flexibly, for seven years—and in three countries (the United Kingdom, Malaysia, and Denmark).¹ Now I work full time in Denmark but just 37 hours a week, maybe 40 on a bad week, and this includes on-call hours.

How does the Danish public health system function with such reduced doctors' working hours? Perhaps it is because Denmark has more doctors (they are still looking for more), and doctors are more efficient at time keeping during an average working day. The department I work in performs over 200 cleft cases and 150 free breast flaps in a year, plus all the skin cancer cases and burns injuries. These figures are similar to those of London teaching hospitals.

The downside? The training takes longer, but who cares if you can still have a good social and family life as well as enjoy your work? And there is no special treatment for women either: both parents are entitled to a shared maternity or paternity leave of up to nine months.

Shariha Khalid Erichsen clinical fellow, plastic surgery
Rigshospitalet, 2100, Copenhagen, Denmark
shariha@doctors.org.uk

Competing interests: None declared.

- 1 Ahmed-Little Y. Implications of shift work for junior doctors. *BMJ* 2007;334:777-8. (14 April.)