- 2 Bonnet JP, Basset T, Dijoux D. Abdominal inflammatory myofibroblastic tumors in children: report of an appendiceal case and review of the literature. *J Pediatr Surg* 1996;31:1311–14.
- 3 Maklouf HR, Sobin LH. Inflammatory myofibroblastic tumors (inflammatory pseudotumors) of the gastrointestinal tract: how closely are they related to inflammatory fibroid polyps. *Hum Pathol* 2002;33:307–15.
- 4 Karnak I, Senocak ME, Ciftci AO, et al. Inflammatory myofibroblastic tumors in children: diagnosis and treatment. *J Pediatr Surg* 2001;36:908–12.
- 5 Lee S-H, Fang Y-C, Luo J-P, et al. Inflammatory pseudotumour associated with chronic persistent Eikenella corrodens infection: a case report and brief review. *J Clin Pathol* 2003;56:868–70.
- 6 Vijayaraghavan R, Sujatha Y, Santosh KV, et al. Inflammatory fibroid polyp of jejunum causing jejuno-jejunal intussusception. *Indian J Gastroenterol* 2004;23:190–2.

Glomangioma of the lungs: a rare differential diagnosis of a pulmonary tumour

Glomangiomas (glomus tumours) are uncommon, and in most cases are benign perivascular tumours, usually located in the dermis of the extremities, especially in the subungual region of the fingers. They also occur in other organs and in other regions of the body. Histogenetically, they are derived from modified smooth muscle cells of the glomus apparatus, which is associated with temperature regulation. These tumours are composed of capillaries, sometimes of dilated endothelial bigger vessels surrounded by nests of uniform, cytoplasm-rich cells with isomorphic round or ovoid nuclei (glomus cells). They also contain a variable amount of smooth muscle tissue. Three different types of glomus tumours are described according to their different components. The common form is predominantly made up of glomus cells, with a smaller amount of vessels and smooth muscle cells. The second variant, called “glomangioma”, contains more partly dilated vessels resembling cavernous haemangiomas and is less circumscribed than the common form. In the most rare type, the “glomangiomyoma”, the glomus cells undergo a transition to elongated smooth muscle cells.

Immunohistochemically, glomus tumours usually show a positivity for vimentin and smooth muscle actin and a negativity for cytokeratin, neuroendocrine markers (chromogranin and others), CD31, CD34 and S-100.¹

These tumours are very rare in the lungs, and, to our knowledge, only 14 cases have been described, including two malignant glomus tumours.^{2–10}

Case report

A 64-year-old man was admitted to the hospital for resection of a carcinoma of the rectum. The preoperative computed tomography showed a round tumour (3.5 cm) in the left lower lung lobe that was removed 2 weeks after the resection of the rectum.

Findings

On macroscopic examination, the round tumour was reddish. Histological examination (haematoxylin and eosin, Elastica-van-Gieson, Prussian Blue and periodic-acid Schiff) showed lung tissue with partly compressed, partly hyperaemic and ectatic

Figure 1 (A) Part of the glomangioma with many partly dilated vessels and the pericapillary cytoplasm-rich cells with small round isomorphic nuclei (haematoxylin and eosin; original magnification $\times 25$). The left inset depicts the typical cells of the glomus tumour at a higher magnification (haematoxylin and eosin; original magnification $\times 100$). The right inset shows cytoplasmic staining of the tumour cells with antibodies against $\alpha 1$ -smooth muscle actin (original magnification $\times 100$). (B) Transition of the glomangioma from the solid, cell-rich part (left) to the more vascularised area (right; CD34; original magnification $\times 50$).

vessels that exceeded the normal density of alveolar capillaries (fig 1A,B). Cytoplasm-rich cells with round isomorphic nuclei were arranged diffusely perivascularly and in nests. Immunohistologically, the tumour cells featured a cytoplasmic staining for $\alpha 1$ -smooth muscle actin (fig 1A), whereas the reactions with antibodies against CD31, CD34, pancytokeratin and TTF-1 were negative.

Comment

On the basis of the histological and immunohistological results, a metastasis of the rectal carcinoma was ruled out and the tumour was identified as a glomangioma of the lung.

The other important differential diagnoses, including carcinoid small-cell lung cancer, melanocytic neoplasms, haemangiopericytoma, smooth muscle neoplasms, paraganglioma and primitive neuroectodermal

tumours, were also excluded by morphology and immunostaining pattern.

M Rössle

Institute of Pathology, Kantonsspital St Gallen, St Gallen, Switzerland

W Bayerle

Institute of Pathology, Kreisklinik München-Pasing, Munich, Germany

U Löhrs

Institute of Pathology, Ludwig-Maximilians-University, Munich, Germany

Correspondence to: M Rössle, Institute of Pathology, Kantonsspital Luzern, 6000 Luzern 16, Switzerland; matthias.roessle@ksl.ch

doi: 10.1136/jcp.2005.031237

Competing interests: None.

References

- 1 Weiss SW, Goldblum JR, eds. Perivascular tumors. *Enzinger and Weiss's soft tissue tumors*. 4th edn. St Louis, MO: Mosby, 2001:985–1001.
- 2 Ueno M, Nakashima O, Mishima M, et al. Pulmonary glomus tumor: CT and MRI findings. *J Thorac Imaging* 2004;131–4.
- 3 Hishida T, Hasegawa T, Asamura H, et al. Malignant glomus tumor of the lung. *Pathol Int* 2003;53:632–6.
- 4 Zhang Y, England DM. Primary pulmonary glomus tumor with contiguous spread to a peribronchial lymph node. *Ann Diagn Pathol* 2003;7:245–8.
- 5 Altorjay A, Arato G, Adame M, et al. Synchronous multiple glomus tumors of the esophagus and lung. *Hepatogastroenterology* 2003;50:687–90.
- 6 Yilmaz A, Bayramgürler B, Aksoy F, et al. Pulmonary glomus tumour: a case initially diagnosed as carcinoid tumour. *Respirology* 2002;7:369–71.
- 7 Gaertner EM, Steinberg DM, Huber M, et al. Pulmonary and mediastinal glomus tumors-report of five cases including a pulmonary glomangiosarcoma: a clinicopathologic study with literature review. *Am J Surg Pathol* 2000;24:1105–14.
- 8 Koss MN, Hochholzer L, Moran CA. Primary pulmonary glomus tumor: a clinicopathologic and immunohistochemical study of two cases. *Mod Pathol* 1998;11:253–8.
- 9 Alt B, Huffer WE, Belchis DA. A vascular lesion with smooth muscle differentiation presenting as a coin lesion in the lung: glomus tumor versus hemangiopericytoma. *Am J Clin Pathol* 1983;80:765–71.
- 10 Tang CK, Tokar C, Foris NP, et al. Glomangioma of the lung. *Am J Surg Pathol* 1978;2:103–9.

Vertebral artery dissection revisited

Vertebral artery dissection is considered to be one of the most difficult dissections that a forensic pathologist has to undertake. Traditionally, vertebral arteries have been dissected using two time-tested procedures. The first procedure is carried out by removing the neck block and decalcifying the block.¹ The neck block can be removed using the anterior or the posterior approach. The anterior approach is less time consuming but is more difficult to carry out. The posterior approach, on the other hand, is more time consuming but is simpler to carry out. The other procedure requires the use of special instruments for dissection.² These are a Swann–Morton number 5 scalpel handle with number 11 blade, Swann–Morton number 3 scalpel handle with number 12 blade

Figure 1 Vertebral arteries in situ (a probe has been placed underneath for better appraisal).

Figure 2 Heart, aorta, its major tributaries along with vertebral arteries.

(curved) and a pair of Universal wire-cutting scissors (curved heavy pattern). These three instruments, although considered to be cheap and readily available, were not so when we tried to find them for our use. As a result, it was not possible for us to use these instruments. We believe that forensic pathologists in third world countries and perhaps even in affluent countries, but working in remote peripheral areas, may be facing the same problem. There may even be a tendency on the part of these forensic pathologists to skip this vital step during postmortem examinations. Keeping this in mind, we developed a

simple procedure, which requires simple instruments.

Method

The routine Y-shaped skin incision is made, after which a horizontal incision is made above both the clavicles. Both the clavicles are then removed from the body by disarticulating the sterno-clavicular and acromio-clavicular joints. This is done to enable the pathologist to have a clear view of the origin of the vertebral artery. The rib cage is then removed. As a wider view of the thoracic

cavity is required, the ribs are not severed from the costo-chondral junction but are cut in the anterior axillary line. Each rib is cut using the bone cutter and the intercostal muscles are cut using the scalpel. We need to be careful while undertaking this part of the dissection, as a blind dissection at the time of removing the uppermost part of the rib cage can lead to the inadvertent severing of the vertebral artery.

After removing the thoracic cage, the pericardial sac is lifted, cut and reflected. The branches of the arch of the aorta are seen. The origin of the vertebral artery is then identified as it originates from the subclavian artery. The vertebral artery is traced upwards in the neck. From this stage onwards, the neck is kept extended by putting a wooden block under the shoulders. The vertebral artery is then dissected in layers, until the thyroid cartilage is reached. At this stage, the vertebral artery can be visualised as going into the transverse foramen of the sixth cervical vertebra. The next part of the dissection is the most difficult. In the absence of the special instruments (mainly the bone-cutting wires) used for the dissection of the vertebral arteries, we had to use normal dissection instruments. We had to cut the transverse foramen from the sixth to the second vertebrae by using these instruments—namely, the ordinary curved scissors, and a toothed and a plain forceps. The scissors are used to cut the transverse processes. The toothed forceps are used to part the pieces of bone and soft tissue, and the plain (non-traumatic) forceps are used to hold the vertebral artery and bring it out of the vertebrae. We have to ascertain that while the transverse foramen is being cut, there is no damage to the artery (this requires practice). When we reach the second cervical vertebra, the artery can be seen to turn outwards after the transverse process of the second cervical vertebra. This is the upper limit to which we can clear the area using these instruments to dissect the artery. The further part can be removed while the brain is being removed. This may leave a small part of the artery, about ½–1 cm, unobserved. But this drawback can be overcome by cutting the artery as deep as possible from the cranial end, while removing the brain. This enables us to visualise the vertebral artery in its entirety.

Conclusion

A simple method for vertebral artery dissection has been described. We hope that this method will encourage more forensic pathologists around the world to undertake it more routinely, which may in turn lead to better administration of justice.

A Aggrawal, P Setia

Department of Forensic Medicine, Maulana Azad Medical College, New Delhi, India

Correspondence to: A Aggrawal, S-299, Greater Kailash-1, 110048 New Delhi, India; dr_anil@hotmail.com

doi: 10.1136/jcp.2005.030742

Accepted for publication 17 October 2005

Competing interests: None declared.

References

- 1 Sheaf MT, Hopster DJ. Organ dissection-cardiovascular system. In: Sheaf MT, Hopster DJ,

eds. *Post mortem technique handbook*. Great Britain: Springer, 2001:113–4.

- 2 Bromilow A, Burns J. Technique for removal of the vertebral arteries. *J Clin Pathol* 1985;**38**:1400–2.

A novel mutation, Ile289Thr, in the ALAS2 gene in a family with pyridoxine responsive sideroblastic anaemia

X-linked sideroblastic anaemia (XLSA; OMIM 301 300) is characterised by accumulation of inorganic iron in erythroblast mitochondria, visualised on staining as distinctive perinuclear rings. It arises from a deficiency of the erythroid specific isoenzyme of δ -aminolaevulinate synthase (ALAS2; E.C. 2.3.1.3.7), caused mainly by mutations affecting the catalytic or substrate-binding domains.^{1,2} ALAS2 uses pyridoxal-5-phosphate as a cofactor to catalyse the first, rate-limiting step of erythroid haem synthesis and pyridoxine treatment can alleviate anaemia in many cases, although the response is variable and affected by factors such as mutation, age and iron load.^{3,4}

A 40-year-old man presented with microcytic anaemia (haemoglobin 7.9 g/dl, mean corpuscular volume 58 fl, and mean corpuscular haemoglobin 15 pg), no evidence of thalassaemia, raised serum ferritin and 88% saturated transferrin. The blood film was dimorphic, with basophilic stippling and Pappenheimer bodies (fig 1A). Bone marrow cytology showed dysplastic erythropoiesis and ringed sideroblasts (fig 1B). After a 10-week course of 200 mg pyridoxine twice daily, the haemoglobin concentration rose to 12.4 g/dl. A liver biopsy, undertaken because of increased liver transaminases, showed heavy parenchymal iron deposition, but no fibrosis or cirrhosis. Screening for *HFE* gene mutations was negative. During fortnightly venesections over 8 months, haemoglobin remained stable, serum ferritin fell to 125 μ g/l and liver transaminases returned to normal. The patient remains on pyridoxine and on a three-monthly maintenance venesection. Family history included a maternal grandfather and a brother (haemoglobin 13.1 g/dl, mean corpuscular volume 82 fl, mean corpuscular haemoglobin 26 pg) with microcytic anaemia.

Sequencing the coding region and intron or exon boundaries of *ALAS2* detected a T918C change on exon 7, predicting substitution of the highly conserved isoleucine by threonine

at codon 289. The Ile289Thr mutation was present in the affected brother but not in an unaffected brother. Screening 150 alleles from 75 healthy women, mostly of northern European ancestry, and 45 alleles from 34 (11 women, 23 men) unrelated patients with congenital or inherited sideroblastic anaemia excluded *ALAS2*T918C as a common ($\geq 5\%$), but not rare, north European polymorphism.

The novel mutation Ile289Thr is therefore probably responsible for the development of sideroblastic anaemia in this family, despite a variable associated phenotype not uncommon for this disorder.³ Ile289 is located in a highly conserved region close to previously reported mutations, Gly291Ser and Lys299Gln, also associated with marked responses to pyridoxine,⁴ and to His285, predicted by homology to have an important steric relationship with bound pyridoxal-5-phosphate.^{4,5}

Acknowledgements

We thank Mrs Angela McVeigh for technical assistance and Dr P J Wilson for clinical data on the patient's family history.

M J Percy, R J G Cuthbert

Department of Haematology, Belfast City Hospital, Belfast, UK

A May

Department of Haematology, Cardiff University School of Medicine, Cardiff, UK

M F McMullin

Department of Haematology, Queen's University, Belfast

Correspondence to: Dr M J Percy, Department of Haematology, Floor C, Belfast City Hospital, Lisburn Road, Belfast BT9 7AB, UK; melanie.percy@bll.n-i.nhs.uk

doi: 10.1136/jcp.2005.030932

Accepted for publication 31 October 2005

References

- 1 Anderson KE, Sassa S, Bishop DF, *et al*. Disorders of heme biosynthesis: X-linked sideroblastic anemia and the porphyrias. In: Scriver CR, Beaudet AL, Sly WS, Valle D, Childs B, Kinzler KW, Vogelstein B, eds. *The metabolic and molecular bases of inherited disease*. Vol 2, 8th edn. New York, NY: McGraw-Hill, 2001:2991–3062.
- 2 Bottomley SS. Sideroblastic anaemias. In: Greer JP, Foerster J, Lukens JN, Rodgers GM, Paraskevas F, Glader B, eds. *Wintrabe's clinical hematology*. Vol 1, 11th edn. Philadelphia: Lippincott Williams & Wilkins, 2004:1011–33.

- 3 May A, Bishop D. The molecular biology and pyridoxine responsiveness of X-linked sideroblastic anaemia. *Haematologica* 1998;**83**:56–70.
- 4 Shoolingin-Jordan PM, Al-Daihan S, Alexeev D, *et al*. 5-Aminolaevulinic acid synthase: mechanism, mutations and medicine. *Biochim Biophys Acta* 2003;**1647**:361–6.
- 5 Alexeev D, Alexeeva M, Baxter RL, *et al*. The crystal structure of 8-amino-7-oxononanoate synthase: a bacterial PLP-dependent, acyl-coA-condensing enzyme. *J Mol Biol* 1998;**284**:401–19.

Cystic dysplasia of testis: a case report

Cystic dysplasia of the testis is a rare congenital defect that results in the formation of numerous irregularly shaped cystic spaces in the mediastinum testis. The clinical presentation of such a disorder usually occurs in the paediatric age group as an asymptomatic scrotal swelling that can mimic testicular tumours. The case presented in this paper is the first identified and diagnosed case of cystic dysplasia in Kashmir, India.

A 2-year-old boy presented with a history of scrotal swelling since birth and was admitted to the Department of Surgery, Shri Mahraja Hari Singh Hospital, Srinagar, Kashmir, associated with Government Medical College, Srinagar, Kashmir, India.

On examination, the scrotum showed an enlarged right testis with a cystic feel. A trans-illumination test was positive. Scrotal ultrasound also showed cystic spaces in the testis. The testis was 7 $\times$ 3 cm with all three coverings and had a cystic feel, with small cystic spaces ranging in size from 0.2 to 1.5 cm on the cut section, affecting the whole testicular mass. Microscopic examination showed that the cysts were lined by low cuboidal flattened epithelium resembling the epithelium of the rete testis, separated from each other by fibrous septae with a complete replacement of normal testicular tissue.

Cystic dysplasia of the rete testis is a rare abnormality often associated with ipsilateral agenesis of the kidney. This malformation is due to a developmental defect of the mesonephric duct, which is the cause of both the dilatation of testicular rete testis and renal agenesis.¹ It is characterised by multiple irregular cystic spaces in the mediastinum of the testis that may affect the whole gonad and is associated with renal malformations (agenesis or cystic dysplasia).²

Embryologically, when the duct system of the testis is formed, seminiferous tubules anastomose with one another to form the rete testis. The rete testis in turn establishes contact with persisting mesonephric tubules, which form vasa-efferentia, the cranial part of the mesonephric duct becomes highly coiled on itself to form the epididymis and the distal part becomes the vas deferens. The seminal vesicle arises on either side of the mesonephric duct as a diverticulum. The part of the mesonephric duct between the opening into the prostatic urethra and the origin of the diverticulum forms the ejaculatory duct.³

A defect in the connection between efferent ductules derived from the mesonephric epithelium and the rete testis and seminiferous tubules seems to be the most likely explanation for the pathogenesis of cystic dysplasia of the testis.^{1,4}

In our patient, teratoma was clinically suspected and orchidectomy was carried out

Figure 1 Blood film and bone marrow smear from the index case. (A) Wright-Giemsa stain of the peripheral blood showing dimorphic red cells and the presence of Pappenheimer bodies (indicated by arrows). (B) Perl's stain of the bone marrow aspirate showing the presence of ringed sideroblasts (indicated by the arrow).