eds. *Post mortem technique handbook*. Great Britain: Springer, 2001:113–4.

- 2 Bromilow A, Burns J. Technique for removal of the vertebral arteries. *J Clin Pathol* 1985;**38**:1400–2.

A novel mutation, Ile289Thr, in the ALAS2 gene in a family with pyridoxine responsive sideroblastic anaemia

X-linked sideroblastic anaemia (XLSA; OMIM 301 300) is characterised by accumulation of inorganic iron in erythroblast mitochondria, visualised on staining as distinctive perinuclear rings. It arises from a deficiency of the erythroid specific isoenzyme of δ -aminolaevulinic synthase (ALAS2; E.C. 2.3.1.3.7), caused mainly by mutations affecting the catalytic or substrate-binding domains.^{1,2} ALAS2 uses pyridoxal-5-phosphate as a cofactor to catalyse the first, rate-limiting step of erythroid haem synthesis and pyridoxine treatment can alleviate anaemia in many cases, although the response is variable and affected by factors such as mutation, age and iron load.^{3,4}

A 40-year-old man presented with microcytic anaemia (haemoglobin 7.9 g/dl, mean corpuscular volume 58 fl, and mean corpuscular haemoglobin 15 pg), no evidence of thalassaemia, raised serum ferritin and 88% saturated transferrin. The blood film was dimorphic, with basophilic stippling and Pappenheimer bodies (fig 1A). Bone marrow cytology showed dysplastic erythropoiesis and ringed sideroblasts (fig 1B). After a 10-week course of 200 mg pyridoxine twice daily, the haemoglobin concentration rose to 12.4 g/dl. A liver biopsy, undertaken because of increased liver transaminases, showed heavy parenchymal iron deposition, but no fibrosis or cirrhosis. Screening for *HFE* gene mutations was negative. During fortnightly venesections over 8 months, haemoglobin remained stable, serum ferritin fell to 125 μ g/l and liver transaminases returned to normal. The patient remains on pyridoxine and on a three-monthly maintenance venesection. Family history included a maternal grandfather and a brother (haemoglobin 13.1 g/dl, mean corpuscular volume 82 fl, mean corpuscular haemoglobin 26 pg) with microcytic anaemia.

Sequencing the coding region and intron or exon boundaries of *ALAS2* detected a T918C change on exon 7, predicting substitution of the highly conserved isoleucine by threonine

at codon 289. The Ile289Thr mutation was present in the affected brother but not in an unaffected brother. Screening 150 alleles from 75 healthy women, mostly of northern European ancestry, and 45 alleles from 34 (11 women, 23 men) unrelated patients with congenital or inherited sideroblastic anaemia excluded *ALAS2*T918C as a common ($\geq 5\%$), but not rare, north European polymorphism.

The novel mutation Ile289Thr is therefore probably responsible for the development of sideroblastic anaemia in this family, despite a variable associated phenotype not uncommon for this disorder.⁵ Ile289 is located in a highly conserved region close to previously reported mutations, Gly291Ser and Lys299Gln, also associated with marked responses to pyridoxine,⁴ and to His285, predicted by homology to have an important steric relationship with bound pyridoxal-5-phosphate.^{4,5}

Acknowledgements

We thank Mrs Angela McVeigh for technical assistance and Dr P J Wilson for clinical data on the patient's family history.

M J Percy, R J G Cuthbert

Department of Haematology, Belfast City Hospital, Belfast, UK

A May

Department of Haematology, Cardiff University School of Medicine, Cardiff, UK

M F McMullin

Department of Haematology, Queen's University, Belfast

Correspondence to: Dr M J Percy, Department of Haematology, Floor C, Belfast City Hospital, Lisburn Road, Belfast BT9 7AB, UK; melanie.percy@bll.n-i.nhs.uk

doi: 10.1136/jcp.2005.030932

Accepted for publication 31 October 2005

References

- 1 Anderson KE, Sassa S, Bishop DF, *et al*. Disorders of heme biosynthesis: X-linked sideroblastic anemia and the porphyrias. In: Scriver CR, Beaudet AL, Sly WS, Valle D, Childs B, Kinzler KW, Vogelstein B, eds. *The metabolic and molecular bases of inherited disease*. Vol 2, 8th edn. New York, NY: McGraw-Hill, 2001:2991–3062.
- 2 Bottomley SS. Sideroblastic anaemias. In: Greer JP, Foerster J, Lukens JN, Rodgers GM, Paraskevas F, Glader B, eds. *Wintrabe's clinical hematology*. Vol 1, 11th edn. Philadelphia: Lippincott Williams & Wilkins, 2004:1011–33.

- 3 May A, Bishop D. The molecular biology and pyridoxine responsiveness of X-linked sideroblastic anaemia. *Haematologica* 1998;**83**:56–70.
- 4 Shoolingin-Jordan PM, Al-Daihan S, Alexeev D, *et al*. 5-Aminolaevulinic acid synthase: mechanism, mutations and medicine. *Biochim Biophys Acta* 2003;**1647**:361–6.
- 5 Alexeev D, Alexeeva M, Baxter RL, *et al*. The crystal structure of 8-amino-7-oxononanoate synthase: a bacterial PLP-dependent, acyl-coA-condensing enzyme. *J Mol Biol* 1998;**284**:401–19.

Cystic dysplasia of testis: a case report

Cystic dysplasia of the testis is a rare congenital defect that results in the formation of numerous irregularly shaped cystic spaces in the mediastinum testis. The clinical presentation of such a disorder usually occurs in the paediatric age group as an asymptomatic scrotal swelling that can mimic testicular tumours. The case presented in this paper is the first identified and diagnosed case of cystic dysplasia in Kashmir, India.

A 2-year-old boy presented with a history of scrotal swelling since birth and was admitted to the Department of Surgery, Shri Mahraja Hari Singh Hospital, Srinagar, Kashmir, associated with Government Medical College, Srinagar, Kashmir, India.

On examination, the scrotum showed an enlarged right testis with a cystic feel. A trans-illumination test was positive. Scrotal ultrasound also showed cystic spaces in the testis. The testis was 7 $\times$ 3 cm with all three coverings and had a cystic feel, with small cystic spaces ranging in size from 0.2 to 1.5 cm on the cut section, affecting the whole testicular mass. Microscopic examination showed that the cysts were lined by low cuboidal flattened epithelium resembling the epithelium of the rete testis, separated from each other by fibrous septae with a complete replacement of normal testicular tissue.

Cystic dysplasia of the rete testis is a rare abnormality often associated with ipsilateral agenesis of the kidney. This malformation is due to a developmental defect of the mesonephric duct, which is the cause of both the dilatation of testicular rete testis and renal agenesis.¹ It is characterised by multiple irregular cystic spaces in the mediastinum of the testis that may affect the whole gonad and is associated with renal malformations (agenesis or cystic dysplasia).²

Embryologically, when the duct system of the testis is formed, seminiferous tubules anastomose with one another to form the rete testis. The rete testis in turn establishes contact with persisting mesonephric tubules, which form vasa-efferentia, the cranial part of the mesonephric duct becomes highly coiled on itself to form the epididymis and the distal part becomes the vas deferens. The seminal vesicle arises on either side of the mesonephric duct as a diverticulum. The part of the mesonephric duct between the opening into the prostatic urethra and the origin of the diverticulum forms the ejaculatory duct.³

A defect in the connection between efferent ductules derived from the mesonephric epithelium and the rete testis and seminiferous tubules seems to be the most likely explanation for the pathogenesis of cystic dysplasia of the testis.^{1,4}

In our patient, teratoma was clinically suspected and orchidectomy was carried out

Figure 1 Blood film and bone marrow smear from the index case. (A) Wright-Giemsa stain of the peripheral blood showing dimorphic red cells and the presence of Pappenheimer bodies (indicated by arrows). (B) Perl's stain of the bone marrow aspirate showing the presence of ringed sideroblasts (indicated by the arrow).

on the right side. The histopathological diagnosis was cystic dysplasia of the testis, but evaluation of the patient for other urogenital anomalies was not possible as the patient did not come for follow-up.

Today, the non-operative management of cystic dysplasia of the testis represents an effective alternative option in these patients and the primary benefit of this approach is the preservation of endocrine function and spermiogenic activity. The natural history of untreated cystic dysplasia of the testis and its effect on normal testicular tissue are, however, still unknown. Therefore, a long-term follow-up is recommended,³ along with evaluation for associated urological anomalies.

M L Bhat, Z Rasool

Department of Pathology, Government Medical College, Srinagar, Kashmir, India

S M Kadri

Regional Institute of Health and Family Welfare (RIHFW) DHS, Srinagar

N Wani

Department of Surgery, Shri Mahraja Hari Sirgh Hospital, Srinagar

G Hassan, D Mumtaz, W Qureshi

Department of Medicine, Government Medical College

B Shiekh

Department of Pathology, Government Medical College

R Dwivedi

VA Medical Center, Minneapolis, Minnesota, USA

Correspondence to: M L Bhat, Department of Pathology, Government Medical College, Srinagar, Kashmir, India; kadrism@sancharnet.in

doi: 10.1136/jcp.2005.032888

References

- 1 **Cimador M, Rosone G, Castagnetti M, et al.** Cystic dysplasia of rete testis associated with ipsilateral renal agenesis. *Case Rep Minerva Pediatr* 2003;**55**:175-9.
- 2 **Camassei FD, Francalanci P, Ferro F, et al.** Cystic dysplasia of the rete testis: report of two cases and review of the literature. *Pediatr Dev Pathol* 2002;**5**:206-10.
- 3 **Glantz L, Hansen K, Caldame A, et al.** Cystic dysplasia of the testis. *Hum Pathol* 1993;**24**:1142-5.
- 4 **Nistal M, Regadera J, Paniagua R.** Cystic dysplasia of the testis. Light and electron microscopic study of three cases. *Arch Pathol Lab Med* 1984;**108**:579-83.
- 5 **Rovellini P, Bianchi A, Bottanelli A, et al.** Cystic dysplasia of the testis: report of a case and review of the literature. *Arch Ital Urol Androl* 2001;**73**:193-8.

Plasma cell satellitism in plasma cell myeloma

Plasma cell satellitism around macrophages has been reported in a patient with plasma cell myeloma (multiple myeloma) and amyloidosis.¹ The formation of the rosettes was attributed to the secretion of amyloid by macrophages.¹⁻³ We report a case of plasma cell satellitism without associated amyloidosis. It is important to recognise this rare morphological finding to prevent confusion with other conditions, such as nucleated red blood cells clustering around macrophages or

Figure 1 (A, B) Plasma cells forming rosette-like structures around a centrally located macrophage.

metastatic tumours exhibiting rosette formation.

The investigation of a 63-year-old man with back pain showed hypercalcaemia, anaemia and high levels of immunoglobulin G, proved to be monoclonal by serum immunofixation. A bone survey showed lytic lesions throughout the shafts of the humeri and femurs. Bone marrow examination contained paucicellular aspirate smears, predominated by plasma cells with large eccentric nuclei and prominent nucleoli. Most of the plasma cells formed distinct rosettes, with tapered basophilic cytoplasmic projections directed towards a centrally located macrophage (fig 1). The hypercellular (70%) bone marrow core biopsy specimen contained 20% plasma cells as scattered cells and focal clusters, but no identifiable rosettes.

Immunohistochemistry for κ immunoglobulin light chains highlighted most of the plasma cells. CD68 immunostains showed a few scattered cells. No amyloid was seen with Congo red stain. Flow cytometric immunophenotypic studies of the aspirate consisted of a monotypic κ -restricted plasma cell population that was CD38, CD138 and CD56 positive, and CD45, CD20, CD25 and CD52 negative. Owing to poor response to chemotherapy, the patient underwent autologous stem cell transplantation. Bone marrow biopsy after 4 months showed 3% myeloma cells, without rosetting in the aspirate.

The mechanism mediating the communication between the myeloma cell and the macrophage is unknown. Previously, one case with plasma cell rosettes around macrophages in the bone marrow aspirate was reported in a patient with plasma cell myeloma and amyloidosis.¹ Clustering of plasma cells around macrophages has also been observed in vitro in myeloma cell cultures, in which amyloid was seen.^{2,3} In this case, the plasma cell satellitism in conjunction with the absence of clinical and morphological signs of amyloidosis suggests that the rosetting is not related to the production or processing of amyloid by macrophages as previously theorised.¹⁻³ Additionally, plasma cell rosetting without associated macrophages has also been reported, raising the possibility that macrophages may not be necessary for this interaction.^{4,5}

N D Dimov*, D L Zynger*, L C Peterson

McGaw Medical Centre of Northwestern University, Chicago, USA

Correspondence to: N D Dimov, Mc Gaw Medical Centre of North Western University, 251 E Huron Street, Feinberg 7 325B, Chicago, IL 60611, USA; n-dimov@md.northwestern.edu

doi: 10.1136/jcp.2005.034710

Accepted for publication 6 January 2006

*These authors contributed equally to this work.

Competing interests: None declared.

References

- 1 **Pillay TS, Sayers G, Bird AR, et al.** Plasmacytic reticulum cell satellitism in multiple myeloma associated with amyloidosis. *J Clin Pathol* 1992;**45**:623-4.
- 2 **Durie BGM, Persky BP, Soehnlen BJ, et al.** Amyloid production in human myeloma stem cell culture, with morphologic evidence of amyloid secretion by associated macrophages. *N Engl J Med* 1982;**307**:1689-92.
- 3 **Zucker-Franklin D, Frangione B.** Amyloid in myeloma stem cell culture. *N Engl J Med* 1983;**308**:1164-5.
- 4 **Wong KF, Chan JK, Chu YC.** Multiple myeloma with tadpolelike cells and rosette formation. *Am J Clin Pathol* 1992;**98**:630-2.
- 5 **O'Malley, DP.** A rosette by any other name. *Arch Pathol Lab Med* 2003;**128**:1053.

BOOK REVIEW

Phages: their role in bacterial pathogenesis and biotechnology

Edited by M K Waldor, D I Friedman, S L Adhya. Washington DC: Published by ASM Press, 2005, \$119-95 (hardcover), pp 468. ISBN 1-55581-307-0

Any book that contains the phrase "rolling global-scale orgy of bacteriophage sex" will grab my attention, but then again, I am a bacteriophage enthusiast. To be honest, I'm really only an amateur and this book is primarily for the professional enthusiast, and at \$120, it is likely to be read only by those who work with bacteriophages regularly. This is a shame, as it contains some fascinating material. Did you know, for instance, that, with a total population estimated at any one time to be 10^{31} , phages are the most numerous organisms on the planet? I didn't know until I read this book. Another amazing fact that had me firing off emails to my long-suffering colleagues was that if all 10^{31} of these phages were laid end to end, they