PostScript

LETTER TO THE EDITOR

An uncommon pattern of cardiac invasion in hepatocellular carcinoma HCC, hepatocellular carcinoma

Hepatocellular carcinoma (HCC), the most common primary malignant tumour of the liver, is a highly vascular neoplasm usually arising in a cirrhotic liver. In addition to the diagnosis of primary lesion, an assessment for extrahepatic metastasis is needed. Herein, an extremely uncommon pattern of heart invasion in a patient with advanced HCC is described.

A 75-year-old man was admitted to the Hospital of Zumarraga, Zumarraga, Spain, for moderate abdominal pain, asthenia and weight loss. He denied having dyspnoea, orthopnoea, cough, palpitation or peripheral oedemas. Three years earlier, HCC had been diagnosed in the eighth segment, and the patient had received several courses of transcatheter arterial chemoembolisation. An objective examination showed hepatosplenomegaly and ascites, with no signs of cardiac failure. Temperature, blood pressure and cardiac frequency were normal. Laboratory data showed raised serum levels of aspartate transaminase and α fetoprotein. Electrocardiogram and chest x ray showed no abnormalities. A total-body computed tomography scan highlighted diffuse infiltration of hepatic lobes (fig 1), mesenteric nodules with ascites (suggestive of carcinomatosis), multiple lymphadenopathy in celiac trunk and portocava veins and a mass in anterior pericardium (fig 2) with no pericardial or

Figure 1 Computed tomography scan of the abdomen showing diffuse infiltration of both hepatic lobes.

Figure 2 Computed tomography scan of the chest showing a rounded, oval-shaped mass in the anterior pericardium.

pleural effusion. Peritoneal fluid showed an exudate but no malignant cells. The cava and suprahepatic veins, and right atrium did not show evidence of tumoral invasion. The patient died 2 weeks later from acute renal failure. An autopsy was not authorised.

Practically all malignant secondary neoplasms may affect the heart. Metastatic spreading to the heart has been identified in about one fifth of all patients who have metastatic cancer. The pericardium may be invaded, mainly by carcinoma of the lung followed by breast carcinoma, lymphoma and leukaemia. Rare metastatic neoplasms of the pericardium include carcinomas of the digestive tract, such as colon, oesophagus, stomach and others.¹ HCC spreads most often to the lungs, peritoneum, adrenal glands and bones. The incidence of cardiac metastasis of HCC has been reported in up to 4.1% of patients.² Classically, several routes of pericardial involvement in malignancy have been described,¹ such as haematogenous route, by direct extension from intrathoracic cancer, by retrograde lymphatic embolisation from the mediastinal lymph nodes, by continuous spreading through the diaphragm with infiltration of the left atrium and by tumour thrombus in the venous system. The mechanism of cardiac involvement in HCC is related to the marked propensity of this tumour for vascular invasion and extension, such as tumour thrombus including the hepatic vein and the inferior vena cava, thus easily reaching the right cardiac cavities.²⁻³ Pericardial involvement through direct invasion of the anterior mediastinum is another rare mechanism that has been reported in a few patients with advanced HCC after several courses of transcatheter arterial chemoembolisation.⁴ This patient represents an extremely

unusual pattern of cardiac involvement from HCC because of intravascular tumour thrombus or growth; transdiaphragmatic infiltration or direct extension from mediastinum or lymph nodes was absent. Thus, the route of isolated pericardial metastasis was probably haematogenous.

E Anton

Department of Internal Medicine, Hospital of Zumarraga, Zumarraga, Spain

A Astiazaran

Department of Radiology, CHDonostia, Donostia, Spain

Correspondence to: E Anton, Department of Internal Medicine, B^o Argixao s/n, 20700-Zumarraga (Guipuzcoa), Spain; ejaranda@hotmail.com

doi: 10.1136/jcp.2005.035238

Accepted for publication 22 November 2005

Competing interests: None declared.

References

- 1 Hanfling SM. Metastatic cancer to the heart: review of the literature and report of 127 cases. *Circulation* 1960;22:474-83.
- 2 Kojiro M, Nakahara H, Sugihara S, et al. Hepatocellular carcinoma with intra-atrial tumor growth. A clinicopathologic study of 18 autopsy cases. *Arch Pathol Lab Med* 1984;108:989-92.
- 3 Nakashima T, Okuda K, Kojiro M, et al. Pathology of hepatocellular carcinoma in Japan: 232 consecutive cases autopsied in ten years. *Cancer* 1983;51:863-77.
- 4 Huang CC, Ng WW, Chiang JH, et al. Hepatocellular carcinoma with mediastinal and pericardial invasion: report of two cases. *Chin Med J* 1999;62:891-5.

CORRECTION

doi: 10.1136/jcp.2004.019323.corr2

We regret that there was an error in the correction that was published in the last issue. Reference 9 in the article by Chatterjee JS, Youssef AHK, Brown RM, et al. Congenital nodular multiple glomangioma: a case report. *J Clin Pathol* 2005;58:102-10 should read as follows: Öztekin HH. Popliteal glomangioma mimicking Baker's cyst in a 9-year-old child: an unusual location of a glomus tumor. *Arthroscopy* 2003;19:e67-e71.