Take-home message

Prostatic stromal sarcoma is not simply a stromal neoplasm as previously believed. Not only the stromal component but also the epithelial component may have a malignant potential.

epithelia of the recurrent tumours were compatible with HGPIN: (1) markedly higher levels of Ki-67 labelling index (which rises higher in prostatic adenocarcinoma and HGPIN than in benign glands^{10–11}) in the epithelia; (2) discontinuous staining for HMWCK (a staining pattern typical of HGPIN¹²); (3) weak staining for AMACR (which stains the majority of prostatic adenocarcinoma and HGPIN¹³). The malignant transformation of the epithelia surrounded by malignant stromal cells implies the existence of an epithelial–stromal interaction—that is, an aberrant stromal micro-environment promoted malignant transformation of the epithelia.¹⁴

Another remarkable finding in the present case was that basal cells, in addition to atypical stromal and secretory cells, existed in multiple recurrent tumours outside the prostate. Although a recent molecular genetic study has demonstrated that both epithelial and stromal components of prostatic phyllodes tumours are clonal,¹⁵ the clonality of basal cells in prostatic stromal tumours has not been examined. In a previous study, McCarthy *et al* concluded that epithelial and stromal components might have different clonal origins.¹⁵ Contrary to McCarthy *et al*'s study, our finding raises the possibility that both components, including basal cells, have a common clonal origin.

In summary, we report a previously unreported case of extraprostatic recurrent PSS in which the epithelial component represented massive HGPIN. It is important to note that mixed epithelial–stromal tumour of the prostate is not solely a stromal neoplasm as previously believed, and that not only the stromal but also the epithelial component of this tumour may have a malignant potential.

Authors' affiliations

Tepei Morikawa, Akiteru Goto, Satoshi Ota, Masashi Fukayama, Department of Human Pathology, Graduate School of Medicine, University of Tokyo, Tokyo, Japan

Kyoichi Tomita, Yuzuri Tsurumaki, Tadaichi Kitamura, Department of Urology, The University of Tokyo Hospital, Tokyo, Japan

Competing interests: None declared.

Correspondence to: Dr T Morikawa, Department of Human Pathology, Graduate School of Medicine, University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, 113-0033 Tokyo, Japan; tmorikawa-ky@umin.ac.jp

Accepted 12 June 2006

REFERENCES

- Gaudin PB, Rosai J, Epstein JI. Sarcomas and related proliferative lesions of specialized prostatic stroma: a clinicopathologic study of 22 cases. *Am J Surg Pathol* 1998;**22**:148–62.
- Bostwick DG, Hossain D, Qian J, *et al*. Phyllodes tumor of the prostate: long-term followup study of 23 cases. *J Urol* 2004;**172**:894–9.
- Chevillat J, Cheng L, Algaba F, *et al*. Mesenchymal tumours. In: Eble JN, Sauter G, Epstein JI, *et al*, eds. *Pathology and genetics of tumours of the urinary system and male genital organs*. Lyon, France: IARC Press, 2004:209–11.
- Lopez-Beltran A, Gaeta JF, Huben R, *et al*. Malignant phyllodes tumor of prostate. *Urology* 1990;**35**:164–7.
- Yum M, Miller JC, Agrawal BL. Leiomyosarcoma arising in atypical fibromuscular hyperplasia (phyllodes tumor) of the prostate with distant metastasis. *Cancer* 1991;**68**:910–15.
- Young JF, Jensen PE, Wiley CA. Malignant phyllodes tumor of the prostate. A case report with immunohistochemical and ultrastructural studies. *Arch Pathol Lab Med* 1992;**116**:296–9.
- Watanabe M, Yamada Y, Kato H, *et al*. Malignant phyllodes tumor of the prostate: retrospective review of specimens obtained by sequential transurethral resection. *Pathol Int* 2002;**52**:777–83.
- Agrawal V, Sharma D, Wadhwa N. Case report: malignant phyllodes tumor of prostate. *Int Urol Nephrol* 2003;**35**:37–9.
- Chen TA, Chou JM, Sun GH, *et al*. Malignant phyllodes tumor of the prostate. *Int J Urol* 2005;**12**:1007–9.
- Tamboli P, Amin MB, Schultz DS, *et al*. Comparative analysis of the nuclear proliferative index (Ki-67) in benign prostate, prostatic intraepithelial neoplasia, and prostatic carcinoma. *Mod Pathol* 1996;**9**:1015–19.
- Hausler O, Epstein JI, Amin MB, *et al*. Cell proliferation, apoptosis, oncogene, and tumor suppressor gene status in adenosis with comparison to benign prostatic hyperplasia, prostatic intraepithelial neoplasia, and cancer. *Hum Pathol* 1999;**30**:1077–86.
- Bostwick DG, Braver MK. Prostatic intra-epithelial neoplasia and early invasion in prostate cancer. *Cancer* 1987;**59**:788–94.
- Jiang Z, Woda BA, Wu CL, *et al*. Discovery and clinical application of a novel prostate cancer marker: alpha-methylacyl CoA racemase (P504S). *Am J Clin Pathol* 2004;**122**:275–89.
- Cunha GR, Hayward SW, Wang YZ. Role of stroma in carcinogenesis of the prostate. *Differentiation* 2002;**70**:473–85.
- McCarthy RP, Zhang S, Bostwick DG, *et al*. Molecular genetic evidence for different clonal origins of epithelial and stromal components of phyllodes tumor of the prostate. *Am J Pathol* 2004;**165**:1395–400.

Squamous differentiation in primary urothelial carcinoma of the urinary tract as seen by MAC387 immunohistochemistry

Antonio Lopez-Beltran, Maria J Requena, Jose Alvarez-Kindelan, Ana Quintero, Ana Blanca, Rodolfo Montironi

J Clin Pathol 2007;**60**:332–335. doi: 10.1136/jcp.2006.038802

Squamous differentiation (SqD) is variably present in urinary tract tumours, but its significance remains unclear. In this study, SqD was assessed by immunohistochemistry using the monoclonal antibody Mac387 in 145 urothelial tumours (bladder, n=115; renal pelvis, n=30). Mac387 detects the myelomonocytic L1 antigen; a member of the calgranulin family shared by epithelial cells and keratinocytes. L1 antigen was shown in SqD in urothelial carcinomas of the bladder or the renal pelvis, including 11 cases with focal SqD unrecognised by

conventional analysis. SqD is more frequent in renal pelvic tumours (p=0.027) and increases with grade/stage mainly in bladder carcinoma (grade, p=0.05; stage, p=0.005). Stage Ta/T1 bladder carcinomas with SqD recurred more (p=0.021). In conclusion, Mac387 efficiently shows SqD in urothelial tumours.

Abbreviations: SqD, squamous differentiation

Urothelial tumours have a great capacity for divergent differentiation and, it is not uncommon to see areas of squamous differentiation (SqD).¹⁻⁵ True incidence is unknown because no strict criteria were defined previously, but some reports suggest that SqD occurs in 16–25% of urothelial carcinomas of the bladder^{1 3 5} and varies according to grade and stage.² The clinical significance of SqD remains uncertain, but seems to be an unfavourable prognostic feature predicting local recurrence after cystectomy.⁶ Also, it might be an indicator of a poor response to radiation therapy or systemic chemotherapy.^{1 7} Information concerning its effect on the recurrence capacity in low-stage bladder urothelial carcinomas is lacking.

Immunohistochemistry might be helpful in the assessment of SqD, and some reports suggest that cytokeratin 14, cytokeratin 19, cytokeratin 10, caveolin 1 or the myelomonocytic L1 antigen could be immunohistochemical markers of SqD.⁸⁻¹²

Our purpose was to evaluate SqD in urothelial tumours as seen by the immunohistochemical detection of L1 antigen in a series of 145 patients with bladder (n = 115) or renal pelvis (n = 30) tumours.

MATERIALS AND METHODS

The retrospective study included specimens from 145 randomly selected patients having primary renal pelvis (n = 30) or bladder (n = 115) urothelial carcinoma retrieved from the files of the Reina Sofia University Hospital, Cordoba, Spain. Treatment included complete transurethral resection of the bladder or cystectomy. Clinical information was obtained from the patient's record. Patient follow-up (mean (SD)), calculated as the number of months from the date of surgery to the date of the most recent cystoscopy (or the last visit or death) was 65 (28) (range 6–120) months. Tumour recurrence in stage Ta/T1 bladder tumours was defined as reappearance of tumour after treatment with at least one tumour-free cystoscopy interval.

Specimens were fixed in formalin and embedded in paraffin. Available H&E stained slides of tumours were reassessed without knowledge of the clinical status. The resulting grade and stage were in accordance with the criteria of the 1973 World Health Organization and the 2002 Tumour, Node and Metastasis, respectively.^{2 13}

Figure 1 Trigone mucosa (A) with suprabasal immunostaining using the L1 antigen (B; A, H&E $\times 200$; B, Mac387 monoclonal antibody $\times 200$). Squamous differentiation (SqD) in urothelial carcinoma showing extensive corneous pearls (C), early corneous pearl formation with dyskeratosis (D), koilocytotic appearance (E), intercellular bridges (F), clear cell features (G) and intracytoplasmic lumina (H; A, H&E $\times 100$; B, C, D, H&E $\times 200$; G, H&E $\times 100$; H, H&E $\times 200$). Diffuse immunostaining of SqD using the L1 antigen in large nests (I), small nests (J), individual cells (K) and intracytoplasmic lumina (L) (I $\times 100$; J, K, L $\times 200$; Mac387 monoclonal antibody).

Table 1 Clinical and pathological variables in 145 cases of urothelial carcinoma of the urinary tract according to squamous differentiation

	Renal pelvis		*p Value	Urinary bladder		*p Value
	Overall	Squamous differentiation		Overall	Squamous differentiation	
Number of patients	30	13 (43.3%)	—	115	24 (21%)	0.027
Age (years; mean (SD))		61.3 (12.7)			67 (9.8)	—
Sex			0.660			0.362
Male	23	10 (43.5%)		95	18 (18.9%)	
Female	7	3 (42.9%)		20	6 (30%)	
Pathological grade			0.439			0.05
I	2	0 (0%)		26	1 (3.8%)	
II	11	5 (45.4%)		41	10 (24.4%)	
III	17	8 (47%)		48	13 (27.1%)	
Pathological stage			0.106			0.005
Ta	—	—		8	— (0%)	
T1	2	— (0%)		65	12 (18.5%)	
T2	10	2 (20%)		37	8 (21.6%)	
T3	15	9 (60%)		5	4 (80%)	
T4	3	2 (67%)		—	—	
Survival status			0.273			0.169
Alive	12	4 (33.3%)		89	14 (15.7%)	
Dead	18	9 (50%)		26	10 (38.4%)	
†Recurrence status			—			0.021
Yes	—	—		18	10 (55.5%)	
No	—	—		55	2 (3.6%)	

* χ^2 test.

†Recurrence status in superficial (stage Ta or T1) bladder urothelial carcinoma (n=73).

The presence of intercellular bridges or keratinisation was indicative of SqD.¹⁻⁵ For immunohistochemistry, 4- μ m-thick sections were deparaffinised in xylene and rehydrated in graded alcohol. Immunohistochemical analysis was carried out using streptavidin–biotin peroxidase. Briefly, sections were incubated overnight with the primary mouse monoclonal antibody detecting myelomonocytic L1 antigen at 4°C (Mac387, Dakopatts, Glostrup, Denmark) diluted at 1:400. Biotinylated secondary antibody and the streptavidin–biotin peroxidase complex were sequentially applied (LSAB2 kit, Dako, Glostrup, Denmark). Diaminobenzidine was used as chromogen (Dakopatts). Endogenous peroxidase activity was blocked by incubating the slides in 1% hydrogen peroxide in methanol for 10 min. For epitope retrieval, the slides were boiled in 10 mM citrate buffer (pH 6.0). Negative and positive controls were included in every procedure. A control group for the immunoreactivity included normal urothelium (n = 5), flat urothelial hyperplasia (n = 5) and normal trigone mucosa (n = 5).

Bivariate and multivariate statistical analyses included the χ^2 test with the SPSS 8.0 for Windows Software. All were two sided tests. A p value ≤ 0.05 was considered significant.

RESULTS

The morphological patterns of keratinising and non-keratinising SqD were: (i) wide nests and cords showing diskeratosis, intercellular bridges and corneous pearls; (ii) small foci often overcome under conventional evaluation showing intercellular bridges, but not diskeratosis or corneous pearls; (iii) additional features suggestive of koilocytosis; (iv) occasional but focally prominent intracytoplasmic lumina in nests of SqD and (v) rare cases had rather clear cells forming nests of variable size well demarcated from concomitant urothelial carcinoma (fig 1).

By immunohistochemistry, the suprabasal cells were labelled in normal trigone mucosa, whereas normal or hyperplastic urothelium and all urothelial carcinomas were negative. Keratinising SqD and the non-keratinising type of SqD, both in tumours of the bladder and the renal pelvis, showed strong homogeneous labelling for the L1 antigen (fig 1) and the intensity of Mac387 staining was dependent on the degree of SqD rather than pathological grade. Mac387 positively stained macrophages, with granulocytes acting as internal control. SqD was more

frequent in renal pelvic than in bladder tumours (p = 0.027). Table 1 shows the demographic features of the series; SqD was seen only in grade 2 and 3 renal pelvic tumours (p = 0.439), but increased from grade 1 to 3 in bladder tumours (p = 0.05). SqD also increased according to stage in the renal pelvis (p = 0.106) and bladder tumours (p = 0.005). Survival status was not related to SqD in both renal pelvis and bladder tumours; patients having stage Ta/T1 bladder tumours with SqD had more chances of recurrence (p = 0.021; table 1). Sensitivity and specificity for SqD using Mac387 reached 100%, while conventional evaluation reached a sensitivity of 54% and 100% specificity. Concerning the clinical significance of SqD in stage T1 bladder cancer, 10 cases with SqD recurred (55.5% of 18 stage T1 recurrent cases) and 2 additional cases with SqD had no recurrences (3.6% of 55 stage T1 non-recurrent cases). In seven stage-T1 cases Mac387 immunohistochemistry helped to delineate SqD that was not already evident with conventional evaluation.

DISCUSSION

Clinical significance of SqD in urothelial tumours remains unsettled with evidence suggesting that it might be an indicator of poor response after radical surgery, radiation or systemic chemotherapy.¹⁻⁷ Most reports to date have concentrated on bladder urothelial tumours showing SqD, but evidence concerning renal pelvis tumours is lacking. Also, the significance of SqD in low-stage bladder tumours in terms of tumour recurrence is unknown. This is in part owing to the fact that SqD in low-stage tumours is often non-keratinising and focal; these foci are often missed during conventional evaluation. The use of immunohistochemistry in assessing SqD is helpful in detecting foci of non-keratinising SqD thus providing an objective identification of the lesion. Our results showed that SqD increased with grade and stage both in the bladder and renal pelvis specimens. Similar results were obtained in earlier studies by Lopez-Beltran *et al*³ and more recently by Sanchez-Carbayo *et al*¹² who also found a significant association between SqD and tumour grade and stage. The absence of differences in renal pelvic tumours might be due to the limited number included since all cases that had SqD were grade 2 or 3 showing similar a incidence. There was no association between SqD and survival status in both renal pelvis and bladder cancer

Take-home messages

- Squamous differentiation (SqD) is more frequent in renal pelvic urothelial tumours and increases with grade/stage.
- The clinical significance of SqD remains uncertain, but seems to be an unfavourable prognostic feature.
- Patients having stage Ta/T1 bladder tumours with SqD had more chance of recurrence.
- Mac387 detects the myelomonocytic L1 antigen; a member of the calgranulin family shared by epithelial cells and keratinocytes.
- Sensitivity and specificity for SqD using Mac387 immunohistochemistry is high.

suggesting that its effect on survival is probably related to the association with higher grade and stage. An original finding in our study is the association with high recurrent tumours in patients with bladder Ta/T1 carcinomas having SqD, which favours SqD as a marker of aggressiveness. In this study, we chose myelomonocytic L1 antigen (a member of the cytoplasmic calcium binding proteins known as calgranulins) expression identified by the monoclonal antibody Mac387, which consistently labelled SqD of all types and allowed more objective identification of SqD than that obtained with routine investigations in some cases.⁸⁻⁹ Our results indicate that Mac387 is also seen in non-keratinised squamous cells, a fact which is corroborated by staining patterns of the non-keratinised mucosa of the bladder trigone.⁹ Our study suggests that Mac387 may be a reliable marker for studying the development of SqD in the urothelium and its tumours where many of the early or partial lesions can be difficult to detect morphologically. This may be due to some forms of SqD being focal or extensive and involving a subjective element in their identification, unlike the typical keratinising types which are easily recognised.

Other valid alternative markers for SqD at the immunohistochemical level include the detection of cytokeratin 14 or cytokeratin 19.¹⁰⁻¹² Recent molecular genetic analyses suggest a characteristic signature of urothelial tumours having SqD; caveolin 1 and keratin 10 have been found to be overexpressed in urothelial carcinomas and cell lines, but more frequently in invasive T2-T4 carcinomas; data on sensitivity and specificity of these markers are not readily available for comparing with Mac387, which seems to be superior to conventional evaluations in our series. In conclusion, our study emphasises the value of immunohistochemistry in assessing SqD in urothelial tumours; in particular, detecting the L1 antigen might result useful in assessing the clinical significance of SqD.

ACKNOWLEDGEMENTS

This work is supported by the Grant FIS 03/0952 (Madrid, Spain).

Authors' affiliations

Antonio Lopez-Beltran, Unit of Anatomic Pathology, Cordoba University Medical School and Reina Sofia University Hospital, Cordoba, Spain

Maria J Requena, Jose Alvarez-Kindelan, Urology service, Reina Sofia University Hospital, Cordoba, Spain

Ana Quintero, Ana Blanca, Biomedical Research Unit, Reina Sofia University Hospital, Cordoba, Spain

Rodolfo Montironi, Institute of Pathological Anatomy and Histopathology, Polytechnic University of the Marche Region, Ancona, Italy

Competing interests: None.

Correspondence to: Professor A Lopez-Beltran, Unit of Anatomical Pathology, Avda Menendez Pidal s/n, E-14004 Cordoba, Spain; em1loboa@uco.es

Accepted 3 July 2006

Published Online First 1 August 2006

REFERENCES

- 1 **Martin JE**, Jenkins BJ, Zuk RJ, *et al*. Clinical importance of squamous metaplasia in invasive transitional cell carcinoma of the bladder. *J Clin Pathol* 1989;**42**:250-3.
- 2 Mostofi KF, ed. In: *Histological typing of urinary bladder tumours. WHO international histological classification of tumours*. Geneva: WHO, 1973.
- 3 **Lopez-Beltran A**, Martin J, Garcia J, *et al*. Squamous and glandular differentiation in urothelial bladder carcinomas. Histopathology, histochemistry and immunohistochemical expression of carcinoembryonic antigen. *Histol Histopathol* 1988;**3**:63-8.
- 4 **Sakamoto N**, Tsuneyoshi M, Enjoji M. Urinary bladder carcinoma with a neoplastic squamous component: a mapping study of 31 cases. *Histopathology* 1992;**21**:135-41.
- 5 **Lopez-Beltran A**, Sauter G, Gasser T, *et al*. Urothelial tumors: infiltrating urothelial carcinoma. In: Eble JN, Sauter G, Epstein JI, Sesterhenn I, eds. *World Health Organization classification of tumors. Pathology and genetics of tumors of the urinary system and male genital organs*. Lyon: IARC press, 2004.
- 6 **Honma I**, Masumori N, Sato E, *et al*. Local recurrence after radical cystectomy for invasive bladder cancer: an analysis of predictive factors. *Urology* 2004;**64**:744-8.
- 7 **Logothetis CJ**, Dexueus FH, Chong C, *et al*. Cisplatin, cyclophosphamide and doxorubicin chemotherapy for unresectable urothelial tumors: the MD Anderson experience. *J Urol* 1989;**141**:33-7.
- 8 **Clark BR**, Kelly SE, Fleming S. Calgranulin expression and association with the keratinocyte cytoskeleton. *J Pathol* 1990;**160**:25-30.
- 9 **Tunçekar MF**, Heryet A, Gatter KC. The L1 antigen and squamous metaplasia in the bladder. *Histopathology* 1991;**19**:245-50.
- 10 **Harnden P**, Southgate J. Cytokeratin 14 as marker of squamous differentiation in transitional cell carcinomas. *J Clin Pathol* 1997;**50**:1032-3.
- 11 **Sanchez-Carbajo M**, Sacci ND, Charytonowicz E, *et al*. Molecular profiling of bladder cancer using cDNA microarrays: defining histogenesis and biological phenotypes. *Cancer Res* 2002;**62**:6973-80.
- 12 **Celis JE**, Celis P, Ossegaard M, *et al*. Proteomics and immunohistochemistry define some of the steps involved in the squamous differentiation of the bladder transitional epithelium: a novel strategy for identifying metaplastic lesions. *Cancer Res* 1999;**59**:3003-9.
- 13 **Greene FL**, Page D, Fleming ID, *et al*. *Morrow M: AJCC cancer staging manual*. New York: Springer-Verlag, 2002.

PostScript

BOOK REVIEWS

Fine needle aspiration cytology: diagnostic principles and dilemmas

Gabrijela Kocjan. New York: Springer, 2005, \$189, pp 236. ISBN 354025 639 3

Gabrijela Kocjan's *Fine needle aspiration cytology: diagnostic principles and dilemmas* is reminiscent

of the type of book (For readers of a certain vintage, and this reviewer, Papworth's "A Primer of Medicine" will come to mind.) that used to be sought out during the interminable long weekend before taking a higher examination: a concise volume, devoid of irrelevant details, that strengthens one's ability to deal with key issues by providing a logical framework on which to hang accumulated knowledge, and freely shares clinical "pearls and pitfalls" accumulated through long experience. One rather hoped the examiners were flipping through the same book.

This monograph of 236 pages is that sort of work. Unlike the ubiquitous "Atlas and Text of Fine Needle Aspiration Cytology", it represents in many ways an update of the type of apologia of clinical cytology written by Lopes-Cordozo or Söderström in the early days of the specialty. The initial chapters are concerned with the "how and why" of fine needle aspiration cytology. Here, the author wisely sacrifices comprehensiveness to her central purpose of showing the enormous potential of clinical cytopathology when practiced in a certain