the committee for autoimmune serology of the International Autoimmune Hepatitis Group. *J Hepatol* 2004;**41**:677–83.

- 6 Bogdanos DP, Baum H, Günsar F, et al. Extensive homology between the major immunodominant mitochondrial antigen in primary biliary cirrhosis and *Helicobacter pylori* does not lead to immunological cross-reactivity. *Scand J Gastroenterol* 2004;**39**:981–7.
- 7 Bogdanos DP, Baum H, Okamoto M, et al. Primary biliary cirrhosis is characterized by IgG3 antibodies cross-reactive with the major mitochondrial autoepitope and its *Lactobacillus* mimic. *Hepatology* 2005;**42**:458–65.

Enteropathy type T cell lymphoma with an unusually late relapse: a case report

Enteropathy type T cell lymphoma (ETTL) is a rare subtype of T cell lymphoma that is strongly associated with coeliac disease.¹ The overall prognosis is poor. In one study, 15 of 19 (79%) patients relapsed within 60 months after the initial diagnosis (median, 6 months), and the overall survival rate at 5 years was 24%.² We describe a case of ETTL with an unusually late relapse. The patient was a 50-year-old woman who developed an ulcerated gastric tumour in 1992, with no associated lymphadenopathy or hepatosplenomegaly. A partial gastrectomy was performed, and at the time of surgery, the tumour was found grossly infiltrating into the peritoneum, posterior rectus sheath and the liver. The pathological diagnosis was diffuse large cell lymphoma of mature T cell lineage (fig 1). On immunohistochemical analysis, the neoplastic cells were found to be CD3+, CD4–, CD5–, CD7+/– and CD8–, and positive for CD56, TIA 1 and granzyme B. She received a full course of VACOP-B (etoposide, doxorubicin, cyclophosphamide, vincristine, prednisone,

bleomycin). Shortly after the diagnosis of T cell lymphoma was made, she was diagnosed with coeliac disease. In view of this finding, the T cell lymphoma was reclassified as ETTL. She was in complete remission and her gastrointestinal symptoms were under control with a gluten-free diet. In 2003, she experienced symptoms of progressive nasal obstruction. Excisional biopsy of a left nasal mass was performed and the pathological diagnosis was diffuse large cell lymphoma of mature T cell phenotype. The immunohistochemical results from the tumour of 1992 and those of 2003 were identical. Molecular analysis of the T cell receptor γ gene also showed an identical rearrangement between these two neoplasms (fig 1), supporting the fact that they were derived from the same neoplastic clone. She was treated with eight cycles of CHOP+R (cyclophosphamide, vincristine, doxorubicin, prednisone, rituxan), followed by additional surgical excision, radiation therapy to the nasal cavity (3600 cGy over 20 fractions) and three cycles of prophylactic intrathecal chemotherapy (methotrexate and cytarabine). After a short remission, she presented with swelling of her right nipple in 2005, which was confirmed to be relapsed T cell lymphoma. She received two cycles of intensive chemotherapy and autologous stem cell transplantation. After a brief remission, she developed jaundice and thrombocytopenia. Circulating malignant cells were found in the peripheral blood (fig 1). She was placed under home palliative care. In summary, our patient represents a highly unusual case of ETTL that relapsed after a long latency time period.

M Al Kaptan

Department of Laboratory Medicine and Pathology,
University of Alberta and Cross Cancer Institute,
Edmonton, Alberta, Canada

D Hui

Department of Medicine, University of Alberta and
Cross Cancer Institute, Edmonton, Alberta, Canada

L Dabbagh

Department of Laboratory Medicine and Pathology,
University of Alberta and Cross Cancer Institute,
Edmonton, Alberta, Canada

T Reiman

Department of Medicine, University of Alberta and
Cross Cancer Institute, Edmonton, Alberta, Canada

R Lai

Department of Laboratory Medicine and Pathology,
University of Alberta and Cross Cancer Institute,
Edmonton, Alberta, Canada

Correspondence to: Dr R Lai, Department of Laboratory
Medicine and Pathology, University of Alberta and
Cross Cancer Institute, 11560 University Avenue,
Edmonton, Alberta, Canada T69 1Z2;
raymond.lai@cancerboard.ab.ca

doi: 10.1136/jcp.2005.036517

Accepted 22 February 2006

Competing interests: None declared.

References

- 1 Isaacson P, Wright D, Ralfkiaer E, et al. Enteropathy-type T-cell lymphoma. In: Jaffe ES, Harris NL, Stein H, Vardiman JW, eds. *WHO classification of tumors*. Lyon: IARC Press, 2001:208–9.
- 2 Gale J, Simmonds PD, Mead GM, et al. Enteropathy-type intestinal T-cell lymphoma: clinical features and treatment of 31 patients in a single center. *J Clin Oncol* 2000;**18**:795–803.

Anaplastic large cell lymphoma presenting with paraneoplastic pemphigus

Paraneoplastic pemphigus is an autoimmune acantholytic mucocutaneous disorder that rarely accompanies an overt or occult neoplasm. Typical features include painful mucosal erosive lesions and pruritic papulosquamous eruptions that often progress to blisters. The antibody is a unique immunoglobulin G that recognises epidermal proteins. We discuss the case of a 23-year-old woman who presented with anaplastic large cell lymphoma complicated by paraneoplastic pemphigus, rapidly resulting in fatal multiorgan failure.

A 23-year-old woman presented with 1 week of fever and a disseminated erythrodermic rash with desquamation of the hands and feet, and erosions in the mouth. A local doctor started treatment with ciprofloxacin and prednisolone.

A diagnosis of toxic epidermal necrolysis was made. On day 4, she required intubation, ventilation and transfer to a tertiary referral centre. A skin punch biopsy showed features of Staphylococcal toxic shock syndrome.

By day 9, multiorgan failure developed and a laparotomy was performed for acute abdominal distension. Retroperitoneal lymphadenopathy, hepatomegaly and splenomegaly were noted, and splenectomy and lymph node biopsy performed.

The histological examination of the spleen and splenic hilar lymph node showed replacement by a nodular infiltrate of small lymphoid cells with vesicular nuclei, multiple nucleoli and a moderate amount of eosinophilic

Figure 1 Left upper panel: gastric enteropathy type T cell lymphoma diagnosed in 1992, composed of a monotonous population of lymphoid cells with a medium cell size, a moderate amount of pale-staining cytoplasm and a round vesicular nucleus. Left lower panel: a circulating large lymphoma cell with a few azurophilic granules in the cytoplasm. Right panel: molecular studies showed that both the gastric and the relapsed nasal tumour had an identical T cell receptor gene arrangement, TCR γ 2, detectable using an automated gene sequencer (AB31300, Applied Biosystems, Foster City, California, USA).

Figure 1 Punch biopsy specimen of skin displaying confluent parakeratosis with underlying suprabasilar clefting and acantholysis. Haematoxylin and eosin staining, $\times 20$.

cytoplasm. Occasional cells with kidney bean-shaped nuclei (hallmark cells) were seen.

Immunohistochemical stains showed membrane and Golgi positivity for CD30 and nuclear, nucleolar and cytoplasmic staining for anaplastic lymphoma kinase protein. A diagnosis of anaplastic large cell lymphoma, small cell variant, anaplastic lymphoma kinase protein positive was made.¹

A bone marrow biopsy showed bone marrow involvement. No cytogenetic abnormality was detected.

Dexamethasone 10 mg four times a day, intravenous and chemotherapy (vincristine 2 mg total dose, adriamycin 35 mg/m², and cyclophosphamide 500 mg/m²) was started on day 10. Granulocyte colony-stimulating factor 5 mcg/kg daily was given.

Figure 2 Postmortem lung specimen displaying segmental destruction of the bronchial walls and occlusion of airway lumen by fresh fibrous tissue. Orcein-Masson, $\times 4$.

On day 10, skin blisters on thighs and forearms developed. Biopsies showed suprabasilar blisters with a lichenoid tissue reaction pattern seen adjacent to the blisters (fig 1). Direct immunofluorescence and indirect immunofluorescence, using rat bladder and monkey oesophagus epithelium, were also negative. A diagnosis of paraneoplastic pemphigus was made.

Despite initial improvement, she subsequently developed septic shock and died on day 19 of admission.

At autopsy, extensive erythroderma and desquamation and loosening of the hair and nails were noted. Residual necrotic tumour was seen in many lymph node groups and in the lungs. The lungs showed focal bronchopneumonia and changes of early bronchiolitis obliterans (fig 2).

Paraneoplastic pemphigus, an autoimmune mucocutaneous disorder with an associated neoplasm and unique immunoglobulin G autoantibody, was first described in 1990.² Most of reported cases are linked to non-Hodgkin's lymphoma, chronic lymphocytic leukaemia or Castleman's disease; characteristically, antibodies to the epidermal proteins desmoglein 1 and 3, and proteins of the plakin family are seen.^{2,3} Pulmonary epithelial damage causing shedding of epithelial cells leading to bronchiolitis obliterans, which results in progressive respiratory failure and often death, are features of paraneoplastic pemphigus.⁴ Respiratory complications occur in approximately 20% of patients, many of whom have circulating antibodies to the respiratory epithelium.⁵ Recent studies suggest that paraneoplastic pemphigus is a heterologous autoimmune syndrome which involves several internal organs in addition to at least five different clinical and immunopathological mucocutaneous variants, and the term "paraneoplastic autoimmune multiorgan syndrome" has been suggested.⁵ In this patient, there was evidence of paraneoplastic autoimmune multiorgan syndrome. This case highlights the need for increased awareness and early recognition of this uncommon erythrodermic condition, paraneoplastic pemphigus, which should prompt timely investigation for an associated, often haematological, malignancy.

A K Davis, M Cole-Sinclair, P Russell
The Alfred Hospital, Melbourne, Australia

Correspondence to: Dr A K Davis, The Alfred Hospital, Commercial Road, Prahran 3181, Melbourne, Victoria, Australia; mandykDavis@hotmail.com

doi: 10.1136/jcp.2005.036137

Accepted 9 April 2006

Competing interests: None declared.

References

- 1 Jaffe E. World Health Organization classification of tumours—tumours of haematopoietic and lymphoid tissues. Lyon: IARC Press, 2001.
- 2 Anhalt GJ, Kim SC, Stanley JR, et al. Paraneoplastic pemphigus. An autoimmune mucocutaneous disease associated with neoplasia. *N Engl J Med* 1990;**323**:1729–35.
- 3 Anhalt GJ. Paraneoplastic pemphigus: the role of tumours and drugs. *Br J Dermatol* 2001;**144**:1102–4.
- 4 Noursari HC, Deterding R, Wojtczak H, et al. The mechanism of respiratory failure in paraneoplastic pemphigus. *N Engl J Med* 1999;**340**:1406–10.

5 Nguyen VT, Ndoye A, Bassler KD, et al. Classification, clinical manifestations, and immunopathological mechanisms of the epithelial variant of paraneoplastic autoimmune multiorgan syndrome: a reappraisal of paraneoplastic pemphigus. *Arch Dermatol* 2001;137:193–206.

Multiple myeloma uncovered by investigating a negative serum iron level

A 67-year-old woman had left nostril obstruction for several weeks, which was thought to be due to a simple nasal polyp. Routine laboratory tests showed anaemia (haemoglobin 7.2 g/l (reference range 11.6–14.0)) and a negative serum iron level ($-68.2 \mu\text{mol/l}$ (7.2–26.9)) was detected with an automatic analyser (Hitachi 7600, Hitachi, Tokyo, Japan). Interference in the direct method of iron measurement was suspected and an investigation of the abnormal absorbance pattern (fig 1) was conducted. We combined the patient's serum with pooled serum at dilution factors 2, 3, 5 and 10, and measured the iron levels. The iron levels were -1.8 , 3.4 , 5.4 and $5.4 \mu\text{mol/l}$, respectively, confirming that the interference was diluted by the pooled serum. Monoclonal immunoglobulins (Igs) can interfere with iron measurement.¹ In our patient, we found significantly high concentrations of IgG (85 g/l (7–16)) and β_2 -microglobulin (3.9 mg/l (1.0–2.7)), combined with low concentrations of IgA (0.3 g/l (0.7–4.0)) and IgM (0.2 g/l (0.4–2.3)). A monoclonal IgG- λ was shown

by immunofixation, and a presumptive diagnosis of multiple myeloma was made. The polypoid mass blocking the patient's left nostril was removed and proved to be an extramedullary plasmacytoma by histopathological examination. Moreover, bone marrow examination showed increasing numbers of plasmacytoid cells (69%), confirming a diagnosis of multiple myeloma.

Blood specimens from patients with multiple myeloma are known to generate erroneous laboratory data through a variety of mechanisms, and monoclonal IgM is more often responsible than monoclonal IgG in this regard.^{2–4} Monoclonal immunoglobulins have often been reported to cause positive interference with direct iron assays,^{1–5} but to our knowledge negative interference in serum iron measurement due to monoclonal Igs has not been reported previously in the English literature. Recent improvement of the reagent buffers used in serum iron tests has considerably decreased the interference. However, the present case shows that such interference may still take place. A negative iron value is usually due to EDTA contamination of sera, which can be suspected on finding low calcium and high potassium concentrations.⁶ However, in our case, serum calcium and potassium levels were normal, so we ruled out the possibility of EDTA contamination. We were extremely satisfied to uncover a hitherto undiagnosed multiple myeloma in this patient. We recommend that laboratory professionals be vigilant whenever confronted with unrealistic test results, and the

possibility of interfering substances (such as drugs, heterophilic antibodies and paraproteins) must be thoroughly investigated.

L C Wu, S S Chuang, C N Lin

Department of Pathology, Chi-Mei Foundation Medical Center, Taipei, Taiwan

C C Su

Departments of Pathology and Anatomic Pathology Clinical, Buddhist Dalin Tzu Chi General Hospital, Chiayi, Taiwan; Department of Pathology, Tzu Chi University, Tainan, Taiwan

M P Lin

Department of Radiology, Chi-Mei Foundation Medical Center, Tainan, Taiwan

Correspondence to: Dr C C Su, Department of Clinical Pathology, Buddhist Dalin Tzu Chi General Hospital, 2, Minsheng Road, Dalin Town, Chiayi County 622, Taiwan; sucupo@yahoo.com.tw

This study was approved by the institutional review board of our hospital.

doi: 10.1136/jcp.2005.033308

Accepted 15 March 2006

Funding: This study was supported by grant DTCRD94(2)-08 from the Buddhist Dalin Tzu Chi General Hospital, Chiayi, Taiwan.

References

- 1 Dorizzi R, Battaglia P, Lora A. Iron measurement in patients with monoclonal immunoglobulin: a further caution. *Clin Chem* 1991;37:589–90.
- 2 Gulian JM, Dalmasso C, Pesquie M, et al. Abnormal bilirubin binding to human serum albumin in a patient with unusual myeloma immunoglobulin G. *Eur J Clin Chem Clin Biochem* 1993;31:115–19.
- 3 John R, Olesky D, Issa B, et al. Pseudohypercalcaemia in two patients with IgM paraproteinaemia. *Ann Clin Biochem* 1997;34:694–6.
- 4 Yu A, Pira U. False increase in serum C-reactive protein caused by monoclonal IgM-lambda: a case report. *Clin Chem Lab Med* 2001;39:983–7.
- 5 Bakker AJ. Influence of monoclonal immunoglobulins in direct determinations of iron in serum. *Clin Chem* 1991;37:690–4.
- 6 Imafuku Y, Meguro S, Kanno K, et al. The effect of EDTA contaminated in sera on laboratory data. *Clin Chim Acta* 2002;325:105–11.

Figure 1 (A) The absorbance (Abs) pattern for determination of iron in a patient with a serum iron concentration (Cons) of $40.8 \mu\text{mol/l}$ ($228 \mu\text{g/dl}$) was normal. The patient's serum was initially admixed with the primary reagent. The absorbance was negative (around -105). When the secondary reagent was added immediately after the 16th detection, the absorbance sharply increased to 550. (B) The iron absorbance pattern of this patient was abnormal (iron concentration $-68.2 \mu\text{mol/l}$ ($-381 \mu\text{g/dl}$)). In the beginning, the absorbance was positive (around 100) and decreased gradually. After the secondary reagent was added, the absorbance further decreased more quickly.

Vulval intraepithelial neoplasia with extra mammary pagets disease: a rare association

Vulval intraepithelial neoplasia (VIN) is the term used for squamous intraepithelial lesions of the vulva, and is classified based on the thickness of epithelium showing dysplasia. The incidence of VIN has nearly doubled over the past two decades and is becoming more frequent in young women aged 20–35 years.¹ Extra mammary Paget's disease comprises $\leq 2\%$ of primary vulval neoplasms, and extra mammary Paget's disease arising in other sites is even rarer. Incidence of an underlying neoplasm detected in cases of vulval extra mammary Paget's disease ranges from 0% to 30% in various studies.² However, it is extremely rare to encounter VIN with extra mammary Paget's disease in the vulva. We present one such rare and interesting association of these two diseases.

A 65-year-old woman presented with whitish plaques on the vulva, along with