5 Nguyen VT, Ndoye A, Bassler KD, et al. Classification, clinical manifestations, and immunopathological mechanisms of the epithelial variant of paraneoplastic autoimmune multiorgan syndrome: a reappraisal of paraneoplastic pemphigus. *Arch Dermatol* 2001;137:193–206.

Multiple myeloma uncovered by investigating a negative serum iron level

A 67-year-old woman had left nostril obstruction for several weeks, which was thought to be due to a simple nasal polyp. Routine laboratory tests showed anaemia (haemoglobin 7.2 g/l (reference range 11.6–14.0)) and a negative serum iron level ($-68.2 \mu\text{mol/l}$ (7.2–26.9)) was detected with an automatic analyser (Hitachi 7600, Hitachi, Tokyo, Japan). Interference in the direct method of iron measurement was suspected and an investigation of the abnormal absorbance pattern (fig 1) was conducted. We combined the patient's serum with pooled serum at dilution factors 2, 3, 5 and 10, and measured the iron levels. The iron levels were -1.8 , 3.4 , 5.4 and $5.4 \mu\text{mol/l}$, respectively, confirming that the interference was diluted by the pooled serum. Monoclonal immunoglobulins (Igs) can interfere with iron measurement.¹ In our patient, we found significantly high concentrations of IgG (85 g/l (7–16)) and β_2 -microglobulin (3.9 mg/l (1.0–2.7)), combined with low concentrations of IgA (0.3 g/l (0.7–4.0)) and IgM (0.2 g/l (0.4–2.3)). A monoclonal IgG- λ was shown

by immunofixation, and a presumptive diagnosis of multiple myeloma was made. The polypoid mass blocking the patient's left nostril was removed and proved to be an extramedullary plasmacytoma by histopathological examination. Moreover, bone marrow examination showed increasing numbers of plasmacytoid cells (69%), confirming a diagnosis of multiple myeloma.

Blood specimens from patients with multiple myeloma are known to generate erroneous laboratory data through a variety of mechanisms, and monoclonal IgM is more often responsible than monoclonal IgG in this regard.^{2–4} Monoclonal immunoglobulins have often been reported to cause positive interference with direct iron assays,^{1–5} but to our knowledge negative interference in serum iron measurement due to monoclonal Igs has not been reported previously in the English literature. Recent improvement of the reagent buffers used in serum iron tests has considerably decreased the interference. However, the present case shows that such interference may still take place. A negative iron value is usually due to EDTA contamination of sera, which can be suspected on finding low calcium and high potassium concentrations.⁶ However, in our case, serum calcium and potassium levels were normal, so we ruled out the possibility of EDTA contamination. We were extremely satisfied to uncover a hitherto undiagnosed multiple myeloma in this patient. We recommend that laboratory professionals be vigilant whenever confronted with unrealistic test results, and the

possibility of interfering substances (such as drugs, heterophilic antibodies and paraproteins) must be thoroughly investigated.

L C Wu, S S Chuang, C N Lin

Department of Pathology, Chi-Mei Foundation Medical Center, Taipei, Taiwan

C C Su

Departments of Pathology and Anatomic Pathology Clinical, Buddhist Dalin Tzu Chi General Hospital, Chiayi, Taiwan; Department of Pathology, Tzu Chi University, Tainan, Taiwan

M P Lin

Department of Radiology, Chi-Mei Foundation Medical Center, Tainan, Taiwan

Correspondence to: Dr C C Su, Department of Clinical Pathology, Buddhist Dalin Tzu Chi General Hospital, 2, Minsheng Road, Dalin Town, Chiayi County 622, Taiwan; sucupo@yahoo.com.tw

This study was approved by the institutional review board of our hospital.

doi: 10.1136/jcp.2005.033308

Accepted 15 March 2006

Funding: This study was supported by grant DTCRD94(2)-08 from the Buddhist Dalin Tzu Chi General Hospital, Chiayi, Taiwan.

References

- 1 Dorizzi R, Battaglia P, Lora A. Iron measurement in patients with monoclonal immunoglobulin: a further caution. *Clin Chem* 1991;37:589–90.
- 2 Gulian JM, Dalmasso C, Pesquie M, et al. Abnormal bilirubin binding to human serum albumin in a patient with unusual myeloma immunoglobulin G. *Eur J Clin Chem Clin Biochem* 1993;31:115–19.
- 3 John R, Olesky D, Issa B, et al. Pseudohypercalcaemia in two patients with IgM paraproteinaemia. *Ann Clin Biochem* 1997;34:694–6.
- 4 Yu A, Pira U. False increase in serum C-reactive protein caused by monoclonal IgM-lambda: a case report. *Clin Chem Lab Med* 2001;39:983–7.
- 5 Bakker AJ. Influence of monoclonal immunoglobulins in direct determinations of iron in serum. *Clin Chem* 1991;37:690–4.
- 6 Imafuku Y, Meguro S, Kanno K, et al. The effect of EDTA contaminated in sera on laboratory data. *Clin Chim Acta* 2002;325:105–11.

Vulval intraepithelial neoplasia with extra mammary pagets disease: a rare association

Vulval intraepithelial neoplasia (VIN) is the term used for squamous intraepithelial lesions of the vulva, and is classified based on the thickness of epithelium showing dysplasia. The incidence of VIN has nearly doubled over the past two decades and is becoming more frequent in young women aged 20–35 years.¹ Extra mammary Paget's disease comprises $\leq 2\%$ of primary vulval neoplasms, and extra mammary Paget's disease arising in other sites is even rarer. Incidence of an underlying neoplasm detected in cases of vulval extra mammary Paget's disease ranges from 0% to 30% in various studies.² However, it is extremely rare to encounter VIN with extra mammary Paget's disease in the vulva. We present one such rare and interesting association of these two diseases.

A 65-year-old woman presented with whitish plaques on the vulva, along with

Figure 1 (A) The absorbance (Abs) pattern for determination of iron in a patient with a serum iron concentration (Cons) of $40.8 \mu\text{mol/l}$ ($228 \mu\text{g/dl}$) was normal. The patient's serum was initially admixed with the primary reagent. The absorbance was negative (around -105). When the secondary reagent was added immediately after the 16th detection, the absorbance sharply increased to 550. (B) The iron absorbance pattern of this patient was abnormal (iron concentration $-68.2 \mu\text{mol/l}$ ($-381 \mu\text{g/dl}$)). In the beginning, the absorbance was positive (around 100) and decreased gradually. After the secondary reagent was added, the absorbance further decreased more quickly.

Figure 1 Low-power photomicrograph showing vulvar intraepithelial neoplasia III/carcinoma in situ changes.

complaints of pruritus. She had no complaints of any discharge per vaginum, and the lower genital tract was normal on colposcopic examination. A biopsy performed from the lesion showed changes of VIN III/severe dysplasia involving almost the whole epidermal thickness. Subsequently, a simple vulvectomy with a 1 cm free margin was performed. Extensive sampling of the specimen showed the presence of VIN III, along with extra mammary Paget's disease involving the superior or upper margins of the resections. Sections from both the lateral resection limits showed the presence of Paget's cells. However, there was no evidence of intraepidermal dysplasia. Paget's cells were larger than surrounding keratinocytes, with the presence of abundant pale to vacuolated cytoplasm and finely granular to vesicular

nucleus. The cells were present singly and in small groups and were confined to the epidermis only (figs 1–3). No koilocytic change was noted in the epidermal lining. Mammography and ultrasound examination of bilateral breasts also did not show any focal lesion. Paget's cells showed positivity for mucin stains such as periodic acid Schiff and mucicarmine. In addition, immunostaining for CK7 was positive, although that for carcinoembryonic antigen and epithelial membrane antigen was non-contributory.

It is currently believed that most cases of vulvar extra mammary Paget's disease are primary—that is, arising within the epidermis—and few are associated with cutaneous sweat gland tumours. Vulvar extra mammary Paget's disease has also been described in

association with endometrial, endocervical, vaginal, urethral and bladder neoplasms. Occasional cases have also been described in association with breast carcinoma.³ Paget's cells are proposed to originate either from the intraepidermal cells of apocrine gland ducts or from pluripotent keratinocyte stem cells. Cytochemically and immunohistochemically, Paget's cells are always adenocarcinoma cells. Extra mammary Paget's cells show stronger positivity for mucin stains and gross cystic disease fluid protein (GCDFP-15) in comparison to their mammary counterparts. In a study by Helm *et al.*,⁴ negativity for the carcinoembryonic antigen was seen more frequently as the grade of lesion increased or when it was associated with an underlying malignancy. An extensive search of the literature showed only a single patient with vulvar Paget's disease having concomitant squamous cell carcinoma in situ/VIN III changes. The case was reported by Brainard *et al.*,⁵ who studied the changes in squamous epithelium in 11 patients with extra mammary Paget's disease and found associated neoplastic changes in two patients. One patient had an underlying adenocarcinoma whereas the other had concomitant VIN III changes.

The association of vulvar Paget's disease and VIN may be just a chance phenomenon, or there may be a common link between the pathogenesis of these two entities. On histopathological examination, it is not difficult to diagnose these entities. However, any patient with the presence of vulvar Paget's disease should also have a thorough check up for breast lesions. As this is a rare association, prognosis of this associated disease is difficult to ascertain. A thorough follow-up of the patient is recommended.

It is important to realise this entity so that thorough sampling can be carried out to exclude an underlying malignancy. Moreover, patients with primary Paget's disease in nature can be treated by wide excision of the lesion with a 1 cm free margin and regular follow-up.

P Bhatia, A Ahuja, P Dey, V Suri

Department of Pathology and Gynecology, PGIMER, Chandigarh, India

Correspondence to: Dr P Bhatia, 290, O-Block, New Doctors' Hostel, PGIMER, Sector-12, Chandigarh-160012, India; prateekbhatia@rediffmail.com

doi: 10.1136/jcp.2006.038521

Accepted 27 March 2006

Competing interests: None declared.

Written consent has been obtained for the publication of this study.

References

- 1 Andreason B, Bock JE. Intraepithelial neoplasia in the vulvar region. *Gynecology* 1985;21:300.
- 2 J Lloyd, AM Flanagan. Mammary and extra mammary Paget's disease. *J Clin Pathol* 2000;53:742–9.
- 3 Sitalkin C, Ackerman AB. Mammary and extra mammary Paget's disease. *Am J Dermatopathol* 1985;7:335–40.
- 4 Helm KF, Goellner JR, Peters MS. Immunohistochemical stains in extra mammary Paget's disease. *Am J Dermatopathol* 1992;14:402–7.
- 5 Brainard JA, Hart WR. Proliferative epidermal lesions associated with anogenital Paget's disease. *Am J Surg Pathol* 2000;24:543–52.

Figure 2 Low-power photomicrograph showing Paget's cells present in vulvar epithelium.

Figure 3 High-power photomicrograph showing Paget's cells present in small groups and nests. Paget's cells show round to oval nucleus with granular to vesicular chromatin and abundant pale to clear cytoplasm. Adjoining vulvar epithelium shows vulvar intraepithelial neoplasia III changes.

CALENDAR OF EVENTS

Diagnostic histopathology of breast disease

14–17 May 2007, Hammersmith Hospital (Imperial College), London.

A 4-day course on the Hammersmith Campus designed for pathologists at

Consultant and Senior Trainee level and suitable for the final examination for MRCPath. The course will provide a comprehensive coverage of the histopathology of breast disease, with special emphasis on areas which pose diagnostic difficulties. The course is based historically on the extensive referral collection of Professor JG Azzopardi, to which has been added new cases from Charing Cross, Nottingham City and Addenbrooke's Hospitals. The course will be slanted towards practical

diagnostic pathology with a specific session dealing with problems encountered in core biopsies and a special workshop dealing with the interpretation of HER2 immunostaining. The participants will be given ample time to study histological preparations, followed by illustrated discussions of the cases. In addition, there will be several daily talks on specific topics, given by eminent breast pathologists and followed by discussions. The topics will include: new issues in the interpretation of breast biopsies, breast immunohistochemistry, problematic breast lesions, dealing with the gross specimen, fibro-epithelial, spindle cell and papillary lesions, proliferative lesions/carcinoma in-situ, sentinel lymph node biopsy and the impact of molecular genetics on diagnostic breast pathology. The faculty will include Anthony Douglas-Jones, Ian Ellis, Bharat Jassani, Andrew Lee, Sarah Pinder, Jorge Reis-Filho and Sami Shousha.

Further details: Sandra Lock, Department of Histopathology, Hammersmith Hospital, Du Cane Road, London W12 0NN, UK. Tel: +44 (0)20 8383 3292; Fax: +44 (0)20 8740 7417; Email:histopathology.shortcourses@imperial.ac.uk

CORRECTION

doi: 10.1136/jcp.2005.035238 corr1

There was an error in the December issue of the journal (Anton E, Larrañaga O. An uncommon pattern of cardiac invasion in hepatocellular carcinoma HCC, hepatocellular carcinoma. *J Clin Pathol* 2006;**59**:1337. The correct name of the second author is Astiazaran A.