EDITORIAL

Is antibiotic prophylaxis ever necessary before transoesophageal echocardiography?

J B Chambers, J L Klein, S R Bennett, M J Monaghan, J C Roxburgh

Evidence strongly suggests that antibiotic prophylaxis should not be used routinely for transoesophageal echocardiography for any indication

Heart 2006;92:435–436. doi: 10.1136/hrt.2005.077446

The British Cardiac Society (BCS) Clinical Practice Committee and Royal College of Physicians Clinical Effectiveness and Evaluation Unit recently published a guidance document¹ recommending antibiotic prophylaxis routinely before transoesophageal echocardiography (TOE), cardiac surgery and percutaneous coronary intervention in patients at moderate or high risk of endocarditis. This guidance is contrary to that of the European Society of Cardiology² and the American Heart Association.³ The purpose of this editorial is to re-examine the evidence for and against antibiotic prophylaxis in TOE.

EVIDENCE FOR ENDOCARDITIS AFTER TOE

The evidence used for the BCS guidance was a single case report.⁴ A 55 year old man was referred for TOE to assess the need for surgery for bileaflet mitral prolapse. One week later he developed anorexia, nausea, light-headedness and malaise and subsequently fever and myalgia. At 17 days after TOE he sought medical help and *Streptococcus sanguis* was grown in both of two blood cultures. A repeat TOE showed no change other than increased nodularity of the flail portion of the posterior mitral leaflet. The appearance was thought to be consistent with infection or an increase in gain setting. The patient was treated with intravenous penicillin and gentamicin for two weeks followed by oral penicillin for a further two weeks, and at the time of writing had remained well. The authors acknowledged that this case was not conclusively caused by the TOE, but thought that the temporal relationship was suggestive.

Systematic studies have failed to find endocarditis whether^{5–6} or not^{7–8} an apparent bacteraemia was demonstrated at the time of TOE. Patients with replacement heart valves are at particularly high risk of endocarditis. In 85 patients with replacement heart valves followed for a mean of 82 days after TOE,⁹ no case of endocarditis was found.

Thus there is one case report suggesting a possible link between endocarditis and TOE, compared against follow-up studies in a total of 432 patients showing no evidence of endocarditis. Similarly, a review of 41 studies of upper

gastrointestinal endoscopy¹⁰ found two cases of endocarditis, in both of which the connection was thought to be unconvincing.

EVIDENCE FOR BACTERAEMIA AFTER TOE

There have been several studies investigating the induction of bacteraemia by TOE. Gorge *et al*⁵ found that all cultures taken before TOE were sterile, but stopped their study early after four (17%) of the next 24 cultures taken 6–12 minutes after TOE were positive. However, in view of the small numbers, this does not represent a significant rise (Fisher's exact test $p > 0.05$). Secondly, the organisms were mainly skin commensals and thus unlikely to represent true bacteraemia following TOE related trauma. A study of 47 patients¹¹ found two positive cultures before, but six positive after TOE. Again the organisms were probably contaminants, diphtheroids in three, micrococci in two and aerobic spore formers in one, all of which are skin commensals.

By contrast a study of 101 patients showed no positive cultures six minutes after the procedure compared to cultures positive for *Staphylococcus epidermidis* in two patients before the procedure.⁷ The rate of positive cultures before TOE or in control subjects not having TOE ranges between 0.07–6.3%.^{7–8 11–13} A number of studies show that bacteraemia is more common before than after TOE^{8 12 14 15} and whether or not antibiotics are given.¹⁶ The presence of bacteraemia is not related to difficulty of intubation or the presence of an intravenous line.⁸ In nearly all studies, blood cultures have been positive only for skin commensals. This strongly suggests contamination. *Streptococcus viridans* is the most likely precursor of subsequent endocarditis and only one instance of viridans bacteraemia has been demonstrated.⁵

Thus, significant bacteraemia appears to be a rare event after TOE. Considering that some sort of allergic reaction occurs in between 7 and 40 of every 1000 courses of penicillin^{17 18} and that there is little evidence for its benefit in dental and other procedures leading to much higher rates of bacteraemia, its routine use for TOE is inappropriate.

CONCLUSION

Routine antibiotic prophylaxis before TOE has been recommended in only two original publications, one a suggestive but inconclusive case report⁴ and the other a small study reporting a statistically insignificant incidence of bacteraemia likely to be caused by contamination.⁵ By contrast, antibiotic prophylaxis has been thought not to be routinely necessary in 11 studies involving 1063 patients.^{6–9 11–16 19} Of these, eight

See end of article for authors' affiliations

Correspondence to:
Dr John Chambers,
Cardiothoracic Centre, St
Thomas' Hospital, London
SE1 7EH, UK;
jboydchambers@aol.com

Published Online First
30 December 2005

studies considered that antibiotic prophylaxis was never routinely recommended, while three^{6 11 13} suggested that prophylaxis could be considered in individual patients with prosthetic valves or after previous endocarditis.

The evidence points overwhelmingly in favour of the European Society of Cardiology guidance which is not to use antibiotic prophylaxis routinely for transoesophageal echocardiography for any indication. It may be reasonable to consider antibiotics in occasional cases—for example, a patient with a replacement heart valve and evidence of poor oral hygiene in whom the study is being performed for an indication other than suspected endocarditis. In these individual cases, practice is governed by clinical common sense rather than evidence.

Authors' affiliations

J B Chambers, J C Roxburgh, Cardiothoracic Centre, Guy's and St Thomas' Hospitals, London, UK

J L Klein, Department of Infection, Guy's and St Thomas' Hospitals, London, UK

S R Bennett, Department of Anaesthesia, Castle Hill Hospital, Cottingham, West Yorkshire, UK

M J Monaghan, Cardiac Department, King's College Hospital, London, UK

REFERENCES

- 1 Ramsdale DR, Elliott TSJ, Wright P, *et al.* Guidance on the prophylaxis and treatment of infective endocarditis in adults. <http://www.rcplondon.ac.uk/pubs/books/endocarditis/endocarditis.pdf>
- 2 Horstkotte D, Follath F, Gutschik E, *et al.* Guidelines on prevention, diagnosis and treatment of infective endocarditis. *Eur Heart J* 2004;**25**:267–76.
- 3 Ajani AS, Bisno AL, Chung KJ, *et al.* Prevention of bacterial endocarditis. Recommendations by the American Heart Association. *JAMA* 1990;**264**:2919–22.
- 4 Foster E, Kusumoto FM, Sobol SM, *et al.* Streptococcal endocarditis temporally related to transesophageal echocardiography. *J Am Soc Echocardiogr* 1990;**3**:424–7.
- 5 Gorge G, Erbel R, Henrichs KJ, *et al.* Positive blood cultures during transesophageal echocardiography. *Am J Cardiol* 1990;**65**:1404–5.
- 6 Roudaut R, Lartigue MC, Texier-Maugein J, *et al.* Incidence of bacteraemia or fever during transesophageal echocardiography: a prospective study of 82 patients. *Eur Heart J* 1993;**14**:936–40.
- 7 Pongratz G, Henneke K-H, von der Grun M, *et al.* Risk of endocarditis in transesophageal echocardiography. *Am Heart J* 1993;**125**:190–3.
- 8 Melendez LJ, Chan K-L, Cheung PK, *et al.* Incidence of bacteremia in transesophageal echocardiography: a prospective study of 140 consecutive patients. *J Am Coll Cardiol* 1991;**18**:1650–4.
- 9 Gal RA, Gaeckle TC, Gadasalli S, *et al.* Chemoprophylaxis before transesophageal echocardiography in patients with prosthetic or bioprosthetic cardiac valves. *Am Heart J* 1993;**72**:115–7.
- 10 Botoman VA, Surawicz CM. Bacteremia with gastrointestinal endoscopic procedures. *Gastrointest Endosc* 1986;**32**:342–6.
- 11 Dhas KL, Hemalatha R, Umesan CV, *et al.* Prospective evaluation of the risk of bacteremia induced by transesophageal echocardiography. *Indian Heart J* 2002;**54**:181–3.
- 12 Voller H, Spielberg C, Schroder K, *et al.* Frequency of positive blood cultures during transesophageal echocardiography. *Am J Cardiol* 1991;**68**:1538–40.
- 13 Shyu K-G, Hwang J-J, Lin S-C, *et al.* Prospective study of blood culture during transesophageal echocardiography. *Am Heart J* 1992;**124**:1541–4.
- 14 Steckelberg JM, Khandheria BK, Anhalt JP, *et al.* Prospective evaluation of the risk of bacteremia associated with transesophageal echocardiography. *Circulation* 1991;**84**:177–80.
- 15 Lamich R, Alonso C, Guma JR, *et al.* Prospective study of bacteremia during transesophageal echocardiography. *Am Heart J* 1993;**125**:1454–5.
- 16 Mentec H, Vignon P, Terre S, *et al.* Frequency of bacteremia associated with transesophageal echocardiography in intensive care unit patients: a prospective study of 139 patients. *Crit Care Med* 1995;**23**:1194–9.
- 17 Parker CW. Allergic reactions to long-term benzathine penicillin prophylaxis for rheumatic fever. *Lancet* 1991;**337**:1308–10.
- 18 Idsoe O, Guthe T, Wilcox RR, *et al.* Nature and extent of penicillin side-reactions, with particular reference to fatalities from anaphylactic shock. *Bull World Health Organ* 1968;**38**:159–88.
- 19 Nikutta P, Mantey-Stiers F, Becht I, *et al.* Risk of bacteremia induced by transesophageal echocardiography: analysis of 100 consecutive procedures. *J Am Soc Echocardiogr* 1992;**5**:168–72.

IMAGES IN CARDIOLOGY

doi: 10.1136/hrt.2005.072694

Blue-grey cutaneous discolouration secondary to amiodarone treatment

A 78 year old man was admitted to the cardiology department because of atrial fibrillation with rapid ventricular response. He was an inhabitant of a remote rural area and had been treated with amiodarone for five years, but without medical supervision for a long time.

Physical examination revealed a blue-grey discolouration of his face. The laboratory tests revealed a 100 fold rise of his serum aminotransferase concentrations (AST 3175 µl, ALT 2662 µl). The drug regimen was discontinued and long term anticoagulation and digoxin treatment were used. A skin biopsy was performed and fixed in formalin. Paraffin sections were stained with haematoxylin and eosin (H&E) stain, Perl's for haemosiderin, and Masson-Fontana for melanin pigment. The epidermis was normal but scattered aggregates of macrophages containing ample fine golden brown pigment granules negative for haemosiderin and melanin were seen in the dermis with a tendency for perivascular localisation. Single, granule laden histiocytes were also present and there was no evidence of inflammatory response. After amiodarone discontinuation, serum aminotransferase

Left: Hyperpigmentation of the face. Right: Haematoxylin and eosin stain (×200) showing pigment containing histiocytes (arrows) around dermal blood vessels.

values returned to near normal within a week's time. The skin discolouration was still evident, but diminished, at six months' follow up.

S Nikolidakis
Z S Kyriakides
C Barbatis
zskyr@otenet.gr

There is no conflict of interest related to this manuscript

The study complies with current ethical considerations