EDITORIAL

Is calcium the key for the assessment of progression/regression of coronary artery disease?

F Cademartiri

Heart 2006;92:1187–1188. doi: 10.1136/hrt.2006.089573

Coronary calcium scoring has become a common test for cardiovascular risk stratification

Coronary artery calcifications are part of the atherosclerotic degeneration of the arterial vessel wall. Since the introduction of electron-beam computer tomography (EBCT) and the Agaston score,¹ coronary calcium has been investigated for several years and has become a common test for cardiovascular risk stratification in several countries.² However, it is not yet accepted within primary prevention guidelines.

The method is simple and was developed to assess the coronary artery burden of calcium. By means of a non-contrast-enhanced EBCT (or more recently multislice computer tomography (MSCT)) scan, we can quantify the amount of calcium in the coronary tree.

Coronary calcium score obtained with EBCT is equivalent to the one obtained with MSCT.^{3–5} Several studies demonstrated a fair correlation between EBCT and spiral-CT/MSCT. In addition, the reproducibility of MSCT (5–12%) is better than EBCT (14%–51%), depending on scan technique and quantification method.^{5–8}

In this issue of the journal, Houslay *et al*⁹ report a substudy from the SALTIRE trial. They investigated, in 102 patients with aortic valve calcifications, the rate of progression of coronary calcifications for a median period of two years, randomising the patients to treatment with 80 mg of atorvastatin versus placebo. The randomised, controlled study concludes that lipid-lowering drugs do not affect significantly the progression of coronary calcifications.

We know that coronary artery calcifications derive from an active metabolic process similar to that taking place within bone tissue.^{10–12} If coronary calcium is the result of the precipitation of calcium, then it may only represent a marker for atherosclerosis. As an active metabolic process, instead, it may become a unique marker of progression/regression of coronary artery disease.

CONCEPTS INHERENT TO CORONARY CALCIUM IMAGING

The interest of the scientific community in coronary calcium is based on the concept that calcium may be an early and/or more specific marker of coronary atherosclerosis and an indicator of future coronary events. Calcification in coronary artery vessel walls is a

very sensitive marker for the presence of atherosclerosis^{13 14} and shows moderate correlation between its extent and the likelihood of coronary obstruction.^{15 16}

Calcification starts very early in the atherosclerotic process and progresses with the volumetric increase of plaque burden.¹⁷ The macroscopic calcified spot (that is, the minimum amount of calcium that can be visualised by EBCT or MSCT) appears in relatively advanced stages of coronary artery disease.^{17 18} This is the basis of the controversy over whether calcium, as detected by EBCT/MSCT, can be considered an early marker of coronary atherosclerosis.¹⁸

Serial studies have demonstrated that it is possible to induce the regression of coronary stenosis with lipid-lowering drugs and that a small reduction of the degree of stenosis corresponds to a considerable reduction of coronary events.^{19 20} Thus, atherosclerotic plaque stabilisation is independent of the recovery of lumen patency. Given the fact that coronary calcium undergoes an active metabolic process, authors reported some time ago that the administration of lipid-lowering drugs in asymptomatic individuals reduced significantly the progression of coronary calcification.^{21–23}

STUDY LIMITATIONS

The scanner technology employed in the study is not the most up to date, and neither is the scoring method—the latest 16/64 slice MSCT scanners may offer more reliable and reproducible quantification of coronary calcium. However, we should consider the calcium score as a simple method for risk stratification, and therefore not necessarily subject to rapid development.

In addition, even though the Agaston score is the reference standard for coronary calcium score, newer and more reliable methods are available (volume and mass).²⁴ These methods are able to reduce further the room for error and improve reproducibility, therefore allowing a better detection of smaller changes in the score, especially in the set-up of serial studies.

The population, although followed up for two years, is rather small compared to other observational studies.

NEW CONCEPTS IN CORONARY RISK ASSESSMENT

Since 2001, MSCT was able to display a promising capability in detecting not only coronary calcium but also the non-calcific component of

Correspondence to:
Dr Filippo Cademartiri,
Department of Radiology,
Erasmus Medical Center,
Dr Molewaterplein, 40,
3015 GD – Rotterdam,
The Netherlands;
filippocademartiri@
hotmail.com

Published Online First
10 April 2006

Abbreviations: MSCT, multislice computer tomography; EBCT, electron-beam computer tomography

plaques.²⁵ These results were improved with the latest generations of MSCT scanners.^{25–29}

It is expected that MSCT will prove to be useful in the assessment of total coronary plaque burden.³⁰ However, only qualitative and semi-quantitative information have been reported until now. Software developments are required as well as further MSCT scanner improvements (that is, beyond 64 slice) in order to achieve the detection of the so called “vulnerable plaque”.

Calcium score alone, as measured by EBCT/MSCT, does not have the capability of assessing the distribution of various morphologic patterns of calcium and their relation to other “soft” plaque components. While MSCT coronary angiography may extend the potential of non-invasive plaque imaging, several limitations are still present. The most important limitation, in terms of serial coronary plaque burden assessment, is radiation exposure.^{31–32}

CONCLUSION

The study of Houslay *et al*⁹ adds new information to the controversy surrounding the usefulness of calcium scoring for serial evaluation of coronary artery disease progression/regression. An important public health message can be drawn from this study. In some regions of the world patients are subjected to repeated CT scanning to determine if their calcification score is increasing. Given the controversial evidence, this may be a questionable medical practice exposing patients (actually asymptomatic individuals) to unnecessary radiation exposure.

The recent introduction of MSCT coronary angiography has brought high expectations in the evaluation of total coronary plaque burden. However, the value of the parameters that can be measured using this technology has yet to be demonstrated.

Conflict of interest statement: I have no conflicts of interest

REFERENCES

- Agatston AS, Janowitz WR, Hildner FJ, *et al*. Quantification of coronary artery calcium using ultrafast computed tomography. *J Am Coll Cardiol* 1990;**15**:827–32.
- Raggi P. Coronary calcium on electron beam tomography imaging as a surrogate marker of coronary artery disease. *Am J Cardiol* 2001;**87**:27A–34A.
- Becker CR, Jakobs TF, Aydemir S, *et al*. Helical and single-slice conventional CT versus electron beam CT for the quantification of coronary artery calcification. *AJR Am J Roentgenol* 2000;**174**:543–7.
- Becker CR, Kleffel T, Crispin A, *et al*. Coronary artery calcium measurement: agreement of multirow detector and electron beam CT. *AJR Am J Roentgenol* 2001;**176**:1295–8.
- Ohnesorge B, Flohr T, Fischbach R, *et al*. Reproducibility of coronary calcium quantification in repeat examinations with retrospectively ECG-gated multislice spiral CT. *Eur Radiol* 2002;**12**:1532–40.
- O'Rourke RA, Brundage BH, Froelicher VF, *et al*. American College of Cardiology/American Heart Association expert consensus document on electron-beam computed tomography for the diagnosis and prognosis of coronary artery disease. *Circulation* 2000;**102**:126–40.
- Wang S, Detrano RC, Secci A, *et al*. Detection of coronary calcification with electron-beam computed tomography: evaluation of interexamination reproducibility and comparison of three image-acquisition protocols. *Am Heart J* 1996;**132**:550–8.
- Bielak LF, Sheedy PF 2nd, Peyser PA. Coronary artery calcification measured at electron-beam CT: agreement in dual scan runs and change over time. *Radiology* 2001;**218**:224–9.
- Houslay ES, Cowell SJ, Prescott RJ, *et al*. Progressive coronary calcification despite intensive lipid-lowering treatment: a randomised controlled trial. *Heart* 2006;**92**:1207–12.
- Bostrom K, Demer LL. Regulatory mechanisms in vascular calcification. *Crit Rev Eukaryot Gene Expr* 2000;**10**:151–8.
- Bostrom KI. Cell differentiation in vascular calcification. *Z Kardiol* 2000;**89**(suppl 2):69–74.
- Canfield AE, Doherty MJ, Wood AC, *et al*. Role of pericytes in vascular calcification: a review. *Z Kardiol* 2000;**89**(suppl 2):20–7.
- Sangiorgi G, Rumberger JA, Severson A, *et al*. Arterial calcification and not lumen stenosis is highly correlated with atherosclerotic plaque burden in humans: a histologic study of 723 coronary artery segments using nondecalcifying methodology. *J Am Coll Cardiol* 1998;**31**:126–33.
- Rumberger JA, Simons DB, Fitzpatrick LA, *et al*. Coronary artery calcium area by electron-beam computed tomography and coronary atherosclerotic plaque area. A histopathologic correlative study. *Circulation* 1995;**92**:2157–62.
- Rumberger JA, Sheedy PF, Breen JF, *et al*. Electron beam computed tomographic coronary calcium score cutpoints and severity of associated angiographic lumen stenosis. *J Am Coll Cardiol* 1997;**29**:1542–8.
- Haberl R, Becker A, Leber A, *et al*. Correlation of coronary calcification and angiographically documented stenoses in patients with suspected coronary artery disease: results of 1,764 patients. *J Am Coll Cardiol* 2001;**37**:451–7.
- Stary HC, Chandler AB, Dinsmore RE, *et al*. A definition of advanced types of atherosclerotic lesions and a histological classification of atherosclerosis. A report from the committee on vascular lesions of the council on arteriosclerosis, American Heart Association. *Circulation* 1995;**92**:1355–74.
- Becker CR, Knez A, Ohnesorge B, *et al*. Visualization and quantification of coronary calcifications with electron beam and spiral computed tomography. *Eur Radiol* 2000;**10**:629–35.
- Waters D, Higginson L, Gladstone P, *et al*. Effects of monotherapy with an HMG-CoA reductase inhibitor on the progression of coronary atherosclerosis as assessed by serial quantitative arteriography. The Canadian coronary atherosclerosis intervention trial. *Circulation* 1994;**89**:959–68.
- Jukema JW, Bruschke AV, van Boven AJ, *et al*. Effects of lipid lowering by pravastatin on progression and regression of coronary artery disease in symptomatic men with normal to moderately elevated serum cholesterol levels. The regression growth evaluation statin study (REGRESS). *Circulation* 1995;**91**:2528–40.
- Achenbach S, Ropers D, Pohle K, *et al*. Influence of lipid-lowering therapy on the progression of coronary artery calcification: a prospective evaluation. *Circulation* 2002;**106**:1077–82.
- Callister TQ, Raggi P, Cooll B, *et al*. Effect of HMG-CoA reductase inhibitors on coronary artery disease as assessed by electron-beam computed tomography. *N Engl J Med* 1998;**339**:1972–8.
- Budoff MJ, Lane KL, Bakhsheshi H, *et al*. Rates of progression of coronary calcium by electron beam tomography. *Am J Cardiol* 2000;**86**:8–11.
- Hong C, Becker CR, Schoepf UJ, *et al*. Coronary artery calcium: absolute quantification in nonenhanced and contrast-enhanced multi-detector row CT studies. *Radiology* 2002;**223**:474–80.
- Schroeder S, Kopp AF, Baumbach A, *et al*. Noninvasive detection and evaluation of atherosclerotic coronary plaques with multislice computed tomography. *J Am Coll Cardiol* 2001;**37**:1430–5.
- Leber AW, Knez A, Becker A, *et al*. Accuracy of multidetector spiral computed tomography in identifying and differentiating the composition of coronary atherosclerotic plaques: a comparative study with intracoronary ultrasound. *J Am Coll Cardiol* 2004;**43**:1241–7.
- Leber AW, Knez A, White CW, *et al*. Composition of coronary atherosclerotic plaques in patients with acute myocardial infarction and stable angina pectoris determined by contrast-enhanced multislice computed tomography. *Am J Cardiol* 2003;**91**:714–8.
- Schroeder S, Kopp AF, Ohnesorge B, *et al*. Accuracy and reliability of quantitative measurements in coronary arteries by multi-slice computed tomography: experimental and initial clinical results. *Clin Radiol* 2001;**56**:466–74.
- Schroeder S, Kopp AF, Baumbach A, *et al*. Non-invasive characterisation of coronary lesion morphology by multi-slice computed tomography: a promising new technology for risk stratification of patients with coronary artery disease. *Heart* 2001;**85**:576–8.
- Mollet NR, Cademartiri F, Nieman K, *et al*. Noninvasive assessment of coronary plaque burden using multislice computed tomography. *Am J Cardiol* 2005;**95**:1165–9.
- Jakobs TF, Becker CR, Ohnesorge B, *et al*. Multislice helical CT of the heart with retrospective ECG gating: reduction of radiation exposure by ECG-controlled tube current modulation. *Eur Radiol* 2002;**12**:1081–6.
- Morin RL, Gerber TC, McCollough CH. Radiation dose in computed tomography of the heart. *Circulation* 2003;**107**:917–22.