REFERENCES

- 1 Aurigemma GP, Gottdiener JS, Shemanski L, et al. Predictive value of systolic and diastolic function for incident congestive heart failure in the elderly: the cardiovascular health study. *J Am Coll Cardiol* 2001;**37**:1042–8.
- 2 Redfield MM, Jacobsen SJ, Burnett JC, et al. Burden of systolic and diastolic ventricular dysfunction in the community. *JAMA* 2003;**289**:194–202.
- 3 Hunt SA. ACC/AHA 2005 guideline update for the diagnosis and management of chronic heart failure in the adult: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines. *J Am Coll Cardiol* 2005;**46**:1–82.
- 4 Burkhoff D, Maurer MS, Packer M. Heart failure with a normal ejection fraction: is it really a disorder of diastolic function? *Circulation* 2003;**107**:656–8.
- 5 Maurer MS, Spevack D, Burkhoff D, et al. Diastolic dysfunction: can it be diagnosed by Doppler echocardiography? *J Am Coll Cardiol* 2004;**44**:1543–9.
- 6 Ommen SR, Nishimura RA. A clinical approach to the assessment of left ventricular diastolic function by Doppler echocardiography. *Heart* 2003;**89**(suppl III):iii18–21.
- 7 Yu C-M, Lin H, Yang H, et al. Progression of systolic abnormalities in patients with "isolated" diastolic heart failure and diastolic dysfunction. *Circulation* 2002;**105**:1195–201.
- 8 Sanderson JE. Heart failure with a normal ejection fraction. *Heart*, (in press).
- 9 Hundley WG, Kitzman DW, Morgan DM, et al. Cardiac cycle-dependent changes in aortic area and distensibility are reduced in older patients with isolated diastolic heart failure and correlate with exercise intolerance. *J Am Coll Cardiol* 2001;**38**:796–802.
- 10 Kawaguchi M, Hay I, Fetics B, et al. Combined ventricular systolic and arterial stiffening in patients with heart failure and preserved ejection fraction: implications for systolic and diastolic reserve limitations. *Circulation* 2003;**107**:714–20.
- 11 Brutsaert DL, Sys SU. Diastolic dysfunction in heart failure. *J Card Fail* 1997;**3**:225–42.
- 12 Gillebert TC, Leite-Moreira AF, De Hert SG. Load dependent diastolic dysfunction in heart failure. *Heart Failure Rev* 2000;**5**:345–55.
- 13 Munagala VK, Hart CYT, Burnett JC, et al. Ventricular structure and function in aged dogs with renal hypertension. *Circulation* 2005;**111**:1128–35.
- 14 Weber T, Auer J, O'Rourke MF, et al. Diastolic dysfunction is accompanied with increased arterial wave reflections and prolonged mechanical systole [abstract]. *J Am Coll Cardiol* 2005;**45**(Suppl):132.
- 15 Anon. European Study Group on diastolic heart failure. How to diagnose diastolic heart failure. *Eur Heart J* 1998;**19**:990–1003.
- 16 Vasan RS, Levy D. Defining diastolic heart failure: a call for standardized diagnostic criteria. *Circulation* 2000;**101**:2118–21.
- 17 Tschöpe C, Kasner M, Westermann D, et al. The role of NT-proBNP in the diagnostics of isolated diastolic dysfunction: correlation with echocardiographic and invasive measurements. *Eur Heart J* 2005;**26**:2277–84.
- 18 Weber T, Auer J, O'Rourke MF, et al. Aortic stiffness, wave reflections, and the risk of coronary artery disease. *Circulation* 2004;**109**:184–9.
- 19 Weissler AM, Harris WS, Schoenfeld CD. Systolic time intervals in heart failure in man. *Circulation* 1968;**37**:149–59.
- 20 Pauca AL, O'Rourke MF, Kon ND. Prospective evaluation of a method for estimating ascending aortic pressure from the radial artery pressure waveform. *Hypertension* 2001;**38**:932–7.
- 21 Schiller NB, Shah PM, Crawford M, et al. Recommendations for quantification of the left ventricle by two-dimensional echocardiography. *J Am Soc Echocardiogr* 1989;**2**:358–67.
- 22 Devereux RB, Alonso DR, Lutas EM, et al. Echocardiographic assessment of left ventricular hypertrophy: comparison to necropsy findings. *Am J Cardiol* 1986;**57**:450–8.
- 23 Ommen SR, Nishimura RA, Appleton CP, et al. Clinical utility of Doppler echocardiography and tissue Doppler imaging in the estimation of left ventricular filling pressures: a comparative simultaneous Doppler-catheterization study. *Circulation* 2000;**102**:1788–94.
- 24 Nichols WW, O'Rourke MF. *McDonald's blood flow in arteries: theoretical, experimental and clinical principles*, 5th ed. London: Hodder Arnold, 2005.
- 25 Gillebert TC, Lew WYW. Influence of systolic pressure profile on rate of left ventricular pressure fall. *Am J Physiol Heart Circ Physiol* 1991;**261**:H805–13.
- 26 Yano M, Kohno M, Kobayashi S, et al. Influence of timing and magnitude of arterial wave reflection on left ventricular relaxation. *Am J Physiol Heart Circ Physiol* 2001;**280**:H1846–52.
- 27 Iketani T, Takazawa K, Ibukiya C. The influence of changes in loading patterns on left ventricular relaxation in humans. *Jpn Circ J* 1998;**62**:581–5.
- 28 Ihara T, Komamura K, Shen YT, et al. Left ventricular systolic dysfunction precedes diastolic dysfunction during myocardial ischemia in conscious dogs. *Am J Physiol* 1994;**267**:H333–43.
- 29 Warner JG Jr, Metzger C, Kitzman DW, et al. Losartan improves exercise tolerance in patients with diastolic dysfunction and a hypertensive response to exercise. *J Am Coll Cardiol* 1999;**33**:1557–62.
- 30 Paulus WJ, Vantrimpont PJ, Shah AM. Acute effects of nitric oxide on left ventricular relaxation and diastolic distensibility in humans. *Circulation* 1994;**89**:2070–8.
- 31 Chen CH, Nakayama M, Talbot M, et al. Verapamil acutely reduces ventricular-vascular stiffening and improves aerobic exercise performance in elderly individuals. *J Am Coll Cardiol* 1999;**33**:1602–9.
- 32 Hay I, Rich J, Ferber P, et al. Role of impaired myocardial relaxation in the production of elevated left ventricular filling pressure. *Am J Physiol Heart Circ Physiol* 2005;**288**:H1203–8.

IMAGES IN CARDIOLOGY

doi: 10.1136/hrt.2006.090829

Angiographic presentation of sinus of Valsalva aneurysm

Angiography of the aorta in a right anterior oblique view.

A 75-year-old patient was admitted because of dyspnoea, and a coronary angiography was performed due to slightly elevated troponin concentration. The coronary arteries were unremarkable. The angiography of the aorta revealed an aneurysm of the right sinus of Valsalva with massive contrast enhancement (see panel and video 1; to view video footage visit the *Heart* website—<http://www.heartjnl.com/supplemental>). Transoesophageal echocardiography showed spontaneous echo contrast in the area of the aneurysm (video 2). The troponin elevation was thus possibly due to microembolisation from the aneurysm. The aneurysm was successfully resected, and the postoperative course was uneventful.

M Handke
M Göpfrich
H Keller

handke@mm31.ukl.uni-freiburg.de

There are no competing interests.

To view video footage visit the
Heart website—<http://www.heartjnl.com/supplemental>