Overexpression of Thrombospondin-1 Decreases Angiogenesis and Inhibits the Growth of Human Cutaneous Squamous Cell Carcinomas

Michael Streit,* Paula Velasco,*
Lawrence F. Brown,[†] Mihaela Skobe,*
Lisa Richard,* Lucia Riccardi,* Jack Lawler,[†] and
Michael Detmar*

From the Cutaneous Biology Research Center, Department of Dermatology,* Massachusetts General Hospital and Harvard Medical School, Charlestown; and the Department of Pathology,[†] Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, Massachusetts

The function of the endogenous angiogenesis inhibitor thrombospondin-1 (TSP-1) in epithelial tumor development has remained controversial. We studied the *in vitro* growth characteristics and the *in vivo* tumor xenograft growth of the human squamous cell carcinoma cell lines A431 and SCC-13, stably transfected to overexpress human TSP-1. Overexpression of TSP-1 inhibited tumor growth of A431 xenotransplants, and completely abolished tumor formation by SCC-13 cells. TSP-1 overexpressing A431 tumors were characterized by extensive areas of necrosis and by decreased tumor vessel number and size. The effects of TSP-1 on tumor cell growth were indirect since tumor cell proliferation rates *in vivo* and *in vitro*, anchorage-dependent and -independent growth *in vitro*, and susceptibility to induction of apoptosis by serum withdrawal were unchanged in TSP-1 overexpressing tumor cells. However, TSP-1 overexpression up-regulated the TSP-1 receptor CD36, leading to enhanced adhesion of A431 cells to TSP-1. These findings establish TSP-1 as a potent inhibitor of angiogenesis and tumor growth in carcinomas of the skin. (Am J Pathol 1999, 155:441–452)

To grow beyond minimal size and to metastasize, tumors need to induce the growth of new blood vessels (angiogenesis) providing a lifeline for tumor sustenance and waste disposal.¹ Tumor development is associated with increased release of angiogenesis factors, most prominently of vascular endothelial growth factor (VEGF).² Several studies have shown that overexpression of angiogenesis factors in experimental tumors leads to enhanced tumor growth and vascularization, and therapeutic inhibition of VEGF activity has been shown to inhibit tumor growth and metastasis.^{3–5}

In contrast, much less is known about the expression and biological role of endogenous inhibitors of angiogenesis during carcinogenesis. Several naturally occurring angiogenesis inhibitors have been identified, including thrombospondin-1 (TSP-1),⁶ TSP-2,⁷ angiostatin,⁸ and endostatin.⁹ TSP-1 is a 420-kd homotrimeric matricellular glycoprotein that regulates attachment, proliferation, migration, and differentiation of various cell types (for review, see Ref. 10). TSP-1 inhibits proliferation and migration of vascular endothelial cells *in vitro* and inhibits neovascularization *in vivo*, contributing to the normal quiescence of the vasculature.¹¹ However, controversial data have been reported regarding the role of TSP-1 in epithelial tumor growth and metastasis. TSP-1 protein expression was shown to be inversely correlated to cellular differentiation in several squamous cell carcinoma (SCC) cell lines,¹² and was shown to induce SCC proliferation, adhesion, migration, and invasion of cells *in vitro*.^{13–15} Enhancement of *in vitro* tumor cell invasion by TSP-1 has also been reported for breast, lung, and pancreatic carcinoma cell lines.^{16–19} Based on the observation that antisense inhibition of TSP-1 in SCC resulted in suppression of tumor growth *in vivo*,²⁰ it was suggested that TSP-1 may promote tumor growth.²¹ In contrast, other studies reported that TSP-1 expression was inversely correlated with malignant progression in human lung, breast, and bladder carcinoma cell lines.^{22,23}

In human skin, TSP-1 is deposited in the basement membrane,²⁴ contributing to the antiangiogenic barrier that separates the avascular epidermis from the vascularized dermis. Recently, we found that TSP-1 expression was down-regulated in squamous cell carcinomas (SCC) of the skin, richly vascularized malignant tumors derived from epidermal keratinocytes (Detmar M, Velasco P, Tognazzi K, Brown LF, unpublished data). Based on these findings, we studied the biological role of TSP-1 for cutaneous carcinoma growth, using stable tumor cell trans-

Supported by NIH/NCI grant CA69184 (to M.D.), by Deutsche Forschungsgemeinschaft grant De483/3–2 (to M.D.), by Deutscher Akademischer Austauschdienst (to M.St.), by the Human Frontier Science Program (M.Sk.), by NIH/HLBI grant HL28749 (to J.L.), and by the Cutaneous Biology Research Center through the Massachusetts General Hospital/Shiseido Co. Ltd. Agreement (M.D.).

Accepted for publication April 11, 1999.

Address reprint requests to Michael Detmar, CBRC/Dept. of Dermatology, Massachusetts General Hospital, Building 149, 13th Street, Charlestown, MA 02129. E-mail: mdetmar@cbrc.mgh.harvard.edu.

fectants in an intradermal xenograft model. Here, we report that TSP-1 overexpression reduced intradermal tumor growth of A431 epidermoid carcinoma cells and completely inhibited intradermal tumor formation of SCC-13 squamous cell carcinomas, although no direct effect on tumor cell proliferation was detected *in vitro* and *in vivo*. TSP-1 overexpressing A431 tumors were characterized by extensive areas of necrosis and by decreased tumor vessel numbers and sizes. TSP-1 overexpression also up-regulated the TSP-1 receptor CD36, leading to enhanced A431 cell adhesion to TSP-1. These findings establish TSP-1 as a potent inhibitor of angiogenesis and growth of cutaneous squamous cell carcinomas.

Materials and Methods

Cell Culture

The human epidermoid carcinoma cell line A431 was obtained from the American Type Culture Collection (Rockville, MD) and was maintained in Dulbecco's minimal essential medium (DMEM; Gibco BRL, Grand Island, NY) supplemented with 10% fetal bovine serum (FBS) and 1% L-glutamine (all purchased from Gibco BRL).

The human cutaneous squamous carcinoma cell line SCC-13,²⁵ kindly provided by Dr. James Rheinwald (Harvard Medical School, Boston), was maintained in DMEM/Ham's F-12 medium (3:1) supplemented with 5% FBS, 1% L-glutamine, 50 μ g/ml insulin, 0.1 U/ml epidermal growth factor, 18.2 mg/ml adenine, and 0.4 μ g/ml hydrocortisone (all purchased from Gibco BRL). Human dermal microvascular endothelial cells (HDMEC) were isolated from neonatal foreskins and cultivated as recently described.²⁶

Cell Transfection and Selection

A 3.643-kb human TSP-1 cDNA sequence, comprising the full TSP-1 coding sequence (nucleotides 67–3689 of the human TSP-1 sequence; GenBank accession code X04665), was kindly provided by Dr. Luisa Iruela-Arispe, University of California, Los Angeles. The TSP-1 cDNA was cloned into a pGEM7Z vector (Promega, Madison, WI). The sequence was verified by restriction mapping and by direct sequencing using the Sanger dideoxy method. After restriction digestion with *SacI*, *SphI*, and *PvuI* (Gibco BRL), a 3.6-kb fragment was gel purified, blunted, and ligated into a pcDNA3.1Zeo(-) expression vector (Invitrogen, San Diego, CA) which contains a CMV-enhancer-promoter and a Zeocin selection cassette. Subconfluent cultures were stably transfected either with pcDNA3.1Zeo(-) vector containing the full-length human TSP-1 cDNA or with pcDNA3.1Zeo(-) vector alone using the SuperFect transfection reagent (Qiagen, Chatsworth, CA) according to the manufacturer's protocol. Forty-eight hours after transfection, cells were split 1:3 into their full growth medium containing 250 μ g/ml Zeocin (Invitrogen) to select transfectants. Stably transfected clones were expanded, and 10 clones were characterized for TSP-1 mRNA and protein expression.

RNA Isolation and Northern Blot Analysis

Total cellular RNA was isolated from stable transfectants and from intradermal tumors using the RNeasy kit (Qiagen), according to the manufacturer's instructions. The isolated RNA was subjected to electrophoresis and transferred to Biotrans nylon supported membranes (ICN Pharmaceuticals, Costa Mesa, CA). ³²P-radiolabeled cDNA probes were prepared with a random primed synthesis kit (Multiprime; Amersham, Arlington Heights, IL). We used a 4.1-kb TSP-1 cDNA probe and a 300-bp human VEGF cDNA probe which recognizes all known VEGF variants. A 2.0-kb human β -actin cDNA probe purchased from Clontech (Palo Alto, CA) was used as a control for equal RNA loading. Blots were washed at high stringency as described²⁷ and exposed to X-OMAT MR film (Kodak, Rochester, NY) or a PhosphorImager screen (Molecular Dynamics, Sunnyvale, CA). mRNA expression was quantitated with a Molecular Dynamics scanning densitometer using the ImageQuant software.

Western Blot Analysis

Western Blot analyses were performed on cell lysates and conditioned media from stably transfected A431 cells, SCC-13, and HDMEC. Cells were grown to confluence in 100-mm dishes, washed with phosphate buffered saline (PBS), and lysed as described.²⁸ Cell lysates were homogenized using a cell shredder (Qiagen), and protein concentrations were determined using the Bio-Rad protein assay (Bio-Rad, Hercules, CA). Conditioned media were obtained from confluent cells grown for 48 hours in serum-free culture medium. TSP-1 was concentrated using heparin beads (Sigma, St. Louis, MO). All samples were boiled in denaturing sample buffer, and equal amounts according to the protein assay were electrophoresed on polyacrylamide gels under reducing conditions.²⁹ Proteins were blotted onto polyvinylidene difluoride membranes (Bio-Rad). To verify equal protein loading membranes were stained with 0.1% Ponceau red (Sigma) diluted in 5% acetic acid. Membranes were incubated overnight in PBS containing 0.1% Tween-20 and 3% bovine serum albumin to block nonspecific binding. Membranes were then incubated with primary antibodies directed against human TSP-1 (clone 133; Genzyme, Cambridge, MA), human CD36 (clone 1A7; NeoMarkers, Fremont, CA), or human VEGF (clone 2352, kindly provided by Dr. Don Senger, Beth Israel Deaconess Medical Center, Boston, MA), washed in PBS/Tween, incubated with horseradish peroxidase-conjugated anti-mouse IgG (Amersham), and analyzed by the enhanced-chemiluminescence system (Amersham). Protein expression was quantitated with a Molecular Dynamics scanning densitometer using the ImageQuant software.

Cell Growth and Apoptosis Assays

To determine whether TSP-1 overexpression influences tumor cell proliferation, we measured anchorage-dependent and -independent cell growth rates. 1×10^5 A431

cells were plated in duplicate into 100-mm culture dishes and total cell numbers per dish were determined after 2, 4, and 6 days using a hemocytometer. Anchorage-independent growth rates were determined by using a soft-agar assay as described.³⁰ Ten thousand control transfected or TSP-1 transfected A431 cells were transferred into six 30-mm cell culture dishes with a 2-mm grid (Nunc, Naperville, IL). The dishes were incubated at 37°C and 5% CO₂, and colonies were counted after 8 days. The effects of conditioned media from TSP-1 transfectants or from vector transfected controls on endothelial cell proliferation were determined using the BrdU labeling and detection kit (Boehringer, Mannheim, Germany). Human dermal microvascular endothelial cells were grown in 24-well plates in the presence of cell culture medium or conditioned media, supplemented with 10% FBS for 24 hours. HDMEC were also co-incubated with either TSP-1 (25 µg/ml) and/or a TSP-1 neutralizing antibody (50 µg/ml, clone AB-1, Neomarkers) or an isotype-specific control antibody (clone MOPC 21, Sigma). Cell proliferation was assayed according to the manufacturer's instructions. The absorbance was determined at 405 nm using a microtiter plate reader (Titertek, Huntsville, AL). The results represent the mean values ± SD of four dishes per group. Apoptosis induced by serum withdrawal was studied in subconfluent A431 cell clones after 6 days in serum-free medium. The percentage of apoptotic cells was determined as described,³¹ using the Fluorescein-FragEL DNA fragmentation kit (Oncogene, Cambridge, MA) according to the manufacturer's instructions, and a Beckton-Dickinson FACS-Scan (Franklin Lanes, NJ).

Cell Adhesion Assay

Twenty-four-well plates were coated with 20 µg/ml human collagen type I (Sigma) or with 20 µg/ml platelet-derived human TSP-1 (Sigma) for 1 hour at 4°C, followed by 100 mg/ml bovine serum albumin (Sigma) to block remaining protein binding sites. A431 cells (1 × 10⁶ cells/ml) in serum-free DMEM medium (Gibco BRL) were mixed with a blocking mouse monoclonal antibody to human CD36 (Neomarkers) or with mouse control IgG (Sigma) at 5 µg/ml. One hundred µl of the cell suspensions were added to each well and were incubated at 37°C for 60 minutes. Unattached cells were removed by three gentle washes with PBS. Attached cells were fixed with 0.1% glutaraldehyde (Sigma), stained with a 0.1% crystal violet solution (Sigma), and washed three times with PBS. Crystal violet adsorbed onto cells was solubilized with 0.2% Triton X-100 (Sigma), and absorbance was measured at 590 nm in a Titertek microtiter plate reader. The absorbance of 1 × 10⁵ fixed cells served as control (100% value).³² The unpaired Student's *t*-test was used for statistical analysis of the results.

Growth of Xenografts in Nude Mice

Confluent A431 or SCC-13 cells, stably transfected with a human TSP-1 expression vector or with the expression

vector alone, were trypsinized and resuspended in serum-free DMEM medium (Gibco BRL) at a density of 2 × 10⁷ cells/ml. Two million tumor cells of each type were injected intradermally into both flanks of five 8-week-old female BALB/c (nu/nu) mice. Two control clones and 3 TSP-1 overexpressing clones were investigated. The smallest and largest tumor diameter were measured weekly, using a digital caliper, and tumor volumes were calculated using the following formula:

$$\text{Volume} = 4/3 \times \pi \times (1/2 \text{ smaller diameter})^2 \times 1/2 \text{ larger diameter}$$

Mice were sacrificed after 4 weeks in the group of A431 injected animals and after 7 weeks in the group of SCC-13 injected animals or earlier if the largest tumor diameter reached 20 mm. All animal studies were approved by the subcommittee on Research Animal Care of the Massachusetts General Hospital.

In Situ Hybridization and Immunohistochemistry

In situ hybridization was performed on 6-µm paraffin sections of tumor xenografts as described.²⁷ The sense and antisense single-stranded RNA probes for human VEGF were transcribed from a pGEM-3Zf(+) vector containing a 204-bp polymerase chain reaction fragment common to all known VEGF splicing variants. A RNA probe to human TSP-1 was transcribed from a pBluescript II KS+ vector containing a 240-bp polymerase chain reaction fragment of the coding region of human TSP-1. Immunohistochemical staining was performed on 6-µm frozen or paraffin sections of tumor xenotransplants as previously described,³³ using monoclonal antibodies against mouse CD31 (Pharmingen, San Diego, CA), human TSP-1 (Genzyme), human CD36 (Neomarkers), and human PCNA antigen (Zymed Laboratories, San Francisco, CA).

Computer-Assisted Morphometric Analysis of Tumor Vessels

To determine the degree of tumor-induced angiogenesis, cryostat sections of tumor xenografts were stained with an anti-mouse CD31 monoclonal antibody. Representative sections obtained from five tumors from each cell clone were analyzed, using a Nikon E-600 microscope (Nikon, Melville, NY). Images were captured with a Spot digital camera (Diagnostic Instruments, Sterling Heights, MI), and morphometric analyses were performed using the IP LAB software program (Scanalytics Inc.; Fairfax, VA). Three different fields at 60x magnification were examined on each section, and the number of vessels per mm², the size distribution, the average, and the total area of tumor blood vessels were determined. The unpaired *t*-test was used to analyze differences in the vessel areas.

Results

Overexpression of TSP-1 in Two Human Cutaneous Tumor Cell Lines

The human squamous cell carcinoma cell lines A431 and SCC-13 were chosen with respect to their endogenous TSP-1 and VEGF secretion and their *in vivo* growth characteristics. A431 cells are characterized by strong secretion of VEGF but little or no TSP-1 secretion (Figure 1), and form fast growing and highly vascularized tumors *in vivo*.³⁴ SCC-13 cells express low levels of VEGF but synthesize substantial amounts of TSP-1 (Figure 1), and grow slowly as well differentiated, rounded tumors. The TSP-1 expression levels of multiple TSP-1 and control transfected clones were determined by Northern Blot analyses (Figure 1A). The highest levels of TSP-1 mRNA were detected in A431 clones 10, 12, and 19 and SCC-13 clones 1, 2, and 3. Western blot analyses confirmed that increased TSP-1 mRNA levels correlated with increased amounts of TSP-1 protein. In TSP-1 transfected A431 cell clones, strong expression of the 180 kd TSP-1 protein was found in cell lysates and in culture supernatants, confirming efficient secretion of TSP-1 (Figure 1B). In contrast, little or no TSP-1 protein was detected in A431 cells transfected with vector only. In SCC-13 cell clones, TSP-1 was already expressed by control transfectants (Figure 1B); three TSP-1 transfected clones showed markedly increased TSP-1 secretion although the amount of cell-associated TSP-1 was not significantly increased. TSP-1 expression was also detected in cell lysates but not in conditioned medium harvested from human dermal microvascular endothelial cells (Figure 1B). The biological activity of transfected human TSP-1 was confirmed in HDMEC proliferation assays. HDMEC proliferation was significantly inhibited after a 24 hour incubation with conditioned media harvested from TSP-1 overexpressing A431 and SCC-13 cell clones, as compared to HDMEC incubated with unconditioned media (Figure 1, C and E). Supernatants obtained from control transfected A431 clones stimulated HDMEC proliferation. Incubation with a TSP-1 neutralizing antibody but not with an isotype-control antibody abolished the inhibition of HDMEC proliferation by culture medium supplemented with TSP-1 and by conditioned medium obtained from TSP-1 overexpressing cell clones (1D).

Growth of TSP-1 Overexpressing Tumor Cells in Vitro and Tumor Growth in Vivo

Our results showed no differences in anchorage-dependent cell proliferation between TSP-1 overexpressing A431 clones and control clones growing in plastic culture dishes (Figure 2A). Anchorage-independent cell growth was studied by determination of colony numbers in a soft agar assay. No significant differences in the number of colonies were observed between the different cell clones (Figure 2B). To compare the susceptibility to induction of apoptosis by serum withdrawal, A431 clones were cultured under serum-free conditions for 6 days. No signif-

icant differences in the percentage of apoptotic cells were found between TSP-1 overexpressing A431 clones ($13.67\% \pm 4.04\%$) and control A431 clones ($12.33\% \pm 1.45\%$). To determine the biological effects of TSP-1 overexpression on the orthotopic tumor growth of A431 and SCC-13 cells *in vivo*, tumor cells were injected intradermally into the flanks of immunodeficient nude mice. Control transfected A431 cell clones formed rapidly growing tumors (Figure 2E), reaching a volume of 2000–3000 mm³ after 4 weeks. Stable overexpression of TSP-1 resulted in a significant inhibition of tumor growth by 50 to 75% after 4 weeks, as compared to control tumors (Figure 2E). Northern Blot analysis of RNA extracted from two representative tumors of each A431 cell clone confirmed that TSP-1 transfected tumor cell clones maintained TSP-1 mRNA expression *in vivo* (data not shown), with the highest TSP-1 mRNA expression and the slowest tumor growth in A431 clone T10. Northern analysis of the same RNA samples demonstrated equal levels of VEGF mRNA expression in TSP-1 overexpressing and control tumors. The effects of transfected TSP-1 on *in vivo* tumor growth were confirmed in transfected SCC-13 cell clones. After 2 weeks, control transfected SCC-13 clones formed slowly growing intradermal tumors (Figure 2F). In contrast, TSP-1 overexpression in SCC-13 cells led to a complete inhibition of *in vivo* tumor growth in all clones tested (clones T2 and T3) over an observation period of up to 12 weeks (Figure 2F).

Histological Characteristics of TSP-1 Overexpressing Tumors

Extensive areas of necrosis were detected in TSP-1 overexpressing tumors (see Figure 4B), whereas only occasional small necrotic foci were found in control tumors (see Figure 4A). However, the fraction of proliferating cells within the viable tumor areas, as determined by staining for the PCNA antigen, was unchanged (data not shown). The expression and distribution of TSP-1 within A431 tumors was assessed by *in situ* hybridization and by immunohistochemistry. Only weak TSP-1 mRNA expression was detected in control tumor cells (Figure 3, A and B), and TSP-1 protein expression was predominantly found in the dermal-epidermal basement membrane zone of adjacent normal skin and in blood vessels, but not in tumor cells (see Figure 5A). In contrast, strong TSP-1 mRNA expression was detected in TSP-1 overexpressing tumor cells (Figure 3, C and D), and immunohistochemistry demonstrated massive TSP-1 deposits in the tumor stroma (see Figure 5B). In accordance with the Northern Blot results, no major differences of VEGF mRNA expression were found between TSP-1 overexpressing and control tumors by *in situ* hybridization (Figure 3, E–H). Up-regulation of VEGF mRNA expression was observed adjacent to areas of necrosis in tissue samples derived from TSP-1 overexpressing tumors and control tumors.

Figure 1. A: Overexpression of human TSP-1 mRNA in stably transfected A431 and SCC-13 cell clones as compared to vector transfected control clones was confirmed by Northern Blot analysis. mRNA expression of human VEGF remained unchanged in TSP-1 overexpressing and control clones. Hybridization with a human β -actin probe served as control for equal loading. B: Western blot analysis demonstrated increased amounts of TSP-1 protein in cell lysates and conditioned media obtained from TSP-1 transfected A431 cells. TSP-1 migrates at approximately 180 kd. Control transfected SCC-13 cell clones synthesized substantial amounts of TSP-1. Increased TSP-1 secretion was detected in all three TSP-1 transfected cell clones (C1–3). The secretion of VEGF is not modified by TSP-1 overexpression in A431 and SCC-13 cell clones. HDMEC synthesize but do not secrete detectable amounts of TSP-1. C: Mitogenic effect of conditioned media obtained from control-transfected A431 clones (C1–3) as compared to unconditioned medium (M). Conditioned media obtained from TSP-1 transfected A431 clones (T10, T12, T19) significantly inhibited HDMEC proliferation, as determined by BrdU incorporation ($P < 0.01$). Bars represent mean values $\pm$ SD of two independent experiments. D: Co-incubation with a TSP-1 neutralizing antibody (anti-TSP-1) but not with an isotype-specific control antibody (IgG) abolished the inhibitory effect of TSP-1 (25 μ g/ml) added to unconditioned medium (open bars) or of conditioned medium harvested from a TSP-1 overexpressing A431 cell clone (T19) (closed bars) on HDMEC proliferation. E: Conditioned media obtained from TSP-1 transfected SCC-13 cell clones (T1–3) but not from control-transfected clones (C1–3) inhibited HDMEC proliferation.

Figure 2. A: No influence of transfected TSP-1 on the anchorage-dependent growth of A431 cell clones (T10, T12, T19), as compared to control transfectants (C1–3). B: No major differences in anchorage-independent growth of TSP-1 transfected and control-transfected A431 clones, as determined in a soft agar assay. Mean values $\pm$ SD of two independent experiments. C, D: Rarefaction of large tumor blood vessels supplying TSP-1 overexpressing A431 xenografts (D), as compared to control-transfected A431 tumors (C). Scale bar = 1 cm. E: Decreased tumor growth of TSP-1 overexpressing A431 cells (T10, T12, T19), as compared to control-transfected clones (C2, C3). Values represent mean values $\pm$ SEM for 10 tumors for each clone and time point. F: Complete inhibition of tumor growth of TSP-1 overexpressing SCC-13 clones T2 and T3 (values identical to T2), as compared to control clones C2 and C3. Note different scale of the y axis, as compared to Figure 2E.

Figure 3. *In situ* hybridization demonstrates strong TSP-1 mRNA expression in 3-week-old TSP-1 transfected A431 xenografts (C, D), whereas little or no TSP-1 mRNA expression was detected in control tumors (A, B). In contrast, VEGF mRNA expression remained unchanged in TSP-1 transfected A431 xenotransplants (G, H), as compared to control tumors (E, F). Scale bar = 200 μm .

Morphometric Analysis of Tumor Vessels in TSP-1 Overexpressing Tumor Xenografts

The morphology of tumor supplying blood vessels was studied on the surface of tumors of identical size derived from TSP-1 transfected A431 clones after 4 weeks and control transfectants after 3 weeks (Figure 2, C and D). A significant rarefaction of large blood vessels was seen selectively on the surface of TSP-1 overexpressing tumor xenografts. To determine the microvascular density within TSP-1 transfected and control A431 tumors, frozen sections of 5 different tumors derived from the same A431 clone were stained with an antibody against mouse CD31 (Figure 4, C and D). Morphometric analysis revealed decreased microvessel densities (Figure 4E) in 3-week-old tumors derived from TSP-1 overexpressing clones T10 (28 ± 10 vessels/ mm^2), T12 (46 ± 12 vessels/ mm^2), and T19 (32 ± 7 vessels/ mm^2), as compared to control clone C2 (60 ± 19 vessels/ mm^2) and C3 (52 ± 20 vessels/ mm^2). Moreover, the average area per vessel was significantly ($P < 0.001$) smaller in TSP-1 overexpressing tumors (Figure 4G) derived from clone T10 (729 ± 213 μm^2), T12 (551 ± 157 μm^2), and T19 (755 ± 189 μm^2), as compared to control tumors derived from clone C2 (1057 ± 307 μm^2) and C3 (1298 ± 504 μm^2). The fraction of large blood vessels with an area of more than 2000 μm^2 was decreased to fewer than 1% of all vessels in TSP-1 overexpressing tumors, as compared to more than 10% in control tumors (Figure 4H). In accordance with these data, the total area covered by blood vessels (Figure 4F) was significantly ($P < 0.001$) smaller in tumors derived from TSP-1 transfected clones T10 ($1.7 \pm 0.6\%$), T12 ($2.6 \pm 1.2\%$) and T19 ($2.2 \pm 0.5\%$), as compared to control transfected clones C2 ($5.8 \pm 2\%$) and C3 ($5.4 \pm 1.7\%$).

Up-Regulation of the TSP-1 Receptor CD36 in TSP-1 Transfected Tumor Cells and Xenografts

It has been suggested that the antiangiogenic effect of TSP-1 is mediated through the CD36 receptor on endothelial cells.³⁵ We studied the expression of CD36 in TSP-1 overexpressing A431 tumors and controls by immunohistological staining. While CD36 expression was nearly absent in control transfected tumors (Figure 5C), strong CD36 expression was found in a substantial number of TSP-1 overexpressing A431 tumor cells (Figure 5D). These findings were confirmed by Western blot analysis, demonstrating strong up-regulation of CD36 in cell lysates obtained from TSP-1 overexpressing A431 (Figure 5E) and SCC-13 (Figure 5F) clones as compared to control clones. When subconfluent A431 control cells (clones 1, 2, and 3) were incubated for 24 hours with conditioned media taken from TSP-1 overexpressing A431 clone T19, a 1.5 to ninefold induction of CD36 expression was detected by Western blot, as compared to incubation with conditioned media from control clones (Figure 5G). In contrast, CD36 was already strongly expressed in untreated HDMEC, and incubation with TSP-1 containing conditioned media did not further increase CD36 protein expression levels (data not shown). To study the biological significance of increased CD36 expression on TSP-1 transfected A431 clones, we studied cell adhesion to immobilized TSP-1 and collagen type I. TSP-1 overexpressing A431 cells showed significantly enhanced cell adhesion onto a TSP-1 matrix ($P < 0.001$) but not onto a collagen type I matrix (Figure 5, H and I). Incubation with a blocking antibody against the CD36 receptor significantly reduced adhesion of TSP-1 overexpressing A431 cells to TSP-1 ($P < 0.001$), but not to collagen type I (Figure 5, H and I). In contrast, the adhe-

Figure 4. Extensive areas of tumor necrosis (*) in TSP-1 overexpressing A431 xenotransplants (B), as compared to control tumors (A); hematoxylin-eosin stain. Scale bar = 200 μ m. Immunohistochemical staining with an anti-mouse CD31 monoclonal antibody revealed inhibition of angiogenesis in TSP-1 overexpressing tumors (D), as compared to control tumors (C). **E:** Moderate reduction of microvascular densities in TSP-1 overexpressing tumors, as measured by the vessel number per mm² tumor area. **F:** Significant decrease of the area covered by blood vessels in TSP-1 overexpressing tumors compared to control tumors ($P < 0.001$). **G:** Significant reduction of the average blood vessel area ($P < 0.001$); **H:** loss of blood vessels of more than 2000 μ m² in TSP-1 overexpressing tumors, as compared to control tumors. CD31-stained blood vessels were evaluated in three different 10 $\times$ fields in sections of five different tumors derived from each A431 cell clone. Data are expressed as mean values $\pm$ SEM.

Figure 5. A–D: Immunohistochemical analysis demonstrates strong TSP-1 staining in TSP-1 transfected A431 xenotransplants, predominantly localized in the tumor stroma (B), as compared to the sparse labeling observed in control tumors (A). Little or no detectable CD36 expression in control A431 tumors (C), but strong CD36 expression in TSP-1 transfected xenotransplants (D). Scale bar = 200 μ m. E, F: Western blot analysis demonstrates increased CD36 expression in cell culture lysates obtained from TSP-1 transfected A431 clones (T10, T12, and T19) (E) and SCC-13 cell clones (T1–T3) (F), as compared to control clones. G: CD36 expression is induced in cultured A431 cells (C1, ninefold; C2, sixfold; C3, 1.5-fold) after 24 h incubation with conditioned medium obtained from TSP-1 overexpressing clones. H: Significantly increased adhesion of TSP-1 transfected A431 cells to immobilized TSP-1, as compared to control cells ($P < 0.001$). Incubation with an anti-CD36 blocking antibody significantly ($P < 0.001$) reduced adhesion of TSP-1 transfected A431 cells to levels observed in control cells. I: No significant difference in adhesion to collagen type I between control and TSP-1 transfected A431 cells. An anti-CD36 blocking antibody did not modify A431 cell adhesion to collagen type I. Scale bars represent mean values $\pm$ SD of two independent experiments.

sion of A431 control cells to TSP-1 or collagen type I was not affected. These results suggest that overexpression of TSP-1 in A431 and SCC-13 tumor cells leads to enhanced expression of its receptor CD36 with potential consequences for tumor cell adhesion.

Discussion

Conflicting results have been reported regarding the function of TSP-1 in tumor progression. We chose two established epithelial skin cancer models to enhance tumor cell TSP-1 expression by stable transfection. Our study was designed to characterize the effects of transfected TSP-1 on the orthotopic tumor growth of experimental skin cancer and to characterize the mechanisms by which TSP-1 exerts these effects.

Overexpression of TSP-1 in A431 cell xenotransplants potently decreased tumor growth, as compared to control tumors transfected with vector only. Increased TSP-1 secretion by stable transfectants was confirmed by Western blot analyses of conditioned media, and the bioactivity of transfected TSP-1 was confirmed in a proliferation assay using human dermal microvascular endothelial cells which by themselves did not secrete substantial amounts of TSP-1. Addition of conditioned media from TSP-1 transfected A431 clones significantly inhibited HD-MEC proliferation, whereas conditioned media from control A431 clones stimulated HDMEC growth. The potent inhibitory effect of conditioned media from TSP-1 transfected cell clones on HDMEC proliferation was TSP-1 specific since it was abolished by co-incubation with a TSP-1-neutralizing antibody. *In situ* hybridizations of tumor xenotransplants demonstrated that TSP-1 mRNA expression was maintained at high levels in TSP-1 transfected tumor cell clones. Together, these data provide evidence for a potent inhibitory effect of TSP-1 on skin cancer growth. In accordance with our results, TSP-1 has been reported to inhibit the *in vivo* growth of MDA-MB-435 breast carcinoma cells³⁶ and of v-src transformed NIH 3T3 fibroblasts.³⁷ Furthermore, TSP-1 negatively affected the tumor forming ability of transformed mouse endothelial cells,³⁸ of human glioblastoma cells,³⁹ and of human B16/F10 melanoma cells.⁴⁰ However, a previous investigation using the human squamous cell carcinoma cell line 11B reported that reduction of TSP-1 secretion by antisense transfection inhibited tumor growth *in vivo*, and the authors suggested that TSP-1 overexpression in squamous cell carcinomas might lead to enhanced tumor growth.²⁰ Our data provide strong evidence against this hypothesis and demonstrate, furthermore, that TSP-1 may even completely prevent the formation of cutaneous squamous cell carcinomas. We could demonstrate that stable TSP-1 transfection of human SCC-13 cells, a slowly growing human squamous cell carcinoma line with formation of highly differentiated tumors *in vivo*, induced a complete inhibition of any detectable tumor growth over an observation period of 12 weeks after xenotransplantation.

The tumor growth inhibition induced by TSP-1 in cutaneous squamous cell carcinomas was not due to direct

TSP-1-mediated inhibition of tumor cell growth. We used three independent *in vitro* assays to investigate TSP-1 effects on A431 cells: (1) Anchorage-dependent cell growth in monolayer culture; (2) anchorage-independent cell growth, as determined by the ability to form colonies in soft agar; and (3) susceptibility to induction of apoptosis by serum withdrawal. No significant differences between TSP-1 transfected A431 clones and control transfected A431 clones were detected in any of these studies. These findings are in accordance with previous findings, showing that TSP-1 transfection did not influence the *in vitro* growth of human MDA-MB-453 breast cancer cells³⁶ and B16/F10 melanoma cells.⁴⁰ However, they are in contrast to a previous investigation, reporting a direct correlation between the levels of TSP-1 secretion by several squamous cell carcinoma cell lines and enhanced *in vitro* cell growth and invasion.⁴¹ Moreover, another study reported that antisense-mediated down-regulation of TSP-1 expression in a squamous cell carcinoma cell line led to decreased *in vitro* cell proliferation, suggesting TSP-1 as a promoter of tumor growth.²⁰ In 3T3 fibroblasts, antibodies to TSP-1 were reported to inhibit cell proliferation, and TSP-1 overexpression induced serum- and anchorage-independent growth.⁴² While there is no obvious explanation for these discrepancies, varying susceptibilities of different cell types (eg, mesenchymal versus epithelial) or of distinct cell lines to TSP-1 may play a role. It has been reported that TSP-1 inhibited proliferation of A2058 malignant melanoma cells *in vitro*,⁴³ and certain TSP-1 effects seem to be specific for distinct cell lines. Our results, obtained by using three independent *in vitro* growth assays, strongly indicate that A431 cell growth is not influenced by TSP-1. This is further supported by the findings that cell proliferation rates of TSP-1 overexpressing A431 xenotransplants were indistinguishable from control transfected A431 xenotransplants, as measured by expression of the proliferation-related antigen PCNA. Similarly, tumor treatment with the angiogenesis inhibitor angiostatin also led to reduced tumor size without changing tumor cell proliferation rates.⁴⁴

The CD36 receptor has been reported to be the receptor on vascular endothelial cells that mediates the anti-angiogenic effects of TSP-1.³⁵ We could not detect differences in endothelial cell CD36 expression between control tumors and TSP-1 overexpressing tumors by immunostaining. However, TSP-1 overexpressing A431 tumors showed increased CD36 expression in tumor cells, as compared to no or little CD36 expression in control tumors. TSP-1 transfected A431 and SCC-13 cell clones also showed highly increased CD36 expression *in vitro*. Moreover, incubation of control A431 clones with conditioned media obtained from TSP-1 overexpressing clones significantly induced CD36 expression. These results suggest that soluble TSP-1 induces the CD36 receptor on tumor cells. They are in accordance with the recent findings that adhesion to TSP-1 induced CD36 in embryonic fibroblasts.⁴⁵ Our results further demonstrate that up-regulation of CD36 expression in TSP-1 transfected A431 cells moderately enhanced cell adhesion to immobilized TSP-1 *in vitro*, in accordance with a previous report dem-

onstrating that HT-1080 fibrosarcoma cells adhered to TSP-1 via their CD36 receptor. The functional consequences of enhanced tumor cell expression of CD36 *in vivo* remain unknown since we did not detect any differences in tumor cell proliferation between control and TSP-1/CD36 overexpressing A431 xenotransplants. However, it is conceivable that the CD36/TSP-1 system might play a role for the invasive or metastatic properties of malignant tumor cells *in vivo*.

Overexpression of TSP-1 in A431 xenotransplants resulted in extensive areas of tumor cell necrosis, possibly due to anti-angiogenic effects of TSP-1. To define the effects of TSP-1 on tumor vasculature, we performed morphometric analyses on CD31 stained tumor sections by using a semiautomated image analysis program. In TSP-1 overexpressing tumors, a moderate reduction of the microvascular density was observed, as measured by the average number of CD31-positive tumor vessels per mm². Similar results have been reported in human breast cancer xenotransplants.³⁶ However, when we compared vessel sizes, we found that the average vessel area was significantly diminished in TSP-1 overexpressing tumors with absence of larger vessels with surface areas of greater than 2,000 μm². These findings demonstrate that TSP-1 reduced the characteristically increased size of tumor vessels and suggest that measuring vascular densities alone may not be sufficient to detect changes in tumor vasculature. They also suggest that determining the total vascular area per unit area may serve as a more sensitive parameter to measure tumor angiogenesis.

It is of interest that the first vascular changes observed during treatment of experimental tumors with an antibody to the angiogenesis factor VEGF also consisted of a dramatic reduction of blood vessel diameters and tortuosity.⁴⁶ Moreover, overexpression of VEGF in the skin of transgenic mice³³ or in MEL-57 melanoma xenotransplants⁴ led to the development of tortuous and dilated blood vessels, and inhibition of the VEGF-inducible α1- and α2-integrins significantly inhibited VEGF-driven tumor angiogenesis *in vivo*, most prominently through reduction of average blood vessel diameters.⁴⁷ To exclude that the reduction in vessel sizes observed in TSP-1 overexpressing A431 xenotransplant tumors was due to down-regulation of VEGF expression, we performed *in situ* hybridizations of tumor xenotransplants. These studies demonstrated unchanged levels of VEGF mRNA expression in TSP-1 overexpressing tumors *versus* controls. Therefore, the reduction in vessel sizes reflects an important role of TSP-1 on the formation of tumor vasculature and demonstrates that similar vascular effects can be obtained by overexpression of TSP-1 or by inhibition of VEGF, suggesting antipodal roles of the two molecules in tumor angiogenesis. In summary, TSP-1 induced a potent growth inhibition of malignant epithelial skin cancer with complete inhibition of tumor development of highly differentiated SCC-13 tumors. The anti-tumoral effect of TSP-1 was not due to direct inhibition of tumor cell proliferation, but was associated with significant inhibition of tumor angiogenesis. Our results suggest that quantitation of both microvascular densities and of total vascular areas

provides a much more sensitive parameter for tumor angiogenesis than determination of vessel densities alone.

References

1. Hanahan D, Folkman J: Patterns and emerging mechanisms of the angiogenic switch during tumorigenesis. *Cell* 1996, 86:353-364
2. Brown LF, Detmar M, Claffey K, Nagy JA, Feng D, Dvorak AM, Dvorak HF: Vascular permeability factor/vascular endothelial growth factor: a multifunctional angiogenic cytokine. *Exs* 1997, 79:233-269
3. Ferrara N: The role of vascular endothelial growth factor in pathological angiogenesis. *Breast Cancer Res Treat* 1995, 36:127-137
4. Claffey KP, Brown LF, del Aguila LF, Tognazzi K, Yeo K-T, Manseau EJ, Dvorak HF: Expression of vascular permeability factor/vascular endothelial growth factor by melanoma cells increases tumor growth, angiogenesis, and experimental metastasis. *Cancer Res* 1996, 56:172-181
5. Skobe M, Rockwell P, Goldstein N, Vosseler S, Fusenig NE: Halting angiogenesis suppresses carcinoma cell invasion. *Nat Med* 1997, 3:1222-1227
6. Iruela-Arispe M, Bornstein P, Sage H: Thrombospondin exerts an antiangiogenic effect on cord formation by endothelial cells in vitro. *Proc Natl Acad Sci USA* 1991, 88:5026-5030
7. Volpert OV, Tolsma SS, Pellerin S, Feige JJ, Chen H, Mosher DF, Bouck N: Inhibition of angiogenesis by thrombospondin-2. *Biochem Biophys Res Commun* 1995, 217:326-332
8. O'Reilly MS, Holmgren L, Shing Y, Chen C, Rosenthal RA, Moses M, Lane WS, Cao Y, Sage EH, Folkman J: Angiostatin: a novel angiogenesis inhibitor that mediates the suppression of metastases by a Lewis lung carcinoma. *Cell* 1994, 79:315-328
9. O'Reilly MS, Boehm T, Shing Y, Fukai N, Vasios G, Lane WS, Flynn E, Birkhead JR, Olsen BR, Folkman J: Endostatin: an endogenous inhibitor of angiogenesis and tumor growth. *Cell* 1997, 88:277-285
10. Bornstein P: Diversity of function is inherent in matricellular proteins: an appraisal of thrombospondin 1. *J Cell Biol* 1995, 130:503-506
11. Tolsma SS, Volpert OV, Good DJ, Frazier WA, Polverini PJ, Bouck N: Peptides derived from two separate domains of the matrix protein thrombospondin-1 have anti-angiogenic activity. *J Cell Biol* 1993, 122:497-511
12. Varani J, Carey TE, Fligiel SEG, McKeever PE, Dixit VM: Tumor type-specific differences in cell-substrate adhesion among human tumor cell lines. *Int J Cancer* 1987, 39:397-403
13. Varani J, Dixit VM, Fligiel SE, McKeever PE, Carey TE: Thrombospondin-induced attachment and spreading of human squamous carcinoma cells. *Exp Cell Res* 1986, 167:376-390
14. Riser BL, Varani J, O'Rourke K, Dixit VM: Thrombospondin binding by human squamous carcinoma and melanoma cells: relationship to biological activity. *Exp Cell Res* 1988, 174:319-329
15. Yabkowitz R, Mansfield PJ, Dixit VM, Suchard SJ: Motility of human carcinoma cells in response to thrombospondin: relationship to metastatic potential and thrombospondin structural domains. *Cancer Res* 1993, 53:378-387
16. Albo D, Arnoletti JP, Castiglioni A, Granick MS, Solomon MP, Rothman VL, Tuszynski GP: Thrombospondin (TSP), and transforming growth factor β 1 (TGF-β) promote human A549 lung carcinoma cell plasminogen activator inhibitor type 1 (PAI-1) production, and stimulate tumor cell attachment in vitro: *Biochem Biophys Res Commun* 1994, 203:857-865
17. Wang TN, Qian XH, Granick MS, Solomon MP, Rothman VL, Tuszynski GP: The effect of thrombospondin on oral squamous carcinoma cell invasion of collagen. *Am J Surg* 1995, 170:502-505
18. Albo D, Berger DH, Wang TN, Hu X, Rothman V, Tuszynski GP: Thrombospondin-1 and transforming growth factor-β1 promote breast tumor cell invasion through up-regulation of the plasminogen/plasmin system. *Surgery* 1997, 122:493-499
19. Albo D, Berger DH, Tuszynski GP: The effect of thrombospondin-1 and TGF-β1 on pancreatic cancer cell invasion. *J Surg Res* 1998, 76:86-90
20. Castle V, Varani J, Fligiel S, Prochownik EV, Dixit V: Antisense-mediated reduction in thrombospondin reverses the malignant phe-

- notype of human squamous carcinoma. *J Clin Invest* 1991, 87:1883–1888
21. Tuszynski GP, Nicosia RF: The role of thrombospondin-1 in tumor progression and angiogenesis. *Bioessays* 1996, 18:71–76
22. Zabrenetzky V, Harris CC, Steeg PS, Roberts DD: Expression of the extracellular matrix molecule thrombospondin inversely correlates with malignant progression in melanoma, lung and breast carcinoma cell lines. *Int J Cancer* 1994, 59:191–195
23. Campbell SC, Volpert OV, Ivanovich M, Bouck NP: Molecular mediators of angiogenesis in bladder cancer. *Cancer Res* 1998, 58:1298–1304
24. Wight TN, Raugi GJ, Mumby SM, Bornstein P: Light microscopic immunolocalization of thrombospondin in human tissues. *J Histochem Cytochem* 1985, 33:295–302
25. Rheinwald JG, Beckett MA: Tumorigenic keratinocyte lines requiring anchorage and fibroblast support cultured from human squamous cell carcinomas. *Cancer Res* 1981, 41:1657–1663
26. Richard L, Velasco P, Detmar M: A simple immunomagnetic protocol for the selective isolation and long-term culture of human dermal microvascular endothelial cells. *Exp Cell Res* 1998, 240:1–6
27. Detmar M, Brown LF, Berse B, Jackman RW, Elicker BM, Dvorak HF, Claffey KP: Hypoxia regulates the expression of vascular permeability factor/vascular endothelial growth factor (VPF/VEGF) and its receptors in human skin. *J Invest Dermatol* 1997, 108:263–268
28. Schmidt R, Streit M, Kaiser R, Herzberg F, Schirner M, Schramm K, Kaufmann C, Henneken M, Schafer KM, Thiel E, Kreuser ED: De novo expression of the $\alpha 5 \beta 1$ -fibronectin receptor in HT29 colon-cancer cells reduces activity of C-SRC: increase of C-SRC activity by attachment on fibronectin. *Int J Cancer* 1998, 76:91–98
29. Laemmli UK: Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970, 227:680–685
30. Schirner M, Herzberg F, Schmidt R, Streit M, Schoening M, Hummel M, Kaufmann C, Thiel E, Kreuser ED: Integrin $\alpha 5 \beta 1$: a potent inhibitor of experimental lung metastasis. *Clin Exp Metastasis* 1998, 16:427–435
31. Gorczyca W, Gong J, Darzynkiewicz Z: Detection of DNA strand breaks in individual apoptotic cells by the in situ terminal deoxynucleotidyl transferase and nick translation assays. *Cancer Res* 1993, 53:1945–1951
32. Gillies RJ, Didier N, Denton M: Determination of cell number in monolayer cultures. *Anal Biochem* 1986, 159:109–113
33. Detmar M, Brown LF, Schön MP, Elicker BM, Velasco P, Richard L, Fukumura D, Monsky W, Claffey KP, Jain RK: Increased microvascular density and enhanced leukocyte rolling and adhesion in the skin of VEGF transgenic mice. *J Invest Dermatol* 1998, 111:1–6
34. Myoken Y, Kayada Y, Okamoto T, Kan M, Sato GH, Sato JD: Vascular endothelial cell growth factor (VEGF) produced by A-431 human epidermoid carcinoma cells and identification of VEGF membrane binding sites. *Proc Natl Acad Sci USA* 1991, 88:5819–5823
35. Dawson DW, Pearce SF, Zhong R, Silverstein RL, Frazier WA, Bouck NP: CD36 mediates the in vitro inhibitory effects of thrombospondin-1 on endothelial cells. *J Cell Biol* 1997, 138:707–717
36. Weinstat-Saslow DL, Zabrenetzky VS, VanHoutte K, Frazier WA, Roberts DD, Steeg PS: Transfection of thrombospondin 1 complementary DNA into a human breast carcinoma cell line reduces primary tumor growth, metastatic potential, and angiogenesis. *Cancer Res* 1994, 54:6504–6511
37. Castle VP, Dixit VM, Polverini PJ: Thrombospondin-1 suppresses tumorigenesis and angiogenesis in serum- and anchorage-independent NIH 3T3 cells. *Lab Invest* 1997, 77:51–61
38. Sheibani N, Frazier WA: Thrombospondin 1 expression in transformed endothelial cells restores a normal phenotype, and suppresses their tumorigenesis. *Proc Natl Acad Sci USA* 1995, 92:6788–6792
39. Hsu SC, Volpert OV, Steck PA, Mikkelsen T, Polverini PJ, Rao S, Chou P, Bouck NP: Inhibition of angiogenesis in human glioblastomas by chromosome 10 induction of thrombospondin-1. *Cancer Res* 1996, 56:5684–5691
40. Volpert OV, Lawler J, Bouck NP: A human fibrosarcoma inhibits systemic angiogenesis and the growth of experimental metastases via thrombospondin-1. *Proc Natl Acad Sci USA* 1998, 95:6343–6348
41. Varani J, L RB, Hughes LA, Carey TE, Fligiel SE, Dixit VM: Characterization of thrombospondin synthesis, secretion and cell surface expression by human tumor cells. *Clin Exp Metastasis* 1989, 7:265–276
42. Castle VP, Ou X, O'Rourke K, Dixit VM: High level thrombospondin 1 expression in two NIH 3T3 cloned lines confers serum- and anchorage-independent growth. *J Biol Chem* 1993, 268:2899–2903
43. Guo N-h, Zabrenetzky VS, Chandrasekaran L, Sipes JM, Lawler J, Kruttsch HC, Roberts DD: Differential roles of protein kinase C and pertussis toxin-sensitive G-binding proteins in modulation of melanoma cell proliferation and motility by thrombospondin 1. *Cancer Res* 1998, 58:3154–3162
44. O'Reilly MS, Holmgren L, Chen C, Folkman J: Angiostatin induces and sustains dormancy of human primary tumors in mice. *Nat Med* 1996, 2:689–692
45. Stomski FC, Gani JS, Bates RC, Burns GF: Adhesion to thrombospondin by human embryonic fibroblasts is mediated by multiple receptors and includes a role for glycoprotein 88 (CD36). *Exp Cell Res* 1992, 198:85–92
46. Yuan F, Chen Y, Dellian M, Safabakhsh N, Ferrara N, Jain RK: Time-dependent vascular regression and permeability changes in established human tumor xenografts induced by an anti-vascular endothelial growth factor/vascular permeability factor antibody. *Proc Natl Acad Sci USA* 1996, 93:14765–14770
47. Senger DR, Claffey KP, Benes JE, Perruzzi CA, Sergiou AP, Detmar M: Angiogenesis promoted by vascular endothelial growth factor: regulation through $\alpha 1 \beta 1$, and $\alpha 2 \beta 1$ integrins. *Proc Natl Acad Sci USA* 1997, 94:13612–13617