WHEN SHOULD WE PAY ATTENTION TO UNFAVORABLE NEWS FROM PREGNANCY REGISTRIES?

In Utero Antiepileptic Drug Exposure: Fetal Death and Malformations. Meador KJ, Baker GA, Finnell RH, Kalayjian LA, Liporace JD, Loring DW, Mawer G, Pennell PB, Smith JC, Wolff MC; NEAD Study Group. *Neurology* 2006;67:407–412.

BACKGROUND: Pregnancy outcomes following in utero exposure to antiepileptic drugs (AEDs) are uncertain, limiting an evidenced-based approach. **OBJECTIVE:** To determine if fetal outcomes vary as a function of different in utero AED exposures. **METHODS:** This ongoing prospective observational study across 25 epilepsy centers in the United States and United Kingdom enrolled pregnant women with epilepsy from October 1999 to February 2004 to determine if differential long-term cognitive and behavioral neurodevelopmental effects exist across the four most commonly used AEDs. This initial report focuses on the incidence of serious adverse outcomes including major congenital malformations (which could be attributable to AEDs) or fetal death. A total of 333 mother/child pairs were analyzed for monotherapy exposures: carbamazepine (n = 110), lamotrigine (n = 98), phenytoin (n = 56), and valproate (n = 69). **RESULTS:** Response frequencies of pregnancies resulting in serious adverse outcomes for each AED were as follows: carbamazepine 8.2%, lamotrigine 1.0%, phenytoin 10.7%, and valproate 20.3%. Distribution of serious adverse outcomes differed significantly across AEDs and was not explained by factors other than in utero AED exposure. Valproate exhibited a dose-dependent effect. **CONCLUSIONS:** More adverse outcomes were observed in pregnancies with in utero valproate exposure versus the other antiepileptic drugs (AEDs). These results combined with several recent studies provide strong evidence that valproate poses the highest risk to the fetus. For women who fail other AEDs and require valproate, the dose should be limited if possible.

COMMENTARY

The study by Meador and colleagues, who form the Neurodevelopmental Effects of Antiepileptic Drugs (NEAD) study group, is one of a number of recently published studies that addresses the impact of AEDs on infants born to women receiving them during pregnancy (1–5). Of interest, the study was funded by the National Institutes of Health for a completely different purpose, namely to address the impact of in utero AED exposure on the cognitive development of children born to mothers with epilepsy. The study is ongoing, and the cognitive outcome will be reported at a future date. Reported here are the important pregnancy outcomes, major fetal malformation or fetal death, associated with specific AEDs. This study confirms the finding of several other registries that the incidence of these adverse pregnancy outcomes is increased in the offspring of mothers receiving valproate (1–5).

How should physicians respond to information coming at an ever-increasing rate from a number of registries? When an adverse outcome is reported to be at a greater frequency for a particular AED, should the response be to stop prescribing that drug to women of childbearing age? The answer to this question is not so simple. One problem is that the stud-

ies and registries have provided conflicting information. For example, the North American pregnancy registry recently reported an increased incidence of cleft lip/cleft palate associated with the use of lamotrigine, with an incidence of approximately 8.9/1,000 cases, as compared with 0.5 to 2.16/1,000 in nonepileptic mothers not receiving AEDs (6). Yet, an increased incidence of this malformation was not seen in any of the other registries or in a large prospective lamotrigine pregnancy registry (2). Confusing the matter further, the British pregnancy registry found a correlation between the likelihood of major fetal malformations and higher lamotrigine dose (1). This correlation was not found in other registries, not even in those that had enrolled a large enough number of women with lamotrigine exposure to provide the power to detect a dose-response relationship.

How, then, do we account for these differences? As with every other type of investigation, details matter. If the studies are carefully reviewed, subtle differences in design begin to emerge that might explain the differing results. For example, there are disparities in how patients are identified for inclusion. In some registries, patients can only be included if they either have had no perinatal testing at the time of enrollment or have had negative testing up to that point. Since patients often do not enroll until their 5th or 6th month of pregnancy, this protocol would exclude patients who had early evidence of fetal malformation, thereby potentially lowering the incidence of malformations

identified in the study. Other registries require that women self-identify for enrollment, after being provided a phone number from their physician. It is the experience of many physicians that only a fraction of women who are told about a registry will actually call to enroll. This fact might increase the representation of motivated women, who might differ in important ways from women who do not enroll. In other registries, women are enrolled by their physician, possibly providing a broader sampling of the population.

Differences in the timing of malformation ascertainment also might factor into the contradictory results seen across registries. Some registries only assessed at birth, while others followed children for months to years after delivery. Control groups also may differ and often are not representative of the type of patients enrolled in the registry. For example, the lamotrigine pregnancy registry compared outcomes of women enrolled throughout the United States with outcomes of patients seen at the Center for Disease Control's Metropolitan Atlanta Congenital Defects Program—even though the Metropolitan Atlanta population probably was not equivalent to the United States as a whole in terms of race, socioeconomic status, or exposure to other potential teratogens, such as cigarettes and alcohol (2). If the control population had a relatively high rate of major malformations compared to the expected rate of the women enrolling in the registry, malformations in the women exposed to AEDs might be missed. In this respect, comparisons between two groups of women in the same registry who have been exposed to different drugs (e.g., women exposed to carbamazepine versus valproic acid) might have greater validity than comparisons to the external control group. Even within the same registry, it is impossible to exclude population bias related to nonrandomized selection of AEDs. For example, if the genetic abnormality that confers risk for juvenile myoclonic epilepsy also confers a higher risk of major malformations in offspring, there might be a bias against valproic acid, a drug of choice for that condition.

The NEAD study was an ideal study from the perspective of several of the issues reviewed here. Infants were followed for several years after birth, which would clearly increase the likelihood of discovering abnormalities that are not immediately apparent postpartum. In contrast to some of the other registries, the association of malformations with important potential confounders, such as socioeconomic status, age of the mother, epilepsy syndrome, AED dose, and compliance, was explored and eliminated in the NEAD study. Drugs were compared to each other, obviating the need for an external comparison group. In addition, rather than being given a national registry phone number to call, patients were asked by their own

physicians to provide consent, which may have increased the likelihood of a representative sample of women. The weakness of the study lies in the relatively low number of patients enrolled. This factor might reduce the importance of the study's finding that there were no differences in rates of fetal death or major malformation among three of the AEDs studied, namely carbamazepine, phenytoin, and lamotrigine. However, the fact that valproic acid was found to convey a higher risk, despite the relatively small sample size, only highlights the importance of the effect. Therefore, the results of this study should be given careful consideration.

In summary, because of differences among pregnancy registries described here and because, for obvious reasons, a randomized trial is impossible, it would be reasonable to wait until a finding has been confirmed in several studies before altering drug selection in women contemplating pregnancy. The fact that valproic acid has been associated with a significantly higher risk of malformations in six studies means that the signal is strong enough to warrant consideration of selecting an alternative AED, when a choice exists. A more thorny issue arises when valproic acid has conveyed seizure control, and conversion to alternative therapy may be associated with a risk of seizure breakthrough. In this case, discussion of the risk/benefit relationship is essential, at the very least.

by Jacqueline A. French, MD

References

1. Morrow J, Russell A, Guthrie E, Parsons L, Robertson I, Waddell R, Irwin B, McGivern RC, Morrison PJ, Craig J. Malformation risks of antiepileptic drugs in pregnancy: a prospective study from the UK Epilepsy and Pregnancy Register. *J Neurol Neurosurg Psychiatry* 2006;77:193–198.
2. Cunnington M, Tennis P. International Lamotrigine Pregnancy Registry Scientific Advisory Committee. Lamotrigine and the risk of malformations in pregnancy. *Neurology* 2005;64:955–960.
3. Wyszynski DF, Nambisan M, Surve T, Alsdorf RM, Smith CR, Holmes LB. Antiepileptic Drug Pregnancy Registry. Increased rate of major malformations in offspring exposed to valproate during pregnancy. *Neurology* 2005;64:961–965.
4. Wide K, Winbladh B, Kallen B. Major malformations in infants exposed to antiepileptic drugs in utero, with emphasis on carbamazepine and valproic acid: a nation-wide population-based register study. *Acta Paediatr* 2004;93:174–176.
5. Vajda FJ, Eadie MJ. Maternal valproate dosage and foetal malformations. *Acta Neurol Scand* 2005;112:137–143.
6. Holmes LB, Wyszynski DF, Baldwin EJ, Habecker E, Glassman LH, Smith CR. Increased risk for non-syndromic cleft palate among infants exposed to lamotrigine during pregnancy (abstract). *Birth Defects Research Part A: Clinical and Molecular Teratology* 2006;76:318.

IT'S ABOUT TIME: LONG-TERM MEMORY OUTCOME FOLLOWING TEMPORAL LOBECTOMY

Verbal Memory Decline after Temporal Epilepsy Surgery?: A 6-Year Multiple Assessments Follow-up Study.

Alpherts WC, Vermeulen J, van Rijen PC, da Silva FH, van Veelen CW; Dutch Collaborative Epilepsy Surgery Program. *Neurology* 2006;67:626–631. **OBJECTIVE:** To assess the long-term effects of temporal lobe epilepsy surgery on verbal memory. **METHODS:** We assessed verbal memory performance as measured by a verbal learning test ("15 Words Test," a Dutch adaptation of Rey's Auditory Verbal Learning Test) before surgery and at three specific times after surgery: 6 months, 2 years, and 6 years in 85 patients (34 left temporal lobe [LTL] vs. 51 right temporal lobe [RTL]). An amygdalo-hippocampectomy and a neocortical temporal resection between 2.5 and 8 cm were carried out in all patients. **RESULTS:** LTL patients showed an ongoing memory decline for consolidation and acquisition of verbal material (both 2/3 SDs) for up to 2 years after surgery. RTL patients at first showed a gain in both memory acquisition and consolidation, which vanished in the long term. Breaking the group up into a mesiotemporal (MTS) group and a non-MTS group showed clear differences. The group with pure MTS showed an overall lower verbal memory performance than the group without pure MTS, in the LTL group more pronounced than in the RTL group. After surgery, both pathology groups showed an ongoing decline for up to 2 years, but the degree of decline was greater for the LTL patients with MTS compared with the non-MTS group. Becoming and remaining seizure-free after surgery does not result in a better performance in the long term. Predictors of postoperative verbal memory performance at 6 years after surgery were side of surgery, preoperative memory score, and age. **CONCLUSIONS:** The results provide evidence for a dynamic decline of verbal memory functions up to 2 years after left temporal lobectomy, which then levels off.

COMMENTARY

Both poorly controlled epilepsy and its treatment are associated with varying degrees of neuropsychological difficulty. Cognitive impairment may result from the underlying neurologic substrate giving rise to a patient's epilepsy, the negative effects of repeated seizures, or the effects associated with antiepileptic drug therapy, to name just a few important factors. Whether negative long-term or cumulative seizure effects on cognition exist is still a subject of debate (1); however, there is little doubt that successful epilepsy treatment that either decreases or eliminates factors associated with impaired cognition (e.g., becoming seizure-free, decreasing/eliminating antiepileptic drugs) can result in improved performance on neuropsychological tests.

For patients with localization-related epilepsy whose seizures cannot be adequately managed with medications, the best hope of successful therapy is surgical resection. Given the importance of the hippocampus and associated medial temporal lobe structures in the acquisition and formation of new memory traces, it follows that when seizures arise from the medial temporal lobe, there is a risk of significant postoperative memory decline; this risk is greater for patients undergoing surgery in the language-dominant hemisphere. Although multiple factors contribute to the surgical decision-making process, memory risks are strongly considered in patient selection and should be

viewed against the backdrop of expected cognitive change in the absence of surgical intervention. In both surgical and nonsurgical scenarios, an important factor is how neuropsychological performance may change over time. In most research studies, time is measured in increments of months and not years, and it is rare that multiple assessments are performed. Consequently, memory change typically is conceptualized as a static rather than dynamic process, and how cognition evolves over time while interacting with related factors, such as preoperative performance levels (2) or patient age (3), is incompletely understood.

Most reports describing post-temporal lobectomy memory change have examined relatively short periods of postoperative follow-up, typically on the order of 6–12 months in duration. As with any CNS lesion, there are multiple determinants of postsurgical functional outcome, including both the functional status of the tissue involved in the surgical resection (functional adequacy) and the functional status of remaining brain structures (functional reserve), particularly of the homologous brain regions to those included in surgery. Both components may be associated with a variety of effects, some of which may be beneficial (e.g., plasticity, leading to functional reorganization) and others that may be detrimental (e.g., degeneration from deafferentation).

When longer postsurgical intervals have been used to study memory change following anterior temporal lobectomy (ATL), it appears that a single follow-up assessment during the first postoperative year does not fully characterize longer-term memory outcomes. In a study of memory, using one of the longest (9–13 years) postsurgical follow-up periods included in any investigation, Rausch et al. observed that verbal memory

continued to decline beyond the first postoperative year (4). However, in addition to studying a relatively small patient cohort, Rausch et al. did not include preoperative MRI findings to characterize patients according to whether or not they displayed evidence of mesial temporal sclerosis (MTS). There were no cognitive evaluations between the initial postoperative assessment performed 1 year after surgery and the long-term follow-up; thus, two issues that could not be addressed were (i) the ongoing deterioration in verbal memory following left ATL and (ii) the time point at which the risk of progressive postoperative memory begins to wane. Also not addressed was the question of whether the presence or absence of MTS, which is a strong predictor of memory change following ATL, was associated with different temporal patterns of memory change following surgery.

The recent report by Alpherts and colleagues reveals a dynamic process of ongoing memory changes that, like the Rausch et al. study, suggests continuing memory decline beyond the first postoperative year. Unlike in the Rausch report, however, progressive postoperative memory deterioration was found to stabilize between the second and sixth postoperative year. As expected, Alpherts et al. observed significant memory decline in left ATL patients when tested 6 months after surgery. Right ATL patients had higher verbal memory scores at their 6-month follow-up compared to preoperative scores. Although memory improvement in right ATL patients is often attributed to practice effects from repeated exposure to the same memory stimuli, these results cannot easily be attributed to a simple practice effect since alternative memory test forms were used. When tested after 2 years, further verbal memory decline was observed in left ATL patients, while verbal memory in right ATL patients returned to preoperative levels. There appears to be a progression of verbal memory impairment following left ATL, even in patients who were rendered seizure-free. No further deterioration in verbal memory was reported when patients were tested during their 6-year follow-up. The progressive decline in verbal memory extending beyond 6 months was attributed to shrinkage of the hippocampal remnant, which has been shown to decrease in size throughout the first preoperative year.

It is when trying to understand the behavior of specific subgroups that many studies fall victim to small sample sizes. For example, using traditional statistical approaches, patients undergoing removal of a healthy, nonatrophied left hippocampus, and those with preoperative evidence of MTS both displayed comparable declines in verbal memory. Although this finding suggests a similar pattern of 2-year decline, regardless of whether MTS was present or not, comparison of the group means suggests that the outcome actually may have resulted from inadequate statistical power. When looking at delayed recall performance at each follow-up assessment, a differential pattern of memory change is suggested. From the second to the sixth

year follow-up, patients with left MTS improved on delayed recall (from 6.6/15 words to 7.3/15 words), whereas patients without MTS continued to decline slightly (from 8.2/15 words to 7.7/15 words). Although this effect was not statistically significant, it is consistent with literature suggesting increased risk for memory decline in patients with larger hippocampal volumes. However, the finding contrasts with one of the main conclusions put forth by Alpherts et al., which is that, based on a verbal learning performance summed across trials (rather than delayed verbal memory recall), the degree of memory decline is greater in left ATL patients with MTS than in those without MTS. Although a small sample, examination of delayed verbal memory performances raises the possibility that patients with left MTS may continue to experience verbal memory decline extending beyond the 2-year assessment point. Because nonsurgical patients with left temporal lobe epilepsy may continue to experience cognitive decline that is unrelated to seizure occurrence or antiepilepsy drug therapy, it is unclear whether the progressive memory decline seen in some left ATL patients more than 2 years after surgery simply reflects the progressive nature of the disease or some other currently unidentified mechanism.

The results of the Alpherts et al. study raise important issues that future reports might be careful to address. For example, is a similar dynamic pattern of postoperative change seen in other neuropsychological measures, in particular, in confrontation naming? Historically, confrontation-naming ability is considered a measure of semantic memory, reflecting general knowledge rather than the acquisition of new context-specific information, yet confrontation naming also is related to hippocampal structure (5) and function (6). In fact, confrontation-naming ability is a robust measure of dysfunction associated with left temporal lobe epilepsy (7,8), with well-established declines following left temporal lobectomy regardless of surgical technique (9). Thus, whether confrontation-naming declines follow the same dynamic time course as does delayed recall will be of considerable interest.

The inclusion of only surgical patients in the Alpherts et al. study resulted in primarily one issue being addressed, that is, the effects of left- versus right-sided ATL on neuropsychological outcomes. Thus, in order to study how memory changes over a similar time interval and to provide information regarding the relative risks of postsurgical memory change (4), it will be necessary to design a study with an appropriately matched nonsurgical control group. A control group of patients without a chronic neurological disease is necessary in any longitudinal study designed to assess ATL for which test-retest effects may exist. As reviewed in the previous issue of *Epilepsy Currents*, Hermann and colleagues determined that without such a control group, lack of cognitive decline on retest (i.e., of the expected practice effect) may incorrectly be interpreted as an

absence of cognitive deterioration, when in fact there is negative cognitive change (10,11). Such studies will provide valuable information for clinicians and patients to consider when evaluating treatment options, while simultaneously maximizing quality of life and overall cognitive health. Neuropsychological assessments performed only during the first postoperative year, appear inadequate to properly characterize longer-term neuropsychological outcomes following ATL.

by David W. Loring, PhD

References

1. Dodrill CB. Neuropsychological effects of seizures. *Epilepsy & Behavior* 2004;5(suppl. 1):21–24.
2. LoGalbo A, Sawrie S, Roth DL, Kuzniecky R, Knowlton R, Fought E, Martin R. Verbal memory outcome in patients with normal preoperative verbal memory and left mesial temporal sclerosis. *Epilepsy & Behavior* 2005;6:337–341.
3. Helmstaedter C, Reuber M, Elger CC. Interaction of cognitive aging and memory deficits related to epilepsy surgery. *Ann Neurol* 2002;52:89–94.
4. Rausch R, Kraemer S, Pietras CJ, Le M, Vickrey BG, Passaro EA. Early and late cognitive changes following temporal lobe surgery for epilepsy. *Neurology* 2003;60:951–959.
5. Seidenberg M, Geary E, Hermann B. Investigating temporal lobe contribution to confrontation naming using MRI quantitative volumetrics. *J Int Neuropsychol Soc* 2005;11:358–366.
6. Sawrie SM, Martin RC, Gilliam FG, Fought RE, Maton B, Hugg JW, Bush N, Sinclair K, Kuzniecky RI. Visual confrontation naming and hippocampal function: a neural network study using quantitative (1)H magnetic resonance spectroscopy. *Brain* 2000;123(Pt 4):770–780.
7. Busch RM, Frazier TW, Haggerty KA, Kubu CS. Utility of the Boston naming test in predicting ultimate side of surgery in patients with medically intractable temporal lobe epilepsy. *Epilepsia* 2005;46:1773–1779.
8. Raspall T, Donate M, Boget T, Carreno M, Donaire A, Agudo R, Bargallo N, Rumia J, Setoain X, Pintor L, Salameiro M. Neuropsychological tests with lateralizing value in patients with temporal lobe epilepsy: reconsidering material-specific theory. *Seizure* 2005;14:569–576.
9. Hermann BP, Perrine K, Chelune GJ, Barr W, Loring DW, Strauss E, Trenerry MR, Westerveld M. Visual confrontation naming following left anterior temporal lobectomy: a comparison of surgical approaches. *Neuropsychology* 1999;13:3–9.
10. Hermann BP, Seidenberg M, Dow C, Jones J, Rutecki P, Bhat-tacharya A, Bell B. Cognitive prognosis in chronic temporal lobe epilepsy. *Ann Neurol* 2006;60:80–87.
11. Harden CL. New Evidence Supports Cognitive Decline in Temporal Lobe Epilepsy. *Epilepsy Curr* 2007; 7: 12–14.

VACCINES, ENCEPHALOPATHIES, AND MUTATIONS

De-Novo Mutations of the Sodium Channel Gene *SCN1A* in Alleged Vaccine Encephalopathy: A Retrospective Study. Berkovic SF, Harkin L, McMahon JM, Pelekanos JT, Zuberi SM, Wirrell EC, Gill DS, Iona X, Mulley JC, Scheffer IE. *Lancet Neurol* 2006;5:488–492. **BACKGROUND:** Vaccination, particularly for pertussis, has been implicated as a direct cause of an encephalopathy with refractory seizures and intellectual impairment. We postulated that cases of so-called vaccine encephalopathy could have mutations in the neuronal sodium channel $\alpha 1$ subunit gene (*SCN1A*) because of a clinical resemblance to severe myoclonic epilepsy of infancy (SMEI) for which such mutations have been identified. **METHODS:** We retrospectively studied 14 patients with alleged vaccine encephalopathy in whom the first seizure occurred within 72 h of vaccination. We reviewed the relation to vaccination from source records and assessed the specific epilepsy phenotype. Mutations in *SCN1A* were identified by PCR amplification and denaturing high performance liquid chromatography analysis, with subsequent sequencing. Parental DNA was examined to ascertain the origin of the mutation. **RESULTS:** *SCN1A* mutations were identified in 11 of 14 patients with alleged vaccine encephalopathy; a diagnosis of a specific epilepsy syndrome was made in all 14 cases. Five mutations predicted truncation of the protein and six were missense in conserved regions of the molecule. In all nine cases where parental DNA was available the mutations arose de novo. Clinical-molecular correlation showed mutations in eight of eight cases with phenotypes of SMEI, in three of four cases with borderline SMEI, but not in two cases with Lennox–Gastaut syndrome. **CONCLUSION:** Cases of alleged vaccine encephalopathy could in fact be a genetically determined epileptic encephalopathy that arose de novo. These findings have important clinical implications for diagnosis and management of encephalopathy and, if confirmed in other cohorts, major societal implications for the general acceptance of vaccination.

COMMENTARY

The report by Berkovic and colleagues forms a nexus between two areas of great controversy in medical research.

The first area of controversy involves the value epilepsy syndromes for use in research and clinical practice. Detractors of the concept of identifying epilepsy syndromes argue that distinguishing specific forms of epilepsy is of little value because (i) only highly trained specialists have the skills and knowledge to identify these syndromes and (ii) most people with epilepsy do not fit into the already recognized syndromic categories. This posture is increasingly untenable in the face of all the advances in genetics that, coupled with precise phenotyping (i.e., recognition and specification of clinical syndromes), are providing an understanding of the genetic basis and associated molecular mechanisms underlying specific and recognizable forms of epilepsy syndromes. Severe myoclonic epilepsy of infancy (SMEI)—also called Dravet syndrome after Dr. Charlotte Dravet, who described the phenomenon—is an example of the value of a syndromic approach in epilepsy. Prior to Dr. Dravet's work, infants with this disorder simply had "bad epilepsy." She described the specific patterns that identify this syndrome (1). Mutations in the *SCN1A* gene coding for part of the voltage-gated sodium channel were first implicated as causal in a range of epilepsy syndromes often referred to as the generalized epilepsy with febrile seizure plus (GEFS+) spectrum (2). Investigators then linked Dravet syndrome to mutations in the *SCN1A* gene (3), thus extending the potential phenotypes of the GEFS+ cluster. Recent analyses indicate that specific manifestation of the epilepsy, including its severity, depends in part on the type of mutation (4). At least one other gene has been implicated in Dravet syndrome, *GABRG2* (5). The association with *SCN1A*, however, is so powerful and so specific that it is now the basis of a genetic diagnostic test.

The second area of controversy concerns severe encephalopathic conditions of infancy and childhood and the possibility that they might be caused by routine childhood vaccinations. In the United States, infants and children receive a large number of vaccinations during the first 18 months of life, including four hepatitis B, four diphtheria–tetanus–pertussis, three polio, one measles–mumps–rubella, one varicella, and four pneumococcal administrations. Encephalopathic conditions, such as that caused by Dravet syndrome (as well as several other specific and nonspecific forms of infant–childhood conditions), often have their initial onset during this 18-month window. Despite numerous studies that have failed to find any positive association between vaccination and encephalopathy, no study can absolutely prove the negative. At best, the Ad Hoc Committee for the Child Neurology Society was willing to conclude in their consensus statement that: "At the present time (1991), there is no means by which a diagnosis of pertussis vaccine-encephalopathy can be established in an individual case." (6) Despite the enormous positive public health impact of childhood vaccinations, on rare occasion, a child develops an encephalopathic condition in close temporal proximity to the receipt of a vac-

cine. These rare occurrences provide an emotionally compelling basis for inferring causality that is hard to dispel, even with the overwhelming quantity of negative data and consensus reports. This phenomenon has led to fear of vaccinations, a flood of lawsuits, and ultimately passage in the United States of the 1986 National Childhood Vaccine Injury Act to provide compensation to alleged victims of vaccine injury. Governments elsewhere have taken other measures (7).

Berkovic and colleagues now shed some light on this vexing problem. In their report, they identify 14 children from a larger series of children with unexplained encephalopathies. These 14 patients met the criteria for vaccine-related injury because of the temporal association between receipt of pertussis vaccine and onset of symptoms (within 72 h) as well as the lack of any other explanation. All 14 children could be assigned a specific syndromic diagnosis. All 11 of the children with SMEI or SMEI-B had mutations in the *SCN1A* gene that, alone, likely explain their neurological conditions. Not all children with encephalopathies following vaccination will have *SCN1A* mutations, as other genes have been implicated in SMEI/Dravet syndrome. There are many efforts under way to understand the genetic basis of other forms of otherwise unexplained childhood encephalopathies, such as West and Lennox–Gastaut syndromes.

The findings of Berkovic et al. are important for at least three reasons. First, they provide a substantial, alternative explanation of cause in cases of alleged vaccine-induced encephalopathy—something the mountains of negative epidemiological data alone could not do. These findings serve as important reassurance concerning the safety of the pertussis vaccine, at least with respect to childhood encephalopathy. Second, this study is a superb demonstration of the power of the syndromic approach to studying and understanding epilepsy and of the persuasive explanatory capabilities of genetic analysis. Without this well-defined phenotype, the role of the *SCN1A* gene likely never would have been recognized, and there would still be the same uncertainty regarding the potential adverse consequences of vaccination. Replication of the findings, of course, is essential. Yet, as it stands, this first report is extremely compelling. Third, the concern over the potential association between vaccines—particularly those with thimerosal, a preservative used in some vaccines—and other disorders of childhood, most notably autism, has gained in momentum during the past several years. The report by Berkovic et al. does not address this association; however, it provides proof of principle that these disorders of early childhood, which many are convinced are caused by vaccinations, may have an entirely different and unrelated cause.

Why is correctly attributing the cause of vaccine-related disorders important? Some parents refuse to have their children vaccinated. This trend has resulted in outbreaks of diseases that

were supposedly eradicated in the developed world. Epidemic outbreaks, such as the recent measles outbreak in Indiana (8) and the huge spike in pertussis cases seen in Sweden after whole cell vaccination stopped in the 1970s, highlight the repercussions of not vaccinating children (7). Measles can have severe and fatal complications, including subacute sclerosing panencephalitis. Pertussis is not a benign disease and has a 0.6% fatality rate. Encephalopathy also may occur in 0.9% of cases (9). These are well-documented and defined risks. The reactions to the perceived dangers of vaccination lead to these risks being realized. Solid, high-quality evidence, such as that presented by Berkovic and colleagues, in addition to the fundamental scientific contribution they represent, may have an important, positive public health impact by allaying fears, at the very least, about the role of pertussis vaccine in causing encephalopathy.

by Anne T. Berg, PhD

References

1. Dravet Ch, Bureau M, Oguni H, Fukuyama Y, Cokar O. Severe myoclonic epilepsy in infancy (Dravet Syndrome). In: Roger J, Bureau M, Dravet Ch, Genton P, Tassinari CA, Wolf P, eds. *Epileptic Syndromes in Infancy, Childhood and Adolescence*. Eastleigh, UK: John Libbey & Co., 2002, 81–103.
2. Scheffer IE, Berkovic SF. Generalized epilepsy with febrile seizures plus a genetic disorder with heterogeneous clinical phenotypes. *Brain* 1997;120:479–490.
3. Claes L, Del-Favero J, Ceulemans B, Lagae L, Van Broeckhoven C, De Jonghe P. De novo mutations in the sodium-channel gene *SCN1A* cause severe myoclonic epilepsy of infancy. *Am J Hum Genet* 2001;68:1327–1332.
4. Kanai K, Hirose S, Oguni H, Fukuma G, Shirasaka Y, Miyajima T, Wada K, Iwasa H, Yasumoto S, Matsuo M, Ito M, Mitsudome A, Kaneko S. Effect of localization of missense mutations in *SCN1A* on epilepsy phenotype severity. *Neurology* 2004;63:329–334.
5. Bonanni P, Malcarne M, Moro F, Veggioni P, Buti D, Ferrari AR, Parrini E, Mei D, Volzone A, Zara F, Heron SE, Bordo L, Marini C, Guerrini R. Generalized epilepsy with febrile seizures plus (GEFS): clinical spectrum in seven Italian families unrelated to *SCN1A*, and *GABRG2* gene mutations. *Epilepsia* 2004;45:149–158.
6. Pertussis immunization and the central nervous system. Ad Hoc Committee for the Child Neurology Society Consensus Statement on Pertussis Immunization and the Central Nervous System. *Ann Neurol* 1991;29:458–460.
7. Gangarosa REJ, Galazka AM, Wolfe CR, Phillips LM, Gangarosa RE, Miller E, Chen RT. Impact of anti-vaccine movements on pertussis control: the untold story. *Lancet* 1998;351:356–361.
8. Parker AA, Staggs W, Dayan GH, Ortega-Sanchez IR, Rota PA, Lowe L, Boardman P, Teclaw R, Graves C, Le Baron CW. Implications of a 2005 measles outbreak in Indiana for sustained elimination of measles in the United States. *N Engl J Med* 2006;355:447–455.
9. Diphtheria, tetanus, and pertussis: recommendations for vaccine use and other preventive measures, recommendations of the Immunization Practice Advisory Committee. *MMWR* 1991;40:1–28.

REVISITING THE WELL-ESTABLISHED SUBJECT OF STROKE AND EPILEPSY

The Epidemiology of Post-Stroke Epilepsy According to Stroke Subtypes. Benbir G, Ince B, Bozluolcay M. *Acta Neurol Scand* 2006;114:8–12. **OBJECTIVES:** Strokes represent the most common etiology of epilepsy in patients over the age of 60 years, with an incidence of 2–4% occurring in different studies. **MATERIALS AND METHODS:** In this observational study, 1,428 patients were included who had stroke and were admitted to our Stroke Unit between the years 1996 and 2005. **RESULTS:** Overall, 51 patients had post-stroke epilepsy (3.6%). Post-ischemic epilepsy occurred in 70.6% of the patients, post-hemorrhagic epilepsy occurred in 21.6% of the patients and epilepsy following venous infarctions occurred in 7.8% of all post-stroke epilepsy patients. Of 1,327 patients having ischemic stroke, 36 patients (2.7%), 11 out of 86 patients with hemorrhagic stroke (12.8%) and 4 of 15 patients with venous infarctions (26.6%) developed epilepsy. Compared with stroke patients without epilepsy, hemorrhagic ($P < 0.001$) and venous infarctions were more common in patients with post-stroke epilepsy ($P < 0.001$). The right hemisphere and the middle cerebral artery (MCA) territory were most commonly observed in ischemic and hemorrhagic stroke patients. **CONCLUSIONS:** Our results indicate that post-stroke epilepsy is more common among patients who have experienced venous infarctions. Hemorrhagic and venous infarctions are more commonly encountered in post-stroke epilepsy patients. Atherosclerotic and cardioembolic strokes were similar to those that occurred in post-stroke epilepsy patients. Localizations in post-stroke epileptic patients showed that the majority occurred in the right hemisphere, in the territory of the MCA. However, prospective, multicentered studies are needed for a better understanding of the epidemiology and social impact of post-stroke epilepsy.

COMMENTARY

Since the time of Hippocrates and of Jackson and Gowers in the mid-19th century, stroke has been reported to

be a cause of seizures and epilepsy, particularly in the elderly. Subsequently, the association has been confirmed by numerous different studies (1–5). Are there new developments on this previously studied topic? In this regard, the study of Benbir et al. does address some important issues. The authors performed a well-designed study, noteworthy for their inclusion of three elements. First, they used the guidelines developed by the International League Against Epilepsy to define epilepsy (6), which in turn underscores the ongoing need for a definition of epilepsy. Second, problems with acute seizures (which can arise from a metabolic disarrangement or other reversible insult) were eliminated altogether by excluding patients with provoked seizures, thus increasing the validity of the study. Third and most important, the authors included patients with venous infarctions, particularly dural sinus thrombosis—patients typically not included in such studies.

Cerebral venous thrombosis is a well-known cause of seizures. However, evidence for this finding comes from descriptive studies or case series in which seizures were part of the clinical presentation (7). Thus far, no prospective studies have been performed to evaluate the development of epilepsy in patients who have venous infarctions. Benbir and colleagues provide some of the best systematic documentation that this type of stroke (i.e., cerebral venous thrombosis) is associated with the development of epilepsy. It is known that a prothrombotic risk factor is identified in about 85% of patients with sinus thrombosis. In the other 15% of cases, sinus thrombosis typically is caused by a precipitating factor (such as a head injury or obstetrical delivery). Some of these conditions are considered risk factors not only for stroke but also for seizures and epilepsy. Another risk factor for poststroke epilepsy identified by Benbir et al. is that all patients who developed epilepsy had a late-onset seizure (2 weeks after stroke), extending the findings of the Seizures After Stroke Study Group published in 2000 (4), which found that that epilepsy occurred in up to 55% of patients with late-onset seizures after ischemic strokes.

Using the Trial of Org 10172 in Acute Stroke Treatment (TOAST) criteria (8), Benbir and colleagues assessed the association between seizures and different subtypes of ischemic stroke. The authors determined that strokes of atherothrombotic and cardioembolic origin were responsible for up to 36% and 28% of the cases with poststroke epilepsy. Surprisingly, there were patients with lacunar infarcts who developed epilepsy; this finding most likely is related to concomitant cortical involvement. When the group was evaluated as whole, atherothrombotic poststroke epilepsy encompassed 2.7% of all patients with atherothrombotic origin, cardioembolic poststroke epilepsy comprised 2.6% of all patients with cardioembolic origin, and poststroke epilepsy following lacunar infarctions constituted 1.3% of all lacunar strokes.

An interesting finding reported by Benbir and colleagues is that the location of the affected arterial territory is a prognostic factor for the development of epilepsy. The middle cerebral artery territory was the most common location to be associated with seizures—a finding that is similar to what occurs in patients with hemorrhagic strokes. This outcome was expected since the middle cerebral artery covers a great amount of cerebral cortex and is a common arterial distribution for major strokes, although one study found that atherothrombotic infarction of occipital topography has been shown to be an independent predictor of early seizures (9).

Status epilepticus was not found in any of the 1,327 patients included in the Benbir et al. study. This finding contradicts the 9% rate for status epilepticus previously reported among 180 patients with poststroke, first-time seizures (10). However, as Benbir et al. indicate, the overall percentage of poststroke patients who typically develop status epilepticus is so low (i.e., 0.14–0.8%) that it could be assumed that after a stroke, status epilepticus is less common than in other disorders (e.g., tumor, hypoxia).

The study by Benbir et al. has merits; the inclusion of patients with cerebral venous thrombosis is perhaps the most valuable of all. The finding that poststroke epilepsy is common with venous infarctions is novel. However, there still is a need for future studies on stroke and epilepsy to focus on prognosis and long-term outcome. Such findings would provide a clearer picture of the key epidemiological issues and risk factors in the development of seizures and epilepsy in poststroke patients, and, thus, would help to better identify the population at greatest risk for developing poststroke epilepsy.

by Jorge G. Burneo, MD, MSPH

References

1. Lossius MI, Ronning OM, Slapo GD, Mowinckel P, Gjerstad L. Poststroke epilepsy: occurrence and predictors—a long-term prospective controlled study (Akershus Stroke Study). *Epilepsia* 2005;46:1246–1251.
2. Lossius MI, Ronning OM, Mowinckel P, Gjerstad L. Incidence and predictors for post-stroke epilepsy. A prospective controlled trial. The Akershus stroke study. *Eur J Neurol* 2002;9:365–368.
3. So EL, Annegers JF, Hauser WA, O'Brien PC, Whisnant JP. Population-based study of seizure disorders after cerebral infarction. *Neurology* 1996;46:350–355.
4. Bladin CF, Alexandrov AV, Bellavance A, Bornstein N, Chambers B, Cote R, Lebrun L, Pirisi A, Norris JW. Seizures after stroke: a prospective multicenter study. *Arch Neurol* 2000;57:1617–1622.
5. De Reuck J, Claeys I, Martens S, Vanwalleghem P, Van Maele G, Phlypo R, Hallez H. Computed tomographic changes of the brain and clinical outcome of patients with seizures and epilepsy after an ischaemic hemispheric stroke. *Eur J Neurol* 2006;13:402–407.
6. Guidelines for epidemiologic studies on epilepsy. Commission

- on Epidemiology and Prognosis, International League Against Epilepsy. *Epilepsia* 1993;34:592–596.
7. Bousser MG. Cerebral Venous Thrombosis. In: Welch KMA, Caplan LR, Reis DJ, Siesjo BK, Weir B, eds. *Primer on cerebrovascular diseases*. San Diego: Academic Press, 1997, 385–389.
 8. Adams HP Jr., Bendixen BH, Kappelle LJ, Biller J, Love BB, Gordon DL, Marsh EE 3rd. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of Org 10172 in Acute Stroke Treatment. *Stroke* 1993;24:35–41.
 9. Arboix A, Comes E, Garcia-Eroles L, Massons JB, Oliveres M, Balcells M. Prognostic value of very early seizures for in-hospital mortality in atherothrombotic infarction. *Eur Neurol* 2003;50: 78–84.
 10. Velioglu SK, Ozmenoglu M, Boz C, Alioglu Z. Status epilepticus after stroke. *Stroke* 2001;32:1169–1172.