Review

T-Cell-Mediated Autoimmunity

Novel Techniques to Characterize Autoreactive T-Cell Receptors

Klaus Dornmair,^{*†} Norbert Goebels,[†]
Hans-Ulrich Weltzien,[‡] Hartmut Wekerle,^{*} and
Reinhard Hohlfeld^{*†}

From the Max-Planck Institute of Neurobiology,^{} Martinsried; the Institute for Clinical Neuroimmunology,[†] Klinikum Großhadern, Ludwig Maximilians University, Munich; Max-Planck-Institute for Immunobiology,[‡] Freiburg, Germany*

Histological samples of autopsy or biopsy tissue provide the best available evidence that autoreactive T cells are involved in the immunopathogenesis of many autoimmune diseases. However, morphology alone does not provide information on the antigen-specific T-cell receptor (TCR) of these cells, let alone on their antigen specificity. In this review article we discuss a number of emerging possibilities for identifying TCR sequences directly from biopsy tissue. We also review the methods for expressing presumably autoreactive TCR molecules and speculate on how the expressed TCR might be used to identify target antigens. Such information should eventually provide new insights into disease pathogenesis which lead to better therapies. (Am J Pathol 2003, 163:1215–1226)

Autoreactive T lymphocytes are key players in autoimmune diseases. They can act both as regulatory and effector cells. Various animal models have been used to show that the transfer of autoreactive T cells is sufficient to induce a model of an autoimmune disease. Thus, the pathogenic importance of autoreactive T cells can be formally demonstrated (reviewed by¹). Obviously, such transfer experiments cannot be done in human autoimmune diseases. Here the evidence for the pathogenic relevance of T cells is usually indirect. For example, cytotoxic T cells have been shown to surround and strikingly invade muscle fibers in inflammatory muscle lesions of patients with polymyositis (Figure 1a). Then CD8⁺ T cells contact muscle fibers and orient their perforin-containing lytic granules vectorially to-

ward the membrane of the target muscle fiber (Figure 1b). This “in flagranti” situation, morphologically speaking, strongly suggests that by recognizing an antigen on the target muscle fiber the T cell is activated to behave as a cytotoxic effector cell. A careful analysis of their antigen receptor repertoire provides further evidence for the pathogenic role of such T cells. To this end, a number of new techniques have recently become available, eg, microdissection of single infiltrating T cells, combined with single-cell polymerase chain reaction (PCR) amplification of their clonotypic, rearranged T-cell receptor (TCR) V β chain.^{2,3}

These advanced techniques allow us to address the following key questions. Are the tissue-infiltrating T cells mono-, oligo-, or polyclonal, ie, are particular TCR α - or β -chains over-represented in the T-cell receptor repertoire? Which antigen(s) do the infiltrating T cells recognize? How is antigen-recognition related to the pathogenesis and to the course of the disease? Can the recognition process be manipulated to influence disease progression? In this article we review these state-of-the-art technologies for identifying and characterizing such infiltrating T cells and their antigen-specific receptors in tissue biopsy specimens.

Figure 2 shows the overall strategy: to morphologically identify putatively pathogenic T cells in the target tissue, to molecularly clone and express both chains of their antigen-specific TCR, and then to try to identify the cognate antigen(s). None of these steps is trivial, mainly because of the high variability of TCR molecules. They are membrane-anchored $\alpha\beta$ - or $\gamma\delta$ -heterodimeric proteins of antibody-like structure. Each chain is composed of variable (V), joining (J), and constant (C) regions. The V and J regions are connected by the small diversity (D) region and/or random nucleotides (N). The genomic organization and the respective numbers of genetic ele-

Supported by Hermann und Lilly Schilling Stiftung and Deutsche Forschungsgemeinschaft grants SFB 571-A1 and 571-A3.

Accepted for publication May 28, 2003.

Address reprint requests to Klaus Dornmair, Ph.D., Max-Planck-Institute of Neurobiology, D-82152 Martinsried, Germany. E-mail: dornmair@neuro.mpg.de.

Figure 1. A: Immunohistochemistry of a muscle biopsy sample from a polymyositis patient with a monoclonal expansion of $\gamma\delta$ -T cells (taken from¹³). The T-cell marker CD3 is stained red with rhodamine fluorescence; the V δ 2 TCR-chain is stained green with FITC, and the double-positive $\gamma\delta$ -T cells are stained yellow. The big brown cells are muscle fibers. The autoaggressive $\gamma\delta$ -T cells surround and invade the muscle fibers, whereas the CD3⁺V δ 2⁻ bystander T cells are not in contact with muscle fibers. Scale bar, 50 μ m. **B:** Autoaggressive T cells excrete the cytotoxic protein perforin into the attacked muscle fiber (m) as shown by confocal laser microscopy of a polymyositis muscle biopsy specimen (taken from¹⁹). Perforin is stained red with rhodamine. The **arrow** indicates the vectorial perforin excretion. Scale bar, 10 μ m.

ments of the TCR V and J regions are delineated in Figure 3a and Table 1. Each chain of the resulting receptor protein has one hypervariable and two variable loops that protrude from a compact β -sheet structure.^{6–8} These loops are termed “complementarity determining regions” (CDR), because they directly contact the antigenic peptide. The high degree of variability (the total diversity of $\alpha\beta$ TCRs may exceed $10^{17.5}$) increases the difficulties to determine the TCR sequences of the infiltrating cells, because the variety of very similar, but not identical proteins and nucleotide sequences is immense.

Identification of T Cells and their Receptors in Human Biopsy Tissue

Availability and Choice of Tissue

The availability of biopsy tissue depends on the disease. In multiple sclerosis (MS), for example, a biopsy is made

only in exceptional cases. Relatively few studies have directly addressed the T-cell repertoire expressed in MS lesions.^{9,10} Many investigations require fresh frozen rather than fixed tissue. In a recent study, the TCR β -chain sequences of single tissue-infiltrating T cells were identified in frozen tissue of two MS patients, using a combination of microdissection and single-cell PCR.²

In contrast, tissue biopsy is routinely taken for diagnostic reasons in other diseases, eg, polymyositis. Polymyositis has the additional advantage that muscle is easily accessible, in contrast to CNS. This is probably one of the reasons why there have been numerous studies of the T-cell repertoire expressed in polymyositic muscle.^{11,12} A monoclonal expansion of $\gamma\delta$ T cells was found in a unique case of polymyositis.^{13,14} In other cases, striking expansions of muscle-infiltrating $\alpha\beta$ TCR+CD8⁺ T cells were observed.^{15–18} The infiltrating T cells made direct contact with the muscle fiber targets and secreted lytic granules toward the attacked muscle fibers,¹⁹ suggesting that the T cells recognize an antigen on the muscle fiber surface. These examples illustrate that the target tissue itself, rather than peripheral blood or cerebrospinal fluid, is the best cell source for this type of study.

Morphological Identification of Autoaggressive T Cells

The quality of the biopsy material is crucial for all types of investigations. Consequently, the appropriate preservation of the tissue is an essential prerequisite. Ideally the tissue should be frozen immediately after excision to prevent any degradation of RNA, DNA, or proteins. Most traditional fixation procedures, however, modify proteins and nucleic acids, thus hindering any further analysis.

One of the first steps in identifying T cells and T-cell receptors in tissue is to apply immunohistochemistry, using monoclonal antibodies against differentiation markers like CD3, CD4, CD8, and antibodies against TCR V- α and V- β family members. Staining with the latter antibodies may provide the first indications of (oligo)clonal expansions, but this method has obvious limitations. First, although the panel of available anti-V- β monoclonal antibodies is quite large, currently there are only three antibodies against TCR V- α , namely anti-V α 2,²⁰ anti-V α 12,²¹ and V α 24.²² Note that V α 24 is not only expressed on “regular” CD4⁺ and CD8⁺ T cells, but also on human NKT cells (reviewed by^{23,24}). Second, not all of the available anti-TCR V-chain family are suited for immunocytochemistry. Third, most antibodies specific for a particular V family do not distinguish between the different subfamilies, which may differ by some amino acids, whereas other antibodies may recognize only one particular allele or subfamily, but not the whole family. Fourth, V-region specific antibodies do not distinguish between different clonotypes (that is, different hypervariable CDR3 regions). These limitations indicate that immunohistochemical methods may provide the first clues of clonal expansions, and may help to identify the expanded TCR

Figure 2. Strategies for identifying unknown T-cell antigens. T cells identified in biopsy samples are excised and their TCR DNA is amplified by single-cell PCR. In case of a monoclonal expansion, a larger piece of tissue containing many cells may be analyzed *in toto*. The DNA fragments obtained by PCR must then be reconstructed to full-length chains. The different genetic elements are displayed in different colors. α chain: V region, green; J region, red; C region, yellow. β chain: V region, turquoise; J region, magenta; C region, gray. They are then inserted into expression vectors, and transfected into appropriate recipient cells. TCR molecules may be expressed 1) as soluble proteins that can be used for affinity chromatography or for screening cDNA-expression libraries, or 2) as a component of the TCR/CD3 signaling complex on the cell surface of lymphoid cell lines. The same colors as above are used for the TCR-protein domains. Protein tags, the accessory CD3 protein-complex, and the cell membrane are given in black. The transfectants may be used to test potential target antigens directly or to screen peptide or cDNA expression libraries.

V-chain family, but they are not suitable for identifying TCR clonotypes.

If expanded T-cell clones are observed in tissue samples, they are usually superimposed on a background of unrelated cells. These may include several

subpopulations: 1) bystander T cells attracted non-specifically by inflammatory cytokines or chemokines, 2) intravascular T cells from blood capillaries, 3) irrelevant clonal expansions as they may occur in healthy, especially older persons,^{25–28} and 4) expanded T-cell

Figure 3. Genetic organization of TCR chains and methods for amplifying functionally rearranged gene products. TCR molecules consist of an α and β , or γ and δ chain. **a:** Organization of TCR genes (see reference⁴ for details). Each of the chains is composed of V and J elements that are connected by hypervariable regions that may contain D elements and random nucleotides (N). The colors used here correspond to the colors used in Figure 2 for the TCR α -chain. The n(D)n region between the V and J regions is given in dotted pink. D genes exist only for the TCR β - and δ -chains; they are omitted in the scheme. The numbers of functional genes are given in Table 1. During T-cell development, germline genomic DNA of the V, (D), and J genes is rearranged. mRNA is generated by splicing and transcription of V(D)J-rearranged genomic DNA. Introns are indicated as dashed lines. The approximate positions of the CDRs are indicated. **b:** Schematic depiction of PCR-based methods to amplify TCR cDNA. Shown are anchored, inverse, and family PCR and CDR3 spectratyping. Forward primers are shown in blue, reverse primers in brown. See text for explanations. **c:** Outline of single-cell PCR. A stained cell in tissue is excised by a laser beam and catapulted into a reaction vessel (red arrow). The photo is taken from reference³. Rearranged genomic DNA or cDNA of this cell is then amplified using pools of PCR primers. **d:** Reconstruction of full-length TCR cDNA from smaller PCR fragments: leader plus beginning of V region (a), fragment of V(D)J amplification by single-cell PCR (b), semisynthetic linker to insert end of J plus beginning of C region (c), if required, and remainder of C region (d). Depending on the size and content of fragment (b), this strategy may be modified.

clones in patients with superimposed infections. The first two groups of bystander cells and T cells from capillaries typically have random distributions of T-cell receptors and therefore provide a continuous background, whereas the latter two groups may include

Table 1. Numbers of Functional Human TCR V-, D-, J-, and C- Genes (Data Taken from Reference 4)

TCR chain	V	D	J	C
α	46	0	50	1
β	48	2	13	2
γ	6	0	5	2
δ	11	3	4	1

The combinatorial diversities of gene elements are thus 2.8×10^6 for $\alpha\beta$ TCR or 3,000 for $\gamma\delta$ -TCR molecules. These numbers are drastically increased by the additional "junctional diversities" of up to 2×10^{11} (reference 5) introduced into the CDR3s by random insertion or deletion of the N nucleotides between the gene elements. This results in theoretical diversities in the range above 10^{17} or 10^{14} , respectively.

expanded clones with preferred TCR expression. They thus may mimic the presence of expanded autoimmune T cells. Further confusion may arise from the fact that in infectious disease, the clonally expanded T cells may be activated, as are some of the autoreactive T cells in autoimmune disease. It is therefore important to distinguish the truly autoaggressive T cells from the different types of expanded, activated, and bystander cells. This is not usually achieved by staining TCR with soluble MHC/peptide complexes, because this would require previous knowledge of the relevant antigens.²⁹ Important distinguishing criteria can be provided by the morphology and location of the inflammatory cells in the tissue. In polymyositis, for example, autoaggressive CD8⁺ T cells make direct contact and invade the target muscle fibers.³⁰ In this case it is relatively simple to identify the potentially autoaggressive T cells and distinguish them from unrelated bystander cells. In other cases, however, the topographical situation may

Table 2. Methods to Analyze Clonal Expansions in Tissue Biopsies

Method	Advantage	Disadvantage
Immunohistochemistry	Covers many V β genes Fast and reliable Discrimination of bystander cells	Only three V α Genes CDR3 and is not detected Antibody crossreactivity Bystanders are co-amplified Statistically inefficient
Anchor PCR	All V α and V β genes Unbiased by V-family Yields full-length sequences	
Inverse PCR	All V α and V β genes Unbiased by V-family Yields full-length sequences	Relatively insensitive Bystanders are co-amplified Statistically inefficient Different sensitivity for different Vs May introduce bias
Family PCR	All V α and V β genes	Bystanders are co-amplified V α analysis not yet described Yields short fragments
CDR3 spectratyping	All V β genes May detect all CDR3 nucleotides Unbiased by V-family	
Single-cell PCR	Focus to interesting cells Combination with immunohistochemistry Unbiased by V-family No statistics required	V α should be surmised Low efficiency Costly, laborious, and slow

For immunohistochemistry fixed or frozen biopsy samples may be used. For all other methods only frozen biopsies are suitable.

be less informative. For example, CD4⁺ T cells may act as “helper” cells by secreting soluble mediators in the local milieu of the inflammatory lesion. This function does not necessarily depend on direct cell-cell contact. Further, the antigen-specific priming of CD4⁺ T cells may occur at other sites outside the lesion.

Characterization of Expanded TCR Chains from Tissue Samples

Several PCR-based techniques can be used to determine the DNA sequences of expanded T-cell clones in homogenized tissue samples. The basic principles of anchored, inverse, and family PCR, and of TCR-CDR3 spectratyping are outlined in Figure 3b. Some advantages and limitations of the different techniques are compared in Table 2. All techniques use cDNA as templates. Anchored and inverse PCR use PCR primers that hybridize to shared TCR chain sequences, irrespective of the V or J family usage. They therefore allow an essentially unbiased expansion of all TCR V-chain families. For anchored PCR (also termed rapid amplification of cDNA ends),^{31,32} a synthetic oligo-C sequence is first added to the N terminus of the cDNA. Then an oligo-G-containing forward-PCR primer is used for amplification. The corresponding reverse-primer hybridizes to the C region. For inverse PCR,³³ the double-stranded cDNA is converted into a closed circle, and the unknown regions are amplified using a primer pair within the C region. The third method, TCR V-family PCR,³⁴ uses a pool of many different primers that hybridize to the V regions. Therefore, a specific primer must be included for each V family. Only the reverse primer (which hybridizes to the C region) is the same for all reactions. If the chosen experimental conditions are optimal, this method can amplify minute amounts of cDNA from particular TCR sequences present in the sample. However, one disadvantage of this method is that not all V sequences may be amplified equally well, depending on the different efficiencies of the

various primers. It therefore might be problematic to directly compare efficiencies between different V families, unless the primer efficiencies have been quantified in earlier experiments and are normalized. It is essential with all techniques, whether anchor, inverse, or family PCR, to clone the PCR products and to sequence a statistically significant number of clones.

A further development of family PCR is the CDR3-spectratyping technique (also termed Immunoscope) (reviewed by³⁵). First, the β -chain cDNA is amplified by 25 different V-specific forward and one C-specific reverse primer. Then the products of these reactions are split, 13 different fluorescence-labeled J-specific primers are added to each aliquot for further “run-off” reactions, and finally the length distributions of the products are analyzed by high-resolution acrylamide gel electrophoresis. If the T-cell populations are polyclonal, the PCR products have Gaussian-like length distributions. In contrast, if the T cells are (oligo)clonal, individual peaks corresponding to PCR products of one defined length appear, skewing the Gaussian distribution. Sequence analysis of prominent peaks may indicate expansions of particular clones. Thus, CDR3 spectratyping provides a broad survey of possible clonal expansions of many TCR chains and may identify clonal expansions irrespective of the hybridization efficiencies of different V primers. A major disadvantage of CDR3 spectratyping, as compared to anchored and inverse PCR, is that the method has not yet been established for TCR α -chains, due to the considerably large number of J- α genes.

In summary, immunocytochemistry is indispensable for identifying potentially autoaggressive T cells in the target tissue, in particular when cytotoxic effector T cells are analyzed. Despite the limitations of anti-V antibody staining, it still gives a quick and broad overview by assigning V-chain family usage to morphologically identifiable cells. This aspect is particularly important because it helps decide whether *in situ* expanded T-cell clones are indeed attacking the target tissue, or whether

they are bystanders. To identify the complete TCR sequences of the expanded clones, one of the PCR-based techniques must be used. All methods are highly sensitive, particularly if nested primer pairs are used. Anchored and inverse PCR have the advantage of yielding full-length products and virtually unbiased product distributions, but inverse PCR may require relatively high amounts of template. Moreover, the products have to be cloned into plasmids, and a statistically significant number of individual *Escherichia coli* clones have to be sequenced. However, the products of TCR family PCR should also be cloned before sequencing, because direct sequencing of the PCR products might reveal only the most prominent CDR3 sequence. The most straightforward indication for clonal expansion in tissue can be obtained by CDR3 spectratyping, because this technique may directly detect clonal expansions not only of particular V, but also of the corresponding CDR3 regions. For final proof of clonal expansion, the products must also be sequenced. If all primers to all V and J families are included, a comprehensive overview may be obtained. However, as mentioned above, the broad variability of J- α elements still impedes investigation of the TCR α -chains and hence a complete analysis.

Identification of Paired TCR Chains Expressed in Individual T Cells

In most studies reported so far, the T-cell expansions observed in autoimmune tissues were oligoclonal. Only a single case of a *monoclonal* expansion has been reported.¹⁴ This was a very special case, because the expanded T cells expressed a $\gamma\delta$ rather than $\alpha\beta$ TCR. Hence, analysis of a tissue homogenate containing oligoclonally expanded T cells with one of the PCR techniques described above would yield several α - and β -chain sequences, but it would be impossible to know which α chain pairs to which β chain. As long as this pairing is not known, however, the structure of the functional pathogenic receptor will not be known. In principle, single-cell PCR can overcome this fundamental problem. For this approach, interesting cells are identified in the biopsy sample by staining with appropriate antibodies. Next, single T cells of interest are excised directly from tissue sections under a microscope using a micromanipulator or a highly focused laser beam (laser-assisted microdissection),^{36,37} and the isolated cells are transferred into a PCR tube where the TCR genes are amplified by single-cell PCR (Figure 3c).

If the α - and β -chain sequences of the expressed TCR are not known, pools of many different V- and J-specific PCR primers, or degenerate primers, must be used. Both types of primers are of course less efficient than defined pairs of specific primers. Nevertheless, unbiased amplification of the TCR β -chain genes from single cells has been successful.^{2,3,38–41} Because RNA might be unstable in a biopsy sample, the authors used rearranged genomic DNA as a template; this is much more resistant to degradation. Obvious disadvantages are that only a single copy of template is present per cell, and the conserved region is sep-

arated from the rearranged V(D)J region by a large intron (Figure 3a). Therefore, it was not feasible to use a single reverse-PCR primer (hybridizing to the C region), as would be used for RT-PCR. Instead, the authors used one pool of 7 reverse J primers covering all 13 different J- β genes (Table 1) together with 8 forward V-primer pools, each containing 2 to 5 individual primers for hybridizing to the V- β regions. With this approach, unknown TCR β -chains could be amplified from single cells excised from tissue sections. In up to 30% of all single cells examined a rearranged TCR β -chain was detected.

Amplification of the TCR α -chains is even more challenging. Although the number of different V α elements is comparable to that of the β chain, the α -chain repertoire may select from a greater number of different J elements (50 functional J α genes as compared to 13 for J β ; Table 1). Thus, the number of primers has to be significantly increased if all possible V and J elements are to be covered. One way to restrict this variability is to stain the putative target T cells with one of the available three anti-V α antibodies. After microdissection of the corresponding cell, the V α chain can then be amplified using only one specific V α primer instead of a pool covering all possible V α sequences. This approach was successfully used to isolate a pair of α and β chains from a freshly prepared, FACS-sorted single cell.⁴² These authors used cDNA as the PCR template, which allowed them to ignore any variability of the J regions and simply employ reverse primers that hybridized to the C region. However, this approach might fail when frozen sections of biopsy samples (rather than fresh cells) are investigated, because of the instability of mRNA as compared to that of DNA.

To define pairs of TCR α - and β -chains we recently stained TCR in biopsy sections of a polymyositis patient with an anti-V α 12 antibody, then isolated stained cells by laser microdissection, and amplified the TCR α - and β -chains from these V α 12+ single cells (C.K. Schneider et al, unpublished results). We successfully amplified pairs of α and β chains, thus demonstrating the usefulness of this approach in case the α chain can be stained. An additional advantage of V α staining is that it shows that the particular V α chain is expressed on the cell surface. This may be of interest, because two α chains (or rarely, two β chains) are synthesized in a significant number of T cells,^{22,43–45} and it is not known which chain is expressed on the cell surface.

"Reconstruction" of TCR cDNA

Once a pair of co-expressed α - and β -TCR chains is identified, a functional heterodimer has to be reconstructed. Only amplification by anchored and inverse PCR yields products that cover the full-length sequences of TCR-DNA. The primer pairs used for family PCR, CDR3 spectratyping, or single-cell PCR mostly yield relatively short fragments because they are designed for high specificity and sensitivity. Therefore, they hybridize to sequences somewhere in the middle of the V regions, in the J regions, or at the beginning of the C regions. Parts of the V and C regions will be missing in such PCR

products, and some products of single cell PCR reactions may not even contain complete J sequences. To express the TCR proteins, it is therefore necessary to reconstruct the complete sequences from fragments. Such a reconstruction from single-cell PCR is outlined schematically in Figure 3d. A typical single-cell PCR fragment ranging from the middle of the V to the middle of the J regions (designated "b") is ligated to a fragment covering the remainder of the J and the beginning of the C region ("c"), and inserted into a cloning plasmid that already contains the beginning of the V regions ("a") and the rest of the C regions ("d"). Fragments "a" and "d" may be obtained from the cDNA of a unrelated TCR with identical V and C regions. Alternatively, the hypervariable regions of an unrelated TCR with identical V and C regions may be replaced by site-directed mutagenesis. The precise strategy depends on the number and lengths of the fragments, restriction sites, and other features.

Expression of Recombinant TCR Molecules

There are basically three strategies for producing the reconstructed autoreactive TCR molecules in a recombinant expression system. First, they can be designed as soluble proteins, ie, as small monomeric (glyco-)proteins, analogous to Fv- or Fab-antibody fragments. These monomeric TCR proteins may carry tags that can be used for detection, oligomerization, or coupling to solid matrices. Second, the TCR can be expressed at the surface of immortalized T cell lines. Here the recombinant TCR is integrated into the CD3 complex of the recipient cells and coupled to their signaling machinery. Third, the TCR can be expressed *in vivo* in a transgenic animal model.

Soluble TCR Molecules

The most straightforward approach is to express the TCR of interest as a soluble molecule. Different $\alpha\beta$ - or $\gamma\delta$ -TCR molecules of murine or human origin have been generated using a broad spectrum of expression protocols. Eukaryotic expression systems often yield correctly folded and processed proteins, but their amount is relatively small. For example, TCR molecules were expressed in myeloma, COS, *Drosophila* cells, and *Baculovirus*-transfected insect cells.^{46–51} In the oxidizing milieu of the periplasm of *E. coli*, folding and formation of disulfide bonds may also be achieved, but again with comparably low yields.⁵² In contrast, expression in the cytosol of bacteria may yield much larger amounts of protein, but the TCR chains have to be refolded *in vitro* from insoluble inclusion bodies.^{53–57} Considerably different constructs have been used. The TCR chains were either co-expressed to directly form heterodimers, expressed as single-chain constructs with the variable domains linked covalently by peptide linkers, or they were expressed separately and later folded and assembled *in vitro*. In all constructs the membrane-spanning regions were truncated and replaced by cleavable lipid linkers⁴⁶ or peptide tags, such as oligo-histidine residues, which facilitate purification or detection.^{51,56} To support dimer-

ization of the chains leucine-zipper domains were introduced,⁵⁰ or the TCRs were fused to other large proteins, eg, antibody domains^{47,48} or the CD3 ζ chain.⁴⁹ These different expression strategies have been used successfully to produce soluble TCR molecules and even protein crystals, allowing structural analysis of $\alpha\beta$ TCRa (reviewed by^{6–8}).

In principle, soluble TCR molecules could be used to identify the corresponding antigen. The main obstacle to this approach, however, is the low affinity of TCR-ligand interactions. Many studies have addressed the kinetics and thermodynamics of TCR-ligand binding, using soluble TCR molecules or MHC/peptide complexes (reviewed by^{7,58}). The results show that the interactions of the monomeric compounds are generally of low affinity and that antigen presentation is dominated by the high avidity gained by oligomerization and by the kinetics of the interaction. Therefore, the classic methods for detecting and analyzing molecular interactions such as affinity chromatography or library screening cannot be applied. These methods require high binding affinities, because they imply washing steps during which the TCR-ligand complexes would dissociate and be lost.

TCR Expression on the Surface of Lymphoid Cell Lines

Expression of a TCR on the surface of a living cell has several advantages to soluble TCR. First, the TCR molecules are oligomerized on a cell surface and may interact with antigens that are oligomerized themselves, for example, bound to a microtiter well or to the surface of an antigen-presenting cell. Therefore, the low affinity of the TCR may be overcome by the high avidity of the oligomerized molecules. Second, immortalized cells can be grown essentially without limitations. Third, and most importantly, TCR activation triggers the production of cytokines, which may be measured by ELISA. Each of these steps amplifies the initial signals. Therefore the overall sensitivity is drastically increased.

The recombinant expression of TCR molecules on the surface of immortalized cells was first described for a TCR β -chain in a recipient cell that lacked the endogenous β -chain.⁵⁹ The recombinant β -chain and the endogenous α -chain were co-expressed as heterodimers. In later experiments two recombinant α - and β -chains were expressed so that the recipient cells expressed both the recombinant and the endogenous receptors.^{60,61} Besides other cell lines, such as TG40⁶² or DS23,⁶³ the commonly used recipient cell lines are the human T-lymphoma Jurkat and its TCR α - or β -chain deletion mutants 18.B3,⁶⁴ J79,⁶⁵ RT3-T5.3,⁵⁹ JBN,⁶¹ 31-13,⁶⁶ and the murine T-hybridoma line 58 $\alpha^- \beta^-$, which lacks endogenous TCR chains.⁶⁷ 58 $\alpha^- \beta^-$ is derived from DO-11.10, which has been generated by fusing chicken-ovalbumin specific T-cell blasts from BALB/C mice to the AKR thymoma BW5147.⁶⁸ 58 $\alpha^- \beta^-$ was later transfected with human CD4 to facilitate the recognition of human HLA class II molecules, and with the mouse CD3 ζ chain to improve signal transduction.⁶⁹ To improve the recognition of murine class-I MHC molecules,

58 $\alpha^- \beta^-$ was also co-transfected with murine CD8 α - and β -chains.⁷⁰ Different plasmids and retroviruses were successfully used to transfect immortalized cell lines like Jurkat and 58 $\alpha^- \beta^-$. Recently, TCR molecules were expressed in spleenocytes, peripheral blood lymphocytes, and T-cell lines (reviewed by⁷¹).

TCR expression can be monitored by FACS analysis using V-region specific antibodies. If appropriate antibodies are unavailable, anti-CD3 antibodies may be used, provided that the recipient mutant cell line lacks the endogenous pendant of the TCR chain to be transfected. This indirect detection strategy exploits the fact that TCR-CD3 complexes can only reach the cell surface if all components are expressed concomitantly. Recognition of the TCR antigen is typically assessed by measuring secreted IL-2. Although this assay might not be as sensitive as a cytotoxicity assay, Jurkat and 58 $\alpha^- \beta^-$ cells grow fast and may be expanded to almost unlimited cell numbers. This is a great advantage to using "real" T cells that are notoriously difficult to grow and can rarely be expanded to more than a few million cells. Further, transfectants can be easily cloned, again in contrast to "real" T cells. It is always advisable to pick single clones during the selection of stable transfectants, because the number of transfected plasmids and the position of vector insertion into the genome of the recipient cell may vary considerably. Analysis of individual clones allows the selection of those with the highest TCR expression levels. Moreover, site-directed mutations may be easily introduced into transfectants by standard techniques. Mutations in the CDR regions help exclude superantigen-like or non-specific activation and thus prove specificity of antigen recognition.⁷²

Many investigators have chosen Jurkat or its derivatives as recipient cells for transfection of human TCR chains, because Jurkat is of human origin and therefore all molecules of the CD3 complex fit to the transfected TCR chains. Further, Jurkat is one of the best-characterized mammalian cell lines. One disadvantage is that there are no sublines lacking both TCR chains. Therefore, transfected Jurkat cells may express not only their own endogenous TCR and the transfected TCR heterodimers, but hybrids of transfected and endogenous chains might also reach the surface. Such transfectants are hardly suited for investigating unknown antigens, because it would be difficult to distinguish the signals of the transfected, parental, and hybrid TCR molecules. In contrast, 58 $\alpha^- \beta^-$ cells and their derivatives lack both endogenous TCR chains⁶⁷ so that no hybrid receptors are formed. Furthermore, even if there were no antibodies to any of the transfected TCR chains available, surface expression might be detected by using anti-CD3 antibodies. It might be a disadvantage that 58 $\alpha^- \beta^-$ is a murine cell line, but it has been shown that human-mouse chimeric TCR molecules consisting of human Vn(D)nJ regions and murine C regions^{69,73-75} as well as full-length human $\alpha\beta$ - or $\gamma\delta$ -TCR molecules may be expressed in the context of murine CD3 molecules.^{62,72} The specificity of the human TCR is preserved in the context of murine CD3.

Recombinant TCR Molecules in Transgenic Animal Models

An obvious way to validate the destructive potential of a presumably autoaggressive human TCR *in vivo* is to use transgenic techniques to insert the TCR into living animals. However, if the target antigen is not known, it is also not known whether the transgenic mice express it. Several TCR transgenic mice that are useful models for human autoimmune diseases have been described so far (reviewed by⁷⁶⁻⁷⁹). In these cases, the antigen(s) recognized by the transgenic TCR were known in advance. Many of these animals expressed TCR molecules of murine origin, but mice with appropriate genetic backgrounds for investigating whether symptoms similar to the human disease might be induced also expressed human autoreactive TCR molecules. These animals expressed the antigen either endogenously, were crossed with mice transgenic for the antigen, or were challenged with the antigen. So far no animal model has been described in which a pathogenic TCR was first detected in a patient and later expressed in a transgenic animal without previous knowledge of the target antigen. TCR-transgenic animals are more useful for studying the pathogenesis and therapy of autoimmune diseases than for identifying unknown antigens.

Using TCRs to Identify Unknown Target Antigens

Both $\alpha\beta$ - and $\gamma\delta$ -T cells have been observed in tissue lesions of patients with autoimmune diseases. The mechanism of antigen recognition of these two types of T cells is quite different. CD8⁺ $\alpha\beta$ T cells recognize class I MHC molecules complexed with peptides of 8 to 10 amino acids in length, and CD4⁺ $\alpha\beta$ T cells recognize class II MHCs with much longer peptides (reviewed in^{7,8}). In contrast, $\gamma\delta$ TCRs have recognition patterns similar to antibodies; they may recognize a range of epitopes, including conformational epitopes of proteins, or carbohydrates, lipids or low molecular weight compounds alone or bound to MHC or MHC-like proteins such as CD1. This recognition is independent of the CD4 or CD8 coreceptors (reviewed in⁸⁰⁻⁸²). *In vivo*, additional costimulatory receptors and accessory molecules are involved.⁸³ The methods for identifying $\alpha\beta$ - and $\gamma\delta$ -T-cell antigens will be discussed separately.

$\alpha\beta$ -TCR Molecules

To identify the antigens of $\alpha\beta$ -TCR molecules, two different components, the antigenic peptide and the HLA-molecule, must be determined. Each individual expresses several relevant polymorphic HLA molecules; it is not known initially which HLA allele presents the relevant antigenic peptide. Immunohistochemistry may reveal if the putatively pathogenic T cells in the biopsy tissue express CD4 or CD8 molecules, and may thus point to MHC class II or class I restriction. The restriction

molecule may be further narrowed down by using genetic and epidemiological data, which may indicate an association of the autoimmune disease with a certain HLA type. Although none of these links are strict, they may provide a clue. If the HLA restriction molecule is not known, autologous EBV-transformed B cells, isolated from the same donor as the TCR, are probably the best source of antigen-presenting cells, because they carry all of the relevant HLA molecules of the patient. To formally determine the specific HLA restriction molecule, human typing cell lines or HLA-transfected mouse fibroblasts may be used as antigen-presenting cells. It should be noted that TCR-transfected human cells (like Jurkat) and mouse cells (like 58 $\alpha^- \beta^-$ cells) may be activated specifically by human and mouse antigen-presenting cells, provided that the correct MHC/peptide complex is expressed. However, we found that higher IL-2 levels may be obtained if 58 $\alpha^- \beta^-$ cells are stimulated by HLA-transfected mouse fibroblasts, presumably because the mouse fibroblasts express the appropriate murine costimulation molecules.

The determination of peptide specificities of TCR molecules has a long tradition in molecular immunology. However, most of these experiments were conducted with previously known antigen specificities, ie, by using peptide derivatives of known parent peptides. If an interesting TCR is cloned from a biopsy sample, the situation is different in that the investigator initially does not have any clue about the antigenic peptide. If the HLA restriction is known (which is usually not the case), certain hints may be obtained from the known peptide-binding motifs of the restricting HLA molecule. A list of HLA peptide-binding motifs may be found under <http://syfpeithi.bmi-heidelberg.com>. On the basis of known anchor positions, a tailor-made peptide library may be constructed in which only amino acids suspected to face the TCR are permuted, but the anchor positions are fixed. Another strategy is to use random libraries containing fixed amino acids facing the TCR. This latter approach revealed the recognition motif of a human CD8 $^+$ T-cell line of known fine-specificity.⁸⁴ Different types of peptide libraries have been described, mostly chemically synthesized pools of peptides but also phage libraries (reviewed in⁸⁵⁻⁸⁷). The basic dilemma of all peptide libraries is that the more comprehensive they are (ie, the more peptides they include), the smaller the concentrations of individual peptides, which finally may be too diluted to even be detected.

$\gamma\delta$ -TCR Molecules

$\gamma\delta$ -TCR molecules do not depend on antigens presented by MHC molecules, but they resemble antibodies in their antigen-recognition properties (reviewed in⁸⁴⁻⁸⁶). While the methods for identifying potentially pathogenic $\gamma\delta$ -T cells in tissue and for analyzing their TCR are essentially the same as for $\alpha\beta$ -T cells, the methods for identifying target antigens are different. One possibility is to construct a soluble TCR that might be used for affinity chromatography or for screening protein expression libraries.

However, both methods depend on high affinities of the receptor ligand pairs. Up to now the affinity of only one $\gamma\delta$ -TCR/antigen pair has been measured.⁸⁸ The reported dissociation constant of 0.1 $\mu\text{mol/L}$ corresponds to a relatively high affinity compared to $\alpha\beta$ TCRs, but it is still more than two orders of magnitude smaller than the affinity of a typical antibody-ligand complex. For another $\gamma\delta$ -TCR/antigen pair the affinity was not high enough to be measured.⁸⁹ Alternatively, a $\gamma\delta$ TCR may be expressed on the surface of immortalized T-cell lines, which may be used as a highly sensitive tool to screen candidate antigen preparations, eg, from candidate tissues, cultured cells, or those obtained by protein-chemical methods. For example, we recently investigated the antigen recognition pattern of a $\gamma\delta$ TCR previously identified in muscle tissue of a patient with polymyositis.^{13,14,72} This is one of the few examples, if not the only known case, of a monoclonal T-cell expansion in the target tissue of a patient with autoimmune disease.¹¹ We reconstructed the cDNA of this TCR, and expressed it in 58 $\alpha^- \beta^-$ cells. The transfectants expressed the human $\gamma\delta$ -TCR in the context of the murine CD3 signaling-complex on their cell surface and were activated by appropriate antibodies. Importantly, they were also activated by extracts of cultured human myoblasts and rhabdomyosarcoma cells (Figure 4A). Recognition was mediated by the CDR3 region of the TCR, because control transfectants with defined exchanges in their TCR sequences could be activated by anti- $\gamma\delta$ -TCR antibodies, but they no longer recognized muscle antigen (Figure 4B). The data suggest that this $\gamma\delta$ TCR indeed recognizes an autoantigen expressed in the target tissue. Although the target antigen has not yet been molecularly defined, this example illustrates that the strategies discussed here may help in the search for unknown (auto-) antigens recognized by infiltrating T cells in human target tissues.

Conclusions

Recent technical progress has provided a series of tools that might eventually allow us to identify the target antigens recognized by infiltrating T cells. Potentially pathogenic T cells are first identified in tissue, detected if possible "in flagranti". Next, the genes of their antigen receptors are cloned. The TCR proteins are then expressed in recombinant expression systems, which finally can be used for screening peptide- or protein-expression libraries. The achievement of each of these individual steps signifies a major breakthrough. Here we have described some recent techniques and provided several examples for each of the individual steps of TCR characterization, including identification and cloning of TCR chains of monoclonally expanded infiltrates or of TCR chains from individual, presumably pathogenic cells, recombinant expression of these molecules, and attempts to identify the target antigens of $\alpha\beta$ or $\gamma\delta$ TCRs. However, the successful combination of all of the successive steps, starting from a patient's biopsy and ending with the clear-cut identification of the target antigen(s), has not

Figure 4. A human $\gamma\delta$ -TCR that recognizes a muscle autoantigen. **A:** Subcellular fractions of human myoblasts and TE671 human rhabdomyosarcoma cells stimulate the $\gamma\delta$ -TCR transfectants. Intact myoblasts and TE671 cells did not activate transfectants. Lysed cells or supernatants after lysis and centrifugation induced IL-2 production. This indicates that the antigen is not presented on the cell surface. Strong stimulation was observed with ultracentrifuged supernatants, showing that the antigen is present in the cytosol. BSA (1 mg/ml) served as negative control. **B:** Recognition of the $\gamma\delta$ -TCR antigen is CDR3 specific as shown by TCR mutagenesis. Activation of different TCR transfectants by TE671 ultracentrifuge supernatant as measured by secreted IL-2. The original $\gamma\delta$ -TCR transfectant is compared to an $\alpha\beta$ -TCR transfectant and to a series of transfectants carrying mutated $\gamma\delta$ -TCR molecules with exchanged variable regions (VJ- γ -, VDJ δ -), amino acid mutations in the CDR3 regions only (n γ -, nD δ -, nDJ δ -), and an alternative constant region, but original VDJ regions (C γ -). Only the original $\gamma\delta$ -TCR transfectant and the mutant with altered C region recognized the antigen, whereas the other transfectants carrying mutated TCR molecules were not stimulated. This clearly shows that the antigen is recognized specifically via the CDR3 regions of both chains, and that the TCR is not stimulated by a superantigen (**A** and **B** reprinted from *J. Immunol* 2002, 169:515–521 with permission from American Association of Immunologists, Inc.⁷²).

yet been accomplished. One of the bottlenecks is that T-cell infiltrates are usually not monoclonal but polyclonal. Therefore, elaborate techniques like single-cell PCR are required to detect the corresponding TCR α - and β -chain pairs. Moreover, identification of the TCR α -chain sequence is especially problematic, although recent results indicate that it is in principle possible. This raises the hope that the successful identification of both TCR chains may be reliably achieved in near future. Another problem that needs to be overcome is the low affinity of many TCR molecules to their antigens, but published work has shown at least some receptors might have sufficient affinity to be analyzed *in vitro*. Despite these problems, the identification of

T-cell antigens therefore seems to be almost within our reach. This is a worthwhile goal, for not only will we eventually be able to explain the pathogenesis of autoimmune diseases, but also to develop new diagnostic and therapeutic approaches.

Acknowledgments

We thank Judy Benson for helpful comments on the manuscript.

References

- Wekele H, Kojima K, Lannes-Vieira J, Lassmann H, Linington C: Animal models. *Ann Neurol* 1994, 36:S47–S53
- Babbe H, Roers A, Waisman A, Lassmann H, Goebels N, Hohlfeld R, Friese M, Schröder R, Deckert M, Schmidt S, Ravid R, Rajewsky K: Clonal expansion of CD8⁺ T cells dominate the T-cell infiltrate in active multiple sclerosis lesions shown by micromanipulation and single-cell polymerase chain reaction. *J Exp Med* 2000, 192:393–404
- Hofbauer M, Wiesener S, Babbe H, Roers A, Wekerle H, Dornmair K, Hohlfeld R, Goebels N: Clonal tracking of autoaggressive T cells in polymyositis by combining laser microdissection, single-cell PCR and CDR3-spectratype analysis. *Proc Natl Acad Sci USA* 2003, 100: 4090–4095
- Lefranc MP, Lefranc G: The T-cell receptor. *Facts Book*. London, Academic Press, 2001 p 18ff
- Janeway CA, Travers P, Walport M, Capra, JD: T cell receptor gene rearrangement. *Immunobiology*. 5th ed. London, Elsevier Science Ltd., 2001, pp 137–141
- Allison TJ, Garboczi DN: Structure of $\gamma\delta$ T-cell receptors and their recognition of non-peptide antigens. *Mol Immunol* 2002, 38:1051–1061
- Lefranc MG, Luz JG, Wilson IA: Structural and thermodynamic correlates of T-cell signaling. *Annu Rev Biophys Biomol Struct* 2002, 31:121–149
- Bankovich AJ, Garcia KC: Not just any T-cell receptor will do. *Immunity* 2003, 18:7–11
- Wucherpfennig KW, Newcombe J, Li H, Keddy C, Cuzner ML, Hafler DA: $\gamma\delta$ T-cell receptor repertoire in acute multiple sclerosis lesions. *Proc Natl Acad Sci USA* 1992, 89:4588–4592
- Oksenberg JR, Panzara MA, Begovich AB, Mitchell D, Erlich HA, Murray RS, Shimonkevitz R, Sherritt M, Rothbard J, Bernard CCA, Steinman L: Selection for T-cell receptor V β -D β -J β gene rearrangements with specificity for a myelin basic protein peptide in brain lesions of multiple sclerosis. *Nature* 1993, 362:68–70
- O'Hanlon TP, Dalakas MC, Plotz PH, Miller FW: Predominant TCR- $\alpha\beta$ variable and joining gene expression by muscle-infiltrating lymphocytes in the idiopathic inflammatory myopathies. *J Immunol* 1994, 152:2569–2576
- Lindberg C, Oldfors A, Tarkowski A: Restricted use of T-cell receptor V genes in endomysial infiltrates of patients with inflammatory myopathies. *Eur J Immunol* 1994, 24:2659–2663
- Hohlfeld R, Engel AG, Li K, Harper MC: Polymyositis mediated by T lymphocytes that express the $\gamma\delta$ receptor. *N Engl J Med* 1991, 324:877–881
- Pluschke G, Rüegg D, Hohlfeld R, Engel AG: Autoaggressive myocytotoxic T lymphocytes expressing an unusual $\gamma\delta$ T-cell receptor. *J Exp Med* 1992, 176:1785–1789
- Mantegazza R, Andreetta F, Bernasconi P, Baggi F, Oksenberg JR, Simoncini O, Mora M, Cornelio F, Steinman L: Analysis of T-cell receptor repertoire of muscle-infiltrating T lymphocytes in polymyositis. *J Clin Invest* 1993, 91:2880–2886
- Bender A, Ernst N, Iglesias A, Dornmair K, Wekerle H, Hohlfeld R: T-cell receptor repertoire in polymyositis: clonal expansion of autoaggressive CD8⁺ T cells. *J Exp Med* 1995, 181:1863–1868
- Nishio J, Suzuki M, Miyasaka N, Kohsaka H: Clonal biases of peripheral CD8 T-cell repertoire directly reflect local inflammation in polymyositis. *J Immunol* 2001, 167:4051–4058

18. Benveniste O, Cherin P, Maisonneuve T, Merat R, Chosidow O, Mouthon L, Guillevin L, Flahault A, Burland MC, Klatzmann D, Herson S, Boyer O: Severe perturbations of the blood T-cell repertoire in polymyositis, but not dermatomyositis patients. *J Immunol* 2001, 167: 3521–3529
19. Goebels N, Michaelis D, Engelhardt M, Huber S, Bender A, Pongratz D, Johnson MA, Wekerle H, Tschopp J, Jenne D, Hohlfield R: Differential expression of perforin in muscle-infiltrating T cells in polymyositis and dermatomyositis. *J Clin Invest* 1996, 97:2905–2910
20. Janson CH, Tehrani MJ, Mellstedt H, Wigzell H: Anti-idiotypic monoclonal antibody to a T-cell chronic lymphatic leukemia: characterization of the antibody, in vitro effector functions and results of therapy. *Cancer Immunol Immunother* 1989, 28:225–232
21. DerSimonian H, Band H, Brenner MB: Increased frequency of T-cell receptor V α 12.1 expression on CD8⁺ T cells: evidence that V α participates in shaping the peripheral T-cell repertoire. *J Exp Med* 1991, 174:639–648
22. Padovan E, Casorati G, Dellabona P, Meyer S, Brockhaus M, Lanzavecchia A: Expression of two T-cell receptor α chains: dual receptor T cells. *Science* 1993, 262:422–424
23. Godfrey DI, Hammond KJL, Poulton LD, Smyth MJ, Baxter AG: NKT cells: facts, functions and fallacies. *Immunol Today* 2000, 21:573–583
24. Park SH, Bendelac A: CD1-restricted T-cell responses and microbial infection. *Nature* 2000, 406:788–792
25. Hingorani R, Choi I-H, Akolkar P, Gulwani-Akolkar B, Pergolizzi R, Silver J, Gregersen PK: Clonal predominance of T-cell receptors within the CD8⁺ CD45RO⁺ subset in normal human subjects. *J Immunol* 1993, 151:5762–5769
26. Posnett DN, Sinha R, Kabak S, Russo C: Clonal populations of T cells in normal elderly humans: the T cell equivalent to “benign monoclonal gammopathy”. *J Exp Med* 1994, 179:609–618
27. Schwab R, Szabo P, Manavalan JS, Weksler M, Posnett DN, Pannetier C, Kourilsky P, Even J: Expanded CD4⁺ and CD8⁺ T-cell clones in elderly humans. *J Immunol* 1997, 158:4493–4499
28. Chamberlain WD, Falta MT, Kotzin BL: Functional subsets within clonally expanded CD8(+) memory T cells in elderly humans. *Clin Immunol* 2000, 94:160–172
29. Hohlfield R: Hunting (auto)immune T cells in neuroimmunological diseases. *Brain* 2003, 126:2–4
30. Hohlfield R, Engel AG: The immunobiology of muscle. *Immunol Today* 1994, 15:269–274
31. Frohman MA, Dush MK, Martin GR: Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene-specific oligonucleotide primer. *Proc Natl Acad Sci USA* 1988, 85:8998–9002
32. Loh EY, Elliott JF, Cwirla S, Lanier LL, Davis MM: Polymerase chain reaction with single-sided specificity: analysis of T-cell receptor δ chain. *Science* 1989, 243:217–220
33. Uematsu Y, Wege H, Straus A, Ott M, Bannwarth W, Lanchbury, Panayi G, Steinmetz MJ: The T-cell receptor repertoire in the synovial fluid of a patient with rheumatoid arthritis is polyclonal. *Proc Natl Acad Sci USA* 1991, 88:8534–8538
34. Choi YW, Kotzin B, Herron L, Callahan J, Marrack P, Kappler J: Interaction of *Staphylococcus aureus* toxin “superantigens” with human T cells. *Proc Natl Acad Sci USA* 1989, 86:8941–8945
35. Pannetier C, Even J, Kourilsky P: T-cell repertoire diversity and clonal expansions in normal and clinical samples. *Immunol Today* 1995, 16:176–181
36. Küppers R, Zhao M, Hansmann ML, Rajewsky K: Tracing B-cell development in human germinal centres by molecular analysis of single cells picked from histological sections. *EMBO J* 1993, 12: 4955–4967
37. Schütze K, Lahr G: Identification of expressed genes by laser-mediated manipulation of single cells. *Nat Biotechnol* 1998, 16:737–742
38. Maryanski JL, Jongeneel CV, Bucher P, Casanova J-L, Walker PR: Single-cell PCR analysis of TCR repertoires selected by antigen in vivo: a high magnitude CD8 response is comprised of very few clones. *Immunity* 1996, 4:47–55
39. Aifantis I, Azogui O, Feinberg J, Saint-Ruf C, Buer J, von Boehmer H: On the role of the pre-T-cell receptor in $\alpha\beta$ versus $\gamma\delta$ T lineage commitment. *Immunity* 1998, 9:649–655
40. Roers A, Montesinos-Rongen M, Hansmann ML, Rajewsky K, Küppers R: Amplification of TCR β gene rearrangements from micromanipulated single cells: T cells rosetting around Hodgkin and Reed-Sternberg cells in Hodgkin’s disease are polyclonal. *Eur J Immunol* 1998, 28:2424–2431
41. Roers A, Hansmann M-L, Rajewsky K, Küppers R: Single-cell PCR analysis of T helper cells in human lymph node germinal centers. *Am J Pathol* 2000, 156:1067–1071
42. Kurokawa M, Tong J, Matsui T, Masuko-Hongo K, Yabe T, Nishioka K, Yamamoto K, Kato T: Paired cloning of the T-cell receptor α and β genes from a single T cell without the establishment of a T-cell clone. *Clin Exp Immunol* 2001, 123:340–345
43. Heath WR, Carbone FR, Bertolino P, Kelly J, Cose S, Miller JFAP: Expression of two T-cell receptor α chains on the surface of normal murine T cells. *Eur J Immunol* 1995, 25:1617–1623
44. Padovan E, Giachino C, Cella M, Valitutti S, Acuto O, Lanzavecchia A: Normal T lymphocytes can express two different T-cell receptor β chains: implications for the mechanism of allelic exclusion. *J Exp Med* 1995, 181:1587–1591
45. Davodeau F, Peyrat M-A, Romagné F, Necker A, Hallet M-M, Vié H, Bonneville M: Dual T-cell receptor β -chain expression on human T lymphocytes. *J Exp Med* 1995, 181:1391–1398
46. Lin AY, Devaux B, Green A, Sagerstrom C, Elliott JF, Davis MM: Expression of T-cell antigen receptor heterodimers in a lipid-linked form. *Science* 1990, 249:677–679
47. Gregoire C, Rebai N, Schweisguth F, Necker A, Mazza G, Auphan N, Millward A, Schmitt-Verhulst AM, Malissen B: Engineered secreted T-cell receptor $\alpha\beta$ heterodimers. *Proc Natl Acad Sci USA* 1991, 88:8077–8081
48. Weber S, Traunecker A, Oliveri F, Gerhard W, Karjalainen K: Specific low-affinity recognition of major histocompatibility complex plus peptide by soluble T-cell receptor. *Nature* 1992, 356:793–796
49. Engel I, Ottenhoff TH, Klausner RD: High-efficiency expression and solubilization of functional T-cell antigen receptor heterodimers. *Science* 1992, 256:1318–1321
50. Chang HC, Bao Z, Yao Y, Tse AG, Goyarts EC, Madsen M, Kawasaki E, Brauer PP, Sacchetti JC, Nathenson SG, Reinherz EL: A general method for facilitating heterodimeric pairing between two proteins: application to expression of α and β T-cell receptor extracellular segments. *Proc Natl Acad Sci USA* 1994, 91:11408–11412
51. Garcia KC, Tallquist MD, Pease LR, Brunmark A, Scott CA, Degano M, Stura EA, Peterson PA, Wilson IA, Teyton L: $\alpha\beta$ T-cell receptor interactions with syngeneic and allogeneic ligands: affinity measurements and crystallization. *Proc Natl Acad Sci USA* 1997, 94:13838–13843
52. Golden A, Khandekar SS, Osburne MS, Kawasaki E, Reinherz EL, Grossman TH: High-level production of a secreted, heterodimeric $\alpha\beta$ murine T-cell receptor in *Escherichia coli*. *J Immunol Methods* 1997, 206:163–169
53. Garboczi DN, Utz U, Ghosh P, Seth A, Kim J, Van Tienhoven EAE, Biddison WE, Wiley DC: Assembly, specific binding and crystallization of a human TCR- $\alpha\beta$ with an antigenic Tax peptide from human T lymphotropic virus type 1 and the class I MHC molecule HLA-A2. *J Immunol* 1996, 157:5403–5410
54. Ding Y-H, Baker BM, Garboczi DN, Biddison WE, Wiley DC: Four A6-TCR/peptide/HLA-A2 structures that generate very different T-cell signals are nearly identical. *Immunity* 1999, 11:45–56
55. Novotny J, Ganju RK, Smiley ST, Hussey RE, Luther MA, Recny MA, Siliciano RF, Reinherz EL: A soluble single-chain T-cell receptor fragment endowed with antigen-combining properties. *Proc Natl Acad Sci USA* 1991, 88:8646–8650
56. Hilyard KL, Reyburn H, Chung S, Bell JL, Strominger JL: Binding of soluble natural ligands to a soluble human T-cell receptor fragment produced in *Escherichia coli*. *Proc Natl Acad Sci USA* 1994, 91:9057–9061
57. Plaksin D, Chacko S, McPhie P, Bax A, Padlan EA, Margulies DH: T-cell receptor V α domain expressed in bacteria: does it dimerize in solution? *J Exp Med* 1996, 184:1251–1258
58. Gao GF, Rao Z, Bell JL: Molecular coordination of $\alpha\beta$ T-cell receptors and coreceptors CD8 and CD4 in their recognition of peptide-MHC ligands. *Trends Immunol* 2002, 23:408–413
59. Ohashi PS, Mak TW, van den Elsen P, Yanagi Y, Yoshikai Y, Calman AF, Terhorst C, Stobo JD, Weiss A: Reconstitution of an active surface T3/T-cell antigen receptor by DNA transfer. *Nature* 1985, 316:606–609
60. Dembic Z, Haas W, Weiss S, McCubrey J, Kiefer H, von Boehmer H,

- steinmetz M: Transfer of specificity by murine α and β T-cell receptor genes. *Nature* 1986, 320:232–238
61. Saito T, Germain RN: Predictable acquisition of a new MHC recognition specificity following expression of a transfected T-cell receptor β -chain gene. *Nature* 1987, 329:256–259
62. Zumla A, McCormack A, George A, Batchelor JR, Lechler R: Use of murine T-cell hybridoma expressing human T-cell receptor α - and β -gene products as tool for the production of human T-cell receptor-specific monoclonal antibodies. *Hum Immunol* 1992, 35:141–148
63. Choi Y, Kappler JW, Marrack P: A superantigen encoded in the open reading frame of the 3' long terminal repeat of mouse mammary tumor virus. *Nature* 1991, 350:203–207
64. Saito T, Weiss A, Miller J, Norcross MA, Germain RN: Specific antigen-Ia activation of transfected human T cells expressing murine T $\alpha\beta$ -human T3 receptor complexes. *Nature* 1987, 325:125–130
65. Geisler C, Rubin B, Caspar-Bauguil S, Champagne E, Vangsted A, Hou X, Gajhede M: Structural mutations of C-domains in members of the Ig superfamily: consequences for the interactions between the T-cell antigen receptor and the ζ 2 homodimer. *J Immunol* 1992, 148:3469–3477
66. Howard FD, Moingeon P, Moebius U, McConkey DJ, Yandava B, Gennert TE, Reinherz EL: The CD3 ζ cytoplasmic domain mediates CD2-induced T-cell activation. *J Exp Med* 1992, 176:139–145
67. Letourneur F, Malissen B: Derivation of a T-cell hybridoma variant deprived of functional T-cell receptor α - and β -chain transcripts reveals a nonfunctional α -mRNA of BW5147 origin. *Eur J Immunol* 1989, 19:2269–2274
68. Kappler JW, Skidmore B, White J, Marrack P: Antigen-inducible, H-2-restricted, interleukin-2-producing T-cell hybridomas: lack of independent antigen and H-2 recognition. *J Exp Med* 1981, 153:1198–1214
69. Blank U, Boitel B, Mège D, Ermonval M, Acuto O: Analysis of tetanus toxin peptide/DR recognition by human T-cell receptors reconstituted into a murine T-cell hybridoma. *Eur J Immunol* 1993, 23:3057–3065
70. Stotz SH, Bolliger L, Carbone FR, Palmer E: T-cell receptor (TCR) antagonism without a negative signal: evidence from T-cell hybridomas expressing two independent TCRs. *J Exp Med* 1999, 189:253–264
71. Kessels HWHG, Wolkers MC, Schumacher TNM: Adoptive transfer of T-cell immunity. *Trends Immunol* 2002, 23:264–269
72. Wiendl H, Malotka J, Holzwarth B, Weltzien HU, Wekerle H, Hohlfield R, Dornmair K: An autoreactive $\gamma\delta$ TCR derived from a polymyositis lesion. *J Immunol* 2002, 169:515–521
73. Vollmer J, Weltzien H-U, Moulon C: TCR reactivity in human nickel allergy indicates contacts with complementarity-determining region 3 but excludes superantigen-like regions. *J Immunol* 1999, 163:2723–2731
74. Vollmer J, Weltzien H-U, Dormoy A, Pistor F, Moulon C: Functional expression and analysis of a human HLA-DQ restricted, nickel-reactive T-cell receptor in mouse hybridoma cells. *J Invest Dermatol* 1999, 113:175–181
75. Madsen LS, Andersson EC, Jansson L, Krogsgaard M, Andersen CB, Engsborg J, Strominger JL, Svejgaard A, Holmdahl R, Wucherpfennig KW, Fugger L: A humanized model for multiple sclerosis using HLA DR2 and a human T-cell receptor. *Nat Genet* 1999, 23:343–347
76. Ji H, Korganow A-S, Mangialaio S, Höglund P, André I, Lühder F, Gonzalez A, Poirot L, Benoist C, Mathis D: Different models of pathogenesis in T-cell dependent autoimmunity: clues from two TCR transgenic systems. *Immunol Rev* 1999, 169:139–146
77. Goverman J: Tolerance and autoimmunity in TCR transgenic mice specific for myelin basic protein. *Immunol Rev* 1999, 169:147–159
78. Fugger L: Human autoimmunity genes in mice. *Curr Opin Immunol* 2000, 12:698–703
79. Boyton RJ, Altmann DM: Transgenic models of autoimmune disease. *Clin Exp Immunol* 2002, 127:4–11
80. Hayday AC: $\gamma\delta$ T cells: a right time and a right place for a conserved third way of protection. *Annu Rev Immunol* 2000, 18:975–1026
81. Morita CT, Mariuzza RA, Brenner MB: Antigen recognition by human gamma delta T cells: pattern recognition by the adaptive immune system. *Springer Semin Immunopathol* 2000, 22:191–217
82. Carding SR, Egan PJ: $\gamma\delta$ T cells: functional plasticity and heterogeneity. *Nat Rev Immunol* 2002, 2:336–345
83. Davis SJ, Ikemizu S, Evans EJ, Fugger L, Bakker TR, van der Merwe AP: The nature of molecular recognition by T cells. *Nat Immunol* 2003, 4:217–224
84. Hemmer B, Fleckenstein BT, Vergelli M, Jung G, McFarland H, Martin R, Wiesmüller K-H: Identification of high-potency microbial and self ligands for a human autoreactive class II-restricted T-cell clone. *J Exp Med* 1997, 185:1651–1659
85. Wiesmüller KH, Fleckenstein B, Jung G: Peptide vaccines and peptide libraries. *Biol Chem* 2001, 382:571–579
86. Jacobsen M, Cepok S, Oertel WH, Sommer N, Hemmer B: New approaches to dissect degeneracy and specificity in T-cell antigen recognition. *J Mol Med* 2001, 79:358–367
87. Borras E, Martin R, Judkowski V, Shukaliak J, Zhao Y, Rubio-Godoy V, Valmori D, Wilson D, Simon R, Houghten R, Pinilla C: Findings on T-cell specificity revealed by synthetic combinatorial libraries. *J Immunol Meth* 2002, 267:79–97
88. Crowley MP, Fahrer AM, Baumgarth N, Hampl J, Gütgemann I, Teyton L, Chien Y-H: A population of murine $\gamma\delta$ T cells that recognize an inducible MHC class Ib molecule. *Science* 2000, 287:314–316
89. Hampl J, Schild H, Litzemberger C, Baron M, Crowley MP, Chien, Y-H: The specificity of a weak $\gamma\delta$ TCR interaction can be modulated by the glycosylation of the ligand. *J Immunol* 1999, 163:288–294