Distinctive Roles of Neutrophils and Monocytes in Anti-Thy-1 Nephritis

Ralf Westerhuis,* Suzanne C. van Straaten,[†]
Mieneke G. A. van Dixhoorn,* Nico van Rooijen,[‡]
Nicole A. M. Verhagen,* Christine D. Dijkstra,[‡]
Emile de Heer,[†] and Mohamed R. Daha*

From the Departments of Nephrology* and Pathology,[†] Leiden University Medical Center, University of Leiden, Leiden; and Department of Cell Biology and Immunology,[‡] Medical Faculty, Free University, Amsterdam, The Netherlands

Anti-Thy-1.1 glomerulonephritis as an experimental model for mesangial proliferative glomerulonephritis was induced in Wistar rats by a single injection of monoclonal IgG2a-anti-Thy-1.1 antibody (ER4G). This transient model is complement-mediated and leads to mesangial-cell (MC) lysis followed by MC proliferation, glomerular microaneurysm formation, glomerular influx of polymorphonuclear leukocytes (PMNs) and macrophages, proteinuria, and hematuria. In this study we investigated the distinctive roles of infiltrating PMNs or monocytes/macrophages by treating rats with an antibody against rat integrin CD11b/CD18 (ED7) or by depletion of monocytes with multilamellar clodronate liposomes, respectively. ED7 administration resulted in reduction of the influx of PMNs in glomeruli during the first 6 days after induction of Thy-1.1 nephritis, whereas treatment with an isotype-matched irrelevant antibody (PEN9) or with phosphate-buffered saline had no effect on macrophage influx. Increased glomerular C3 and C6 deposition on days 1 and 3 was seen in the ED7-treated rats but not seen in the control groups. In addition, the ED7-treated group showed an increased number of aneurysmatic glomeruli and more severe hematuria. Monocyte/macrophage depletion led to a significant reduction of mesangial matrix expansion, although mesangial proliferation, proteinuria, and hematuria remained unaltered. These results, together with the known effects of PMN-derived enzymes on C3 cleavage, suggest that a reduction in the influx of PMNs results in sparing of C3 and consequently of more complement activation in the glomerulus with increased complement-mediated damage. Our data indicate that infiltrating PMNs and monocytes/macrophages play distinctive roles during inflammation in this model of MC glomerulonephritis. (*Am J Pathol* 2000, 156:303–310)

The contribution of polymorphonuclear leukocytes (PMNs) and macrophages in glomerular injury has been well established. The influx of neutrophils during anti-thymus nephritis is well documented,¹ but its role in the pathogenesis is largely unexplored.² Injection of antibodies against Thy-1.1, a transmembrane glycoprotein on mesangial cells,^{3,4} results in complement-dependent mesangial-cell lysis,^{1,5,6} apoptosis,⁷ and subsequent mesangial proliferation⁸ and extracellular-matrix expansion.⁹ Previous studies have shown that mesangial-cell injury and the subsequent proliferative response depend on complement activation and that the response can be suppressed by decapentation with cobra venom factor.^{2,10} Similar results were demonstrated by Bagchus et al,¹¹ who found no glomerular lesions after the injection of a non-complement-fixing monoclonal anti-Thy-1 antibody. During the development of mesangioproliferative glomerulonephritis, the glomeruli are infiltrated by neutrophils^{1,2,12} and monocytes.¹³

Glomerular infiltration by neutrophils is characteristic of acute experimental glomerulonephritis.^{14,15} Proteinuria resulting from neutrophil-mediated glomerular injury has been shown in nephrotoxic serum nephritis^{9,16} and crescentic glomerulonephritis¹⁷ and after intrarenal injection of phorbol myristate acetate (PMA)¹⁸ or of cobra venom factor, which causes complement activation and subsequent PMN chemotaxis.^{12,19} In each of these described models, PMN depletion markedly diminished proteinuria and resulted in reduction or attenuation of renal disease.¹⁷

The β_2 -integrin CD11b/CD18 has been shown to be involved in the influx of PMNs and monocytes/macrophages in different reperfusion models in rabbit²⁰ and experimental allergic encephalomyelitis (EAE) and acute colitis in rats.^{21,22} In the latter models, a mouse monoclonal antibody (mAb) against CD11b (ED7) was able to reduce the influx of inflammatory cells (predominantly PMNs) into the inflamed tissues and also the severity of the induced disease. In EAE, no effect of administration of anti-CD11b antibodies ED7 or ED8 was observed on the cellular infiltration (predominantly monocytes), but clinical severity was significantly reduced in the anti CD11b-treated EAE rats.²¹

The involvement of monocytes and macrophages in various glomerulonephritides has been clearly estab-

Accepted for publication September 16, 1999.

Address reprint requests to M. R. Daha, Ph.D., Department of Nephrology, Leiden University Medical Center, Building 1, C3P, PO Box 9600, 3200 RC Leiden, The Netherlands. E-mail: daha@mail.medfac.leidenuniv.nl.

lished.²³ In particular their involvement has been associated with proliferative forms of glomerulopathies.^{24,25} Experimental evidence has been provided by rodent models of glomerulonephritis.^{26–28} However, in some cases infiltrating monocytes may serve merely to remove immune complexes and not to contribute to the glomerular injury.²⁹ Recent interest has focused on the role of macrophages in the pathogenesis of focal glomerulosclerosis.^{30,31} Both glomerular hypercellularity and expansion of the extracellular matrix are thought to be of primary importance in the development of capillary obsolescence and glomerulosclerosis. In the remnant kidney model in the rat, Van Goor and colleagues²³ showed that macrophages play a central role in the development of focal glomerulosclerosis. As described by Floege et al, the development of glomerulosclerosis in this model is preceded by mesangial proliferation and mesangial matrix expansion.³²

In this study we investigated the role of glomerular monocytes and PMNs in the development of mesangial injury and subsequent glomerular hypercellularity and mesangial matrix expansion. The PMN infiltration into the glomeruli during Thy-1 nephritis was affected by pretreatment with monoclonal antibody ED7. Monocyte depletion was performed using a macrophage suicide technique, by injecting liposomes, in which clodronate was encapsulated.³³ This study shows that each of these inflammatory cells plays a distinctive role in the pathogenesis of mesangioproliferative glomerulonephritis.

Materials and Methods

Animals

Female inbred Wistar rats (160–185 g) were obtained from the animal facilities of the Department of Pathology, Leiden University Medical Center, Leiden, The Netherlands. The rats were housed in accredited animal facilities and fed pelleted food, and they had access to water *ad libitum*. Experiments were performed in accord with Dutch legislation for the care and use of laboratory animals.

Antibodies

mAb against Thy-1.1 (mouse IgG2a/ κ against rat CD90) was derived from hybridoma ER4G as previously described.¹¹ It was purified from ascitic fluid by affinity purification on protein A-Sepharose 4B (Pharmacia, Uppsala, Sweden). Rabbit anti-mouse IgG was obtained from Jansen (Beerse, Belgium), and goat anti-rat fibrinogen-fluorescein isothiocyanate (FITC) was from Nordic (Tilburg, The Netherlands). The ED1 antibody is a murine monoclonal IgG1 to a cytoplasmic antigen present in monocytes, macrophages, and dendritic cells.³⁴ PC10 (Dako, Glostrup, Denmark) is a murine immunoglobulin M (IgM) mAb against proliferating cell nuclear antigen (PCNA), which is expressed by actively proliferating mesangial cells.³⁵ Mouse mAb directed against C6 was kindly provided by Dr. W. G. Couser (Division of Nephrology, University of Washington, Medical Center, Seattle,

WA). FITC-conjugated rabbit anti-rat C3 antibodies were generated in our own laboratory. Rat mAb anti-mouse IgG1-horseradish peroxidase (HRP) was from Sanbio (Uden, The Netherlands). The accumulation of platelets was detected by mAb PI-1.³³ The glomerular influx of PMNs was assessed with FITC-labeled mAb W3/13. After the original description of its tissue distribution³⁶ and purification,³⁷ the antigen recognized by mAb W3/13 has been identified as leukosialin,³⁸ which is present on T cells, neutrophils, and brain from the rat. After molecular cloning of leukosialin, it has been designated as CD43. The expression and function of CD43 on neutrophil granulocytes have been described in several studies.^{39,40}

It is well established that, during the course of anti-Thy-1 nephritis, an influx of neutrophils and monocytes/macrophages takes place.¹ However, there has been no T cell infiltration, during this type of experimental mesangial cell-proliferative glomerulonephritis, that has been detectable with mAb R73 against the T cell receptor, although CD43-positive PMNs were observed with mAb W3/13. Therefore, our study and several earlier ones chose mAb W3/13 for immunohistochemical detection of PMNs in kidney sections of rats.¹ ED7 is an mAb against cell adhesion molecule CD11b/CD18 and was prepared as described earlier.⁴¹ PEN9 is specific for penicillin and was used as an isotype control for ED7. PEN9 has been characterized.¹³

Preparation of Liposomes

Multilamellar liposomes were prepared as described earlier.³³ In brief, 86 mg phosphatidylcholine (a gift from Lipoid KG, Ludwigshafen, Germany) and 8 mg cholesterol, molar ratio 6:1, were dissolved in 20 ml methanol/chloroform (1:1) in a round-bottomed flask. The thin film that formed on the interior of the flask after low-vacuum rotary evaporation at 37°C was dispersed in 10 ml phosphate-buffered saline (PBS; 10 mmol/L, pH 7.4), containing either 2.5 g dichloromethylene diphosphonate (Cl₂MDP; a gift from Boehringer Mannheim GmbH, Mannheim, Germany) or PBS, by gentle rotation for 10 min. Free Cl₂MDP was removed by rinsing the liposomes with PBS and centrifuging them for 30 min at 100,000 × *g* at 16°C. The liposomes were then resuspended in 4 ml of PBS; 2 ml were intravenously injected per rat. Controls received 2 ml of PBS-encapsulated liposomes. The efficacy of depletion was evaluated by immunohistology with ED1 on kidney biopsies taken after 24 hours. Pilot experiments had shown that administration of higher concentrations of Cl₂MDP-liposomes led to a profound complement depletion, resulting in a decrease of proteinuria development and subsequent reduction of anti-Thy-1-induced pathology (data not shown). Therefore a liposome dose was chosen that did not affect the CH50 levels in the blood.

Induction of Anti-Thy-1 Glomerulonephritis

Mesangial proliferative glomerulonephritis was induced by injection of mAb against Thy1.1 (ER4G) at a dose of 1

mg/kg intravenously in the rat tail vein. The animals were housed in metabolic cages, and 24-hour urine protein excretion was measured daily, using the biuret standard method. Urine containing 0.1% merthiolate was used. Hematuria was assessed with dipsticks (Hema-Combisticks; Bayer Diagnostics, Mijdrecht, The Netherlands), which have been shown to be as sensitive as direct microscopic enumeration of erythrocytes in full urine. Moreover, the used dipstick can be examined to distinguish between small differences in the amounts of erythrocyturia. Because the values are comparable with those obtained by a colorimetric method and a direct counting of erythrocytes, we used the values obtained from dipsticks.¹

Experimental Design

For ED7-treatment, 21 young female Wistar rats were divided into three groups of 7 rats each. To determine the role of PMNs during anti-Thy-1.1 nephritis, one group of rats was injected, 2 hours before and 3 days after injection of ER4G, with 0.5 mg of mAb ED7. As controls, we used two groups of seven rats each, injected with PBS alone or with an isotype-matched irrelevant antibody, PEN9 (0.5 mg dissolved in equal amounts of PBS as for ED7). All antibodies were administered by tail vein injection.

From all rats, renal biopsies were taken 1, 3, 6, and 10 days after injection of ER4G. Urine samples (24 hours) were collected before the experiment and on days 1, 2, 4, 6, 8, and 10.

For macrophage depletion studies, 54 female Wistar rats were used. One group of rats ($n = 24$) was injected with 1 ml/100 g of body weight liposome-encapsulated Cl_2MDP at day -1 and with 1 mg/kg mAb ER4G at day 0. Kidney biopsies ($n = 8$ at each time point) were performed at 1, 4, and 24 hours after ER4G injection. At days 2, 6, and 14, eight animals were sacrificed. Corresponding groups of rats served as controls; six animals received liposome-encapsulated Cl_2MDP followed by saline injection; in nine animals, only ER4G was injected at day 0; eight animals received liposome-encapsulated PBS followed by ER4G on day 0; four animals received liposome-encapsulated PBS followed by saline injection; six animals received saline only.

Blood samples were collected from the tail vein before the injection of Cl_2MDP and immediately before the administration of ER4G to evaluate CH50 levels.

All intravenous injections, blood samples, and biopsies were performed under ether anesthesia. At sacrifice, the kidneys were perfused *in situ* with PBS. Cortical tissue was processed for light and immunofluorescence microscopy. All experiments were carried out in two distinct sets. One complete set of experiments is presented in this study.

Histological Examination

Kidney tissue was obtained for light and immunofluorescence microscopy. For light microscopy, tissues were fixed in methacarn solution, dehydrated in graded eth-

anols, and embedded in paraffin. Sections (4- μm each) were stained with periodic acid/Schiff (PAS) reagent. The degrees of glomerular mesangiolysis (glomerular aneurysms)⁶ and glomerular extracellular matrix expansion were graded semiquantitatively in 25 representative glomerular cross sections per rat, as described previously by Floege et al.⁴² Mean values per biopsy were calculated for the number of proliferating (PCNA⁺) cells and monocytes/macrophages per glomerular cross section. An indirect immunoperoxidase method was used for the identification of monocytes with mAb ED1 and for PCNA with PC10. Specific antibody binding was revealed by the reaction with hydrogen peroxide and diaminobenzidine (Sigma Chemical Co., St. Louis, MO). Normal rat spleen served for ED1 as positive control. Immunofluorescence microscopy was performed on tissue samples snap-frozen in CO_2 -ice-cooled isopentane and stored at -70°C . Cryostat sections of 3 μm each were obtained and, after air-drying, fixed in acetone for 10 minutes at room temperature. The slides were washed twice in PBS and examined for the presence of rat C3 by a directly FITC-conjugated antibody. For the detection of PMNs and C6, tissue sections were preincubated with PBS containing a 1:300 dilution of a 30% H_2O_2 solution (Merck, Darmstadt, Germany), to block endogenous peroxidases. Thereafter the slides were washed and incubated with digoxigenin (DIG)-conjugated mouse monoclonal anti-rat C6 or the DIG-conjugated mAb W3/13,¹ in 0.5% Boehringer blocking reagent (TNB) after incubation with HRP-conjugated Sheep F(ab') anti-DIG fragments. The slides were subsequently incubated with tyramide-FITC (NEN-Dupont Research Products, Boston, MA) for 30 minutes at room temperature.¹ Mean values per biopsy were calculated for the number of PMNs per glomerular cross-section. Glomerular C3, C6, fibronectin, and IgG2a depositions were scored by a semiquantitative method as previously described,⁴² with scores ranging from 0 to 4 (0 = 0–5%, 1 = 6–25%, 2 = 26–50%, 3 = 51–75%, 4 > 75% of the glomerular cross-section).

The slides were coded, and 20 glomerular cross-sections per rat were scored independently for each parameter by two experienced microscopists. Photographs were taken on Kodak TX-400 film on a Leitz microscope equipped with a 4-mm BG 38 + 5-mm BG 12 filter for FITC.

Statistical Analysis

Mean values ($\pm\text{SD}$) for each parameter were calculated and compared in an independent Student's *t*-test. A *P* value smaller than 0.05 was considered significant.

Results

Anti-CD-11b Treatment

Glomerular binding of mouse mAb ER4G showed a mesangial pattern with no differences between the ED7-treated and control groups. The antibody was detectable in the rat glomerulus until day 3. The influx of PMNs into the glomerular cross-section as assessed by detection

with W3/13 was maximal on day 1. The PBS- and the PEN9-treated groups revealed 2.0 ± 0.2 and 2.1 ± 0.2 PMNs per glomerular cross-section, respectively. The number of PMNs detected in the ED7-treated group was significantly reduced to 1.6 ± 0.1 PMNs per glomerular cross-section ($P < 0.05$). A comparable reduction of PMNs was measured on days 3 and 6 of the experiment with highly significant values (Figures 1 and 2). Equal results of glomerular PMN counts were obtained by using PAS-stained sections.

The influx of monocytes/macrophages into the glomerular area did not reveal any differences between the various groups. In all rats, we observed a peak influx of this cell type at day 1, followed by a subsequent decline to baseline levels at day 10 (data not shown).

The glomerular deposition of C3 and C6 was scored semiquantitatively after indirect immunohistochemical staining of frozen kidney sections of each rat. Significantly higher scores for C3 and C6 deposition were found on day 1 for the rats treated with ED7 as compared with the PBS- and the PEN9-injected rats. C6 deposition was seen only in rats treated with ED7 on day 3. At later time points, no C6 deposition was detected in any of the experimental animals. Glomerular C3 deposition was highest on day 1 in the ED7-treated group and significantly increased as compared with the controls. On day 6, only ED7-treated rats exhibited some C3 deposition (Figures 1 and 2).

When the biopsies of the rats were analyzed by light microscopy, we observed a significantly higher maximum of glomerular microaneurysms ($37.1 \pm 16\%$) on day 3 for the rats receiving ED7. The glomerular lesions of the control groups were approximately 50% less compared with ED7-treated animals (Figures 2 and 3). The extracellular matrix/fibronectin score was not significantly different between the ED7-treated and the control groups. Also, the number of proliferating glomerular cells was not changed after reduction of glomerular influx of PMNs.

Hematuria and Proteinuria

Hematuria from day 4 and later in this experiment showed, for the ED7-treated rats, higher values as compared with the control groups. Because of the high standard deviations in all groups, this increase was significant only on days 2, 8, and 10 (Figure 3). For proteinuria we did not find significant differences between the groups (data not shown).

Monocyte and Macrophage Depletion Studies

Rats treated with clodronate liposomes exhibited a nearly complete reduction of glomerular monocyte infiltration. Although control rats showed about 4.2 ± 1.8 cells per glomerulus, the monocyte-depleted rats had 0.2 ± 0.25 ED1⁺ cells ($P < 0.001$; Figure 4).

A reduction of $15 \pm 23\%$ of circulating monocytes was found. It is interesting that no additional effect of ED7 was seen on the clodronate-mediated reduction in circulating monocytes.

Figure 1. A: PMN influx in ED7-treated (white bars), PEN9-treated (hatched bars), and PBS-treated (black bars) rats. Immunohistochemical staining for intraglomerular PMNs is shown as mean $\pm$ SEM number of W3/13-positive cells/glomerular cross section per group. ED7 pretreatment led to significant reduction of PMN influx during anti-Thy-1.1 nephritis until day 6. At day 10, no significant difference was detected. B: Immunohistochemical staining for glomerular C3 deposition after induction of anti-Thy-1.1 nephritis, using a semiquantitative scoring system (shown as mean $\pm$ SEM). Higher scores for C3 were observed on days 1 and 3 for the ED7-pretreated group as compared with the control groups. Only the ED7-pretreated rats showed residual C3 deposition on day 6. C: Immunohistochemical detection for glomerular C6 deposition was found to be higher on day 1 in the ED7 group (shown as mean $\pm$ SEM). On day 3 only little C6 deposition was still found in this group, whereas the control groups did not show positivity for C6.

The influx of PMNs into the glomeruli was not affected by the monocyte/macrophage depletion. Glomerular deposition of C3 and C6 was not changed at all by macrophage depletion. All groups showed the expected

Figure 2. **A** and **B:** Immunohistochemical staining for PMN influx in ED7- and PEN9-/PBS-treated rats was performed using mAb W3/13. **A:** ED7 treatment led to significant reduction of PMN influx during anti-Thy-1.1 nephritis. A representative glomerulus obtained on day 1 is shown. **B:** The control groups showed significantly higher values of W3/13-positive cells. **C** and **D:** Immunohistochemical staining for glomerular C3 deposition on day 1 after induction of anti-Thy-1.1 nephritis are shown. **D:** Rats treated with control antibody PEN9 or PBS showed less deposition of C3 as compared with rats treated with ED7 (**C**). **E** and **F:** Glomerular microaneurysms were scored in PAS-stained kidney sections. The ED7-treated group of rats developed significantly more glomerular microaneurysms (**E**) on day 3 as compared with rats treated with PBS or PEN9 (**F**).

degree of complement deposition as described for the control groups in the ED-7 experiment.

In the PAS-staining, we observed no differences for glomerular microaneurysms/mesangiolytic. However, a significant difference in extracellular matrix scores was found between the two groups (Figure 4). The glomerular deposition of fibronectin corresponded to the difference in mesangial extracellular matrix expansion (data not shown).

We did not observe significant differences in proliferating (PCNA⁺) mesangial cells between the different groups of rats (data not shown).

Proteinuria and Hematuria

In monocyte-depleted Wistar rats, proteinuria and hematuria were comparable with that in the control rats (data not shown).

Figure 3. **A:** Course of hematuria during the experiment detected by dip stick. At all time points, hematuria of the ED7-treated group was highest compared with the control groups. Because of the high standard deviations, only the differences on days 2, 8, and 10 were significant. The data show mean values per group $\pm$ SEM. **B:** Detection of glomerular microaneurysms by scoring PAS-stained kidney sections (mean values per group $\pm$ SEM). The ED7-pretreated group developed a significantly higher percentage of glomeruli with microaneurysms on day 3 after induction of nephritis. The aneurysms detected on the other days of the experiment were not significantly different.

Discussion

In this study we demonstrate that treatment of rats with ED-7, a mouse anti-CD11b mAb, selectively reduces glomerular PMN influx during the course of anti-Thy-1.1 nephritis. PMN reduction was associated with enhanced hematuria, a higher percentage of glomeruli with microaneurysms, enhanced glomerular deposition of C3 and C6, and no significant change in proteinuria, suggesting that microaneurysm formation is an important factor in determining the degree of hematuria. It is known that PMNs contribute to tissue injury during ischemic reperfusion of the pulmonary, coronary, cerebral, and splanchnic circulation. Injection of antibodies against CD18 confers protection against ischemic-reperfusion injury of these organs in many experimental models.⁴³

Previously it was shown that ED7 is able to reduce the influx of neutrophils during experimental acute colitis²² in rat. Also the damage in the colon was found to be much less. In a peritoneal recruitment assay, ED7 treatment reduced the influx of myelomonocytic cells.²¹ Also, in a lung reperfusion model, an anti-CD18 antibody prevented vascular injury associated with a reduction of myeloperoxidase in the examined tissues.^{20,44}

Figure 4. **A:** Macrophage influx in macrophage-depleted (black bars) and control (white bars) rats. Immunohistochemical staining for intraglomerular macrophages is shown as mean number of ED1-positive cells per glomerulus per group $\pm$ SEM. Macrophage depletion led to an almost complete reduction of infiltrating macrophages on days 1 and 2 after induction of anti-Thy-1.1 nephritis. At day 6, no significant difference was detected. **B:** Extracellular matrix expansion was scored using PAS staining of rat kidneys. During anti-Thy-1.1 nephritis, glomerular extracellular matrix expansion (shown as mean $\pm$ SEM) was significantly reduced at days 6 and 14 after depletion of macrophages as compared with rats not depleted (white bars).

No effect of ED7 treatment on monocyte influx during anti-Thy-1.1 nephritis was observed. The reason for the different behaviors of the two cell types in this model is unknown. A possible explanation could be the major pathogenetic role of complement in this model, in which several chemoattractants are released during the initial phase of injury, which may contribute to the attraction of monocytes.

Earlier experiments assessed the influence of mAb against CD18, CD11a, CD11b, very late antigen-4 (VLA-4), intercellular adhesion molecule-1 (ICAM-1), and E-selectin on glomerular neutrophil accumulation and proteinuria in Long-Evans rats with nephrotoxic-serum nephritis.⁴⁵ This model is characterized by up-regulation of vascular cell adhesion molecule-1 (VCAM-1), ICAM-1, and E-selectin expression, rapid neutrophil infiltration into the glomerulus, and proteinuria. Treatment of animals with antibodies against CD18, CD11b, and ICAM-1 caused 63%, 46%, and 54% reduction, respectively, in proteinuria and 79%, 66%, and 54% reduction, respectively, in glomerular neutrophil counts, suggesting an

important role for these adhesion molecules in this model of experimental nephritis.⁴⁵ Moreover, these data indicate that neutrophils play a major pathogenetic role in experimental nephrotoxic nephritis, specifically in glomerular permselectivity, whereas other factors, for example, complement activation, seem to have a secondary role. These findings are further supported by another study investigating nephrotoxic-serum nephritis.⁴⁶

Our own results for the reduction of glomerular PMN counts are in agreement with the above mentioned studies. On the other hand, we did not find a reduction of hematuria or proteinuria in this complement-mediated experimental glomerulonephritis, suggesting that neutrophils do not significantly contribute to loss of permselectivity in this model. This is in agreement with earlier studies.^{2,6,10} In contrast, PMNs seem to contribute to the repair process during anti-Thy-1.1 nephritis. Maybe proteolytic enzymes released by these cells have a complement-clearing function, and consequently a reduction in PMN influx may be responsible for the observed pronounced and sustained C3 and C6 deposition in rats treated with ED7, compared with the controls. In the literature a complement-clearing function of proteases released from PMNs has been described. It was found that proteolytic enzymes derived from PMNs in pleural empyema or from isolated PMNs can inactivate C3 bound to Sepharose.⁴⁷ Furthermore, cleavage of C3b and C3bi bound to human erythrocytes was shown to depend on purified leukocyte enzymes or crude extracts from human PMNs,^{48,49} indicating a complement-clearing function of PMN-related enzymes. Because complement is the major pathogenetic factor leading to MC lysis in this experimental model, this could be an explanation, on one hand, for the increased damage and hematuria observed in this study. On the other hand, it was found⁵⁰ that incubation of highly purified human C1 inhibitor with equally pure human leukocyte proteinase 3 resulted in a dose- and time-dependent inactivation of C1 inhibitor hemolytic activity, which may lead to an increase of activated complement factors.

The major difference in the above mentioned nephrotoxic nephritis is that, during Thy-1 nephritis, neutrophils and complement together play a role, whereas, in nephrotoxic nephritis, neutrophils seem to be the main factor affecting the permselectivity of the damaged rat kidneys. Neutrophils in nephrotoxic nephritis seem to contribute to mesangial injury, whereas neutrophils during Thy-1 nephritis do not seem to play a major role.

Concerning the role of macrophages, we found that selective depletion of macrophages led to a reduced glomerular matrix expansion. In contrast to ED7 treatment, glomerular complement deposition, microaneurysm formation, and hematuria remained unaffected. These data are in agreement with recently published investigations.⁵¹ It was found that, after induction of anti-Thy-1.1 nephritis, influx of macrophages is strain dependent. Lewis rats showed high amounts of infiltrating macrophages into the rat glomerulus, whereas F344 rats showed no enhancement of these infiltrating cells. Moreover, after transplantation of F344 kidneys in Lewis rats and vice versa, there was influx of macrophages only in

kidneys transplanted to Lewis rats and not in kidneys transplanted to F344 rats. In F344 rats, the influx of macrophages was low, and the extracellular matrix expansion was marginal. In contrast, high amounts of infiltrating macrophages led to a highly significant increase of extracellular matrix in Lewis rats.⁵¹ The mechanisms leading to these differences are still unclear; all grafts showed an equal induction of monocyte chemotactic protein-1 in both rat strains. These data indicate a predominant factor located outside the kidney that leads to mononuclear infiltrates in this experimental disease.

The potential mechanisms by which macrophages mediate glomerular-cell proliferation and the development of mesangial matrix expansion could involve release of cytokines by the macrophage itself or could involve stimulation by the macrophage of other cell types to release cytokines within the glomerulus.^{52,53} During tissue repair processes, macrophages can release TGF- β 1,⁵⁴ a growth factor that has clearly been shown to be involved in mesangial matrix expansion.⁵⁴⁻⁵⁶ A second macrophage-derived product with a possible high impact is nitric oxide, which is abundantly secreted by infiltrating monocytes in this model.⁵⁷ Monocytes have been shown to induce increased transcription of TGF- β and fibronectin.⁵⁸ In coculture studies, Mosquera demonstrated that monocyte-derived culture supernatants could induce mesangial cells to synthesize fibronectin *in vitro*.⁵⁹ As shown in Figure 3, the mesangial expansion in monocyte-depleted rats was not completely inhibited. This may be because 1) especially major histocompatibility complex-II^{neg} resident macrophages are less sensitive to liposome-mediated elimination (E De Heer, V Cattell, unpublished results), and 2) cytokines from other sources, for instance platelet-derived growth factor, may be able to induce mesangial matrix expansion, albeit less effectively.⁵² In conclusion our study presents evidence that monocytes are specifically involved in the expansion of the mesangial extracellular matrix, whereas other immunopathological processes in the mesangium (complement activation, platelet aggregation, and mesangial proliferation) remain unaffected. These findings indicate that the mesangial alterations occur through distinct signaling pathways.

Taken together we hypothesize that a reduced influx of PMNs and a subsequent reduction in generation of C3-cleaving enzymes from PMNs may lead to a relative sparing of activated C3, which, on its own, results in more injury (glomerular aneurysms) and up-regulation of hematuria via a complement-dependent mechanism. Furthermore, we found monocytes being selectively involved in mesangial extracellular matrix expansion, whereas the induction of mesangial proliferation is induced through a separate pathway.

References

1. Van Dixhoorn MG, Gorter A, Sato T, van der Wal AM, van Eendenburg JD, Rozing J, Daha MR, De Heer E: Induction of microhematuria by an IgA isotype switch variant of a monoclonal anti-Thy-1.1 antibody in the rat. *Kidney Int* 1996, 50:1612-1623
2. Yamamoto T, Wilson CB: Complement dependence of antibody-induced mesangial cell injury in the rat. *J Immunol* 1987, 138:3758-3765
3. Sato T, Van Dixhoorn MG, Schroeijers WE, Huizinga TW, Reutelingsperger CP, van Es LA, Daha MR: Apoptosis of cultured rat glomerular mesangial cells induced by IgG2a monoclonal anti-Thy-1 antibodies. *Kidney Int* 1996, 49:403-412
4. Sato T, Van Dixhoorn MG, Schroeijers WE, van Es LA, Daha MR: Efficient induction of apoptosis in cultured rat glomerular mesangial cells by dimeric monoclonal IgA anti-Thy-1 antibodies. *Kidney Int* 1997, 51:173-181
5. Van Dixhoorn MG, Timmerman JJ, van Gijlswijk-Janssen DJ, Muizert Y, Verweij C, Discipio RG, Daha MR: Characterization of complement C6 deficiency in a PVG/c rat strain. *Clin Exp Immunol* 1997, 109:387-396
6. Brandt J, Pippin J, Schulze M, Hansch GM, Alpers CE, Johnson RJ, Gordon K, Couser WG: Role of the complement membrane attack complex (C5b-9) in mediating experimental mesangioproliferative glomerulonephritis. *Kidney Int* 1996, 49:335-343
7. Floege J, Alpers CE, Burns MW, Pritzl P, Gordon K, Couser WG, Johnson RJ: Glomerular cells, extracellular matrix accumulation, and the development of glomerulosclerosis in the remnant kidney model. *Lab Invest* 1992, 66:485-497
8. Floege J, Eng E, Young BA, Johnson RJ: Factors involved in the regulation of mesangial cell proliferation *in vitro* and *in vivo*. *Kidney Int* 1993, 39 (Suppl.): S47-54
9. Rehan A, Johnson KJ, Wiggins RC, Kunkel RG, Ward PA: Evidence for the role of oxygen radicals in acute nephrotoxic nephritis. *Lab Invest* 1984, 51:396-403
10. Yamamoto T, Wilson CB: Quantitative and qualitative studies of antibody-induced mesangial cell damage in the rat. *Kidney Int* 1987, 32:514-525
11. Bagchus WM, Hoedemaeker PJ, Rozing J, Bakker WW: Glomerulonephritis induced by monoclonal anti-Thy 1.1 antibodies: a sequential histological and ultrastructural study in the rat. *Lab Invest* 1986, 55:680-687
12. Rehan A, Wiggins RC, Kunkel RG, Till GO, Johnson KJ: Glomerular injury and proteinuria in rats after intrarenal injection of cobra venom factor: evidence for the role of neutrophil-derived oxygen free radicals. *Am J Pathol* 1986, 123:57-66
13. de Haan P, de Jonge AJ, Verbrugge T, Boersma DM: Three epitope-specific monoclonal antibodies against the hapten penicillin. *Int Arch Allergy Appl Immunol* 1985, 76:42-46
14. Ferrario F, Castiglione A, Colasanti G, Barbiano DB, Bertoli S, D'Amico G: The detection of monocytes in human glomerulonephritis. *Kidney Int* 1985, 28:513-519
15. Wilson CB, Dixon FJ: The renal response to immunological injury. *The Kidney*. Edited by BM Brenner, FC Rector, Jr. Philadelphia, W.B. Saunders, 1981, pp 1237-1350
16. Boyce NW, Holdsworth SR: Hydroxyl radical mediation of immune renal injury by desferrioxamine. *Kidney Int* 1986, 30:813-817
17. Brady HR: Leukocyte adhesion molecules and kidney diseases. *Kidney Int* 1994, 45:1285-1300
18. Rehan A, Johnson KJ, Kunkel RG, Wiggins RC: Role of oxygen radicals in phorbol myristate acetate-induced glomerular injury. *Kidney Int* 1985, 27:503-511
19. Shah SV: Role of reactive oxygen metabolites in experimental glomerular disease. *Kidney Int* 1989, 35:1093-1106
20. Horgan MJ, Wright SD, Malik AB: Antibody against leukocyte integrin (CD18) prevents reperfusion-induced lung vascular injury. *Am J Physiol* 1990, 259:L315-L319
21. Huitinga I, Damoiseaux JG, Dopp EA, Dijkstra CD: Treatment with anti-CD18 antibodies ED7 and ED8 suppresses experimental allergic encephalomyelitis in Lewis rats. *Eur J Immunol* 1993, 23:709-715
22. Palmen MJ, Dijkstra CD, van der Ende MB, Pena AS, van Rees EP: Anti-CD11b/CD18 antibodies reduce inflammation in acute colitis in rats. *Clin Exp Immunol* 1995, 101:351-356
23. van Goor H, van der Horst ML, Fidler V, Grond J: Glomerular macrophage modulation affects mesangial expansion in the rat after renal ablation. *Lab Invest* 1992, 66:564-571
24. Monga G, Mazzucco G, di Belgiojoso GB, Busnach G: The presence and possible role of monocyte infiltration in human chronic proliferative glomerulonephritides: light microscopic, immunofluorescence, and histochemical correlations. *Am J Pathol* 1979, 94:271-284
25. Hooke DH, Hancock WW, Gee DC, Kraft N, Atkins RC: Monoclonal

- antibody analysis of glomerular hypercellularity in human glomerulonephritis. *Clin Nephrol* 1984, 22:163–168
26. Cook HT, Smith J, Salmon JA, Cattell V: Functional characteristics of macrophages in glomerulonephritis in the rat: O2-generation, MHC class II expression, and eicosanoid synthesis. *Am J Pathol* 1989, 134:431–437
27. Hara M, Batsford SR, Mihatsch MJ, Bitter-Suermann D, Vogt A: Complement and monocytes are essential for provoking glomerular injury in passive Heymann nephritis in rats: terminal complement components are not the sole mediators of proteinuria. *Lab Invest* 1991, 65:168–179
28. Schreiner GF, Cotran RS, Unanue ER: Modulation of Ia and leukocyte common antigen expression in rat glomeruli during the course of glomerulonephritis and aminonucleoside nephrosis. *Lab Invest* 1984, 51:524–533
29. Cattell V, Largen P, De Heer E, Cook T: Glomeruli synthesize nitrite in active Heymann nephritis: the source is infiltrating macrophages. *Kidney Int* 1991, 40:847–851
30. Magil AB, Cohen AH: Monocytes and focal glomerulosclerosis. *Lab Invest* 1989, 61:404–409
31. Matsumoto K, Atkins RC: Glomerular cells and macrophages in the progression of experimental focal and segmental glomerulosclerosis. *Am J Pathol* 1989, 134:933–945
32. Floege J, Burns MW, Alpers CE, Yoshimura A, Pritzl P, Gordon K, Seifert RA, Bowen-Pope DF, Couser WG, Johnson RJ: Glomerular cell proliferation and PDGF expression precede glomerulosclerosis in the remnant kidney model. *Kidney Int* 1992, 41:297–309
33. Bagchus WM, Jeunink MF, Rozing J, Elema JD: A monoclonal antibody against rat platelets. I. Tissue distribution in vitro and in vivo. *Clin Exp Immunol* 1989, 75:317–323
34. Dijkstra CD, Dopp EA, Joling P, Kraal G: The heterogeneity of mononuclear phagocytes in lymphoid organs: distinct macrophage subpopulations in rat recognized by monoclonal antibodies ED1, ED2 and ED3. *Adv Exp Med Biol* 1985, 186:409–419
35. Skalli O, Ropraz P, Trzeciak A, Benzouana G, Gillesen D, Gabbiani G: A monoclonal antibody against alpha-smooth muscle actin: a new probe for smooth muscle differentiation. *J Cell Biol* 1986, 103:2787–2796
36. Williams AF, Galfre G, Milstein C: Analysis of cell surfaces by xenogeneic myeloma-hybrid antibodies: differentiation antigens of rat lymphocytes. *Cell* 1977, 12:663–673
37. Brown WR, Barclay AN, Sunderland CA, Williams AF: Identification of a glycoprotein-like molecule at the cell surface of rat thymocytes. *Nature* 1981, 289:456–460
38. Killeen N, Barclay AN, Willis AC, Williams AF: The sequence of rat leukosialin (W3/13 antigen) reveals a molecule with O-linked glycosylation of one third of its extracellular amino acids. *EMBO J* 1987, 6:4029–4034
39. Lopez S, Seveau S, Lesavre P, Robinson MK, Halbwachs-Mecarelli L: CD43 (sialophorin, leukosialin) shedding is an initial event during neutrophil migration, which could be closely related to the spreading of adherent cells. *Cell Adhes Commun* 1998, 5:151–160
40. Seveau S, Lopez S, Lesavre P, Guichard J, Cramer EM, Halbwachs-Mecarelli L: Leukosialin (CD43, sialophorin) redistribution in uropods of polarized neutrophils is induced by CD43 cross-linking by antibodies, by colchicine or by chemotactic peptides. *J Cell Sci* 1997, 110:1465–1475
41. Damoiseaux JG, Dopp EA, Neefjes JJ, Beelen RH, Dijkstra CD: Heterogeneity of macrophages in the rat evidenced by variability in determinants: two new anti-rat macrophage antibodies against a heterodimer of 160 and 95 kd (CD11/CD18). *J Leukoc Biol* 1989, 46:556–564
42. Floege J, Johnson RJ, Gordon K, Iida H, Pritzl P, Yoshimura A, Campbell C, Alpers CE, Couser WG: Increased synthesis of extracellular matrix in mesangial proliferative nephritis. *Kidney Int* 1991, 40:477–488
43. Harlan JM, Liu DY: Adhesion: Its Role in Inflammatory Disease. New York, W.H. Freeman, 1992
44. Kaslovsky RA, Horgan MJ, Lum H, McCandless BK, Gilboa N, Wright SD, Malik AB: Pulmonary edema induced by phagocytosing neutrophils: protective effect of monoclonal antibody against phagocyte CD18 integrin. *Circ Res* 1990, 67:795–802
45. Mulligan MS, Johnson KJ, Todd RF, Issekutz TB, Miyasaka M, Tamatani T, Smith CW, Anderson DC, Ward PA: Requirements for leukocyte adhesion molecules in nephrotoxic nephritis. *J Clin Invest* 1993, 91:577–587
46. Wu X, Pippin J, Lefkowitz JB: Attenuation of immune-mediated glomerulonephritis with an anti-CD11b monoclonal antibody. *Am J Physiol* 1993, 264:F715–F721
47. Suter S, Nydegger UE, Roux L, Waldvogel FA: Cleavage of C3 by neutral proteases from granulocytes in pleural empyema. *J Infect Dis* 1981, 144:499–508
48. Gaither TA, Hammer CH, Gadek JE, Katusha K, Santaella M, Frank MM: Cleavage of membrane-bound C3b and C3bi by viable human neutrophils (PMN). *Mol Immunol* 1983, 20:623–635
49. Carlo JR, Spitznagel JK, Studer EJ, Conrad DH, Ruddy S: Cleavage of membrane bound C3bi, an intermediate of the third component of complement, to C3c and C3d-like fragments by crude leucocyte lysosomal lysates and purified leucocyte elastase. *Immunology* 1981, 44:381–391
50. Leid RW, Ballieux BE, van dH I, van Kleyburg DK, Hagen EC, van Es LA, van der Woude FJ, Daha MR: Cleavage and inactivation of human C1 inhibitor by the human leukocyte proteinase, proteinase 3. *Eur J Immunol* 1993, 23:2939–2944
51. De Heer E, Prodjosudjadi W, Davidoff A, van der Wal A, Bruijn JA, Paul LC: Control of monocyte influx in glomerulonephritis in transplanted kidneys in the rat. *Lab Invest* 1998, 78:1327–1337
52. Isaka Y, Fujiwara Y, Ueda N, Kaneda Y, Kamada T, Imai E: Glomerulosclerosis induced by in vivo transfection of transforming growth factor-beta or platelet-derived growth factor gene into the rat kidney. *J Clin Invest* 1993, 92:2597–2601
53. Mattana J, Singhal PC: Macrophage supernatants have both stimulatory and suppressive effects on mesangial cell proliferation. *J Cell Physiol* 1993, 154:289–293
54. Wahl SM, McCartney-Francis N, Mergenhagen SE: Inflammatory and immunomodulatory roles of TGF-beta. *Immunol Today* 1989, 10:258–261
55. Border WA, Okuda S, Languino LR, Sporn MB, Ruoslahti E: Suppression of experimental glomerulonephritis by antiserum against transforming growth factor beta 1. *Nature* 1990, 346:371–374
56. Border WA, Okuda S, Languino LR, Ruoslahti E: Transforming growth factor-beta regulates production of proteoglycans by mesangial cells. *Kidney Int* 1990, 37:689–695
57. Cattell V, Lianos E, Largen P, Cook T: Glomerular NO synthase activity in mesangial cell immune injury. *Exp Nephrol* 1993, 1:36–40
58. Ding G, Pesek-Diamond I, Diamond JR: Cholesterol, macrophages, and gene expression of TGF-beta 1 and fibronectin during nephrosis. *Am J Physiol* 1993, 264:F577–F584
59. Mosquera JA: Increased production of fibronectin by glomerular cultures from rats with nephrotoxic nephritis: macrophages induce fibronectin production in cultured mesangial cells. *Lab Invest* 1993, 68:406–412