Reith lectures to see that the practice of psychiatry now confers a title to make authoritative pronouncements on a range of social problems that extends from adolescent unchastity to world government; and for millions of readers throughout the world Dr. Spock continues to resolve, in effortless prose, the fundamental problems of human relationships. To-day Freud and Marx might fairly compete for the title to the biggest popular success since Jesus Christ; and Freud, I think, would win.

The explanation of this phenomenon is not, in my judgment, far to seek. The medical profession has, I think—and here I imply no undervaluation of the skill and devotion of its practitioners—simply inherited the mantle of the church in an age in which profound religious conviction is exceptional and intimations of immortality are seldom taken seriously. In spite of occasional abuses of the National Health Service by disgruntled patients, and a steady trickle of actions against hospitals for negligence, the contemporary prestige of the doctor stands extremely high. And naturally so. For if this life is all we have, it is inevitable that we should look with new reverence upon the man whose hands alone can temporarily bar the gates of death. Equally it is reasonable, as the supernatural sanction of moral systems fades, to seek an identification of morality and mental health. As a personal and social goal health has much to commend it. It is in harmony with science, and science is in harmony with the age. Besides, failure to achieve the goal of health evokes pity and sympathy: failure to attain virtue merely provokes censure. Goodness, too, is often priggish: health never. So if we once owed it to God to be good, we now owe it to science to be healthy.

To-day splendid possibilities, certainly the prospect of splendid new empires, are displayed before the medical world. Indeed, we may well be on the brink of an age in which the power of science to influence human behaviour

will achieve a new dimension. Yet the question, Whose behaviour is to be influenced, and in what direction? remains, and will remain, as obstinate as ever. It can be solved, not by any one profession alone, but only by the good sense, the generosity, and the tolerance of the whole community. Meanwhile let us not forget that all our misfits, failures, and deviants—from the Committee of 100 to the utterly feeble and feckless—are in their several ways attempting to register protests against the demands or the values of the world in which they have to live. To treat those protests as symptomatic of mental disorder is a subtle device for drawing their sting—and one which the merits of the society against which those protests are directed cannot always be said to justify.

REFERENCES

- ¹ Szasz, Thomas S., *Psychiatric Expert Testimony—Its Covert Meaning and Social Function*. *Psychiatry*, 1957, 20, 313.
- ² — Ibid.
- ³ Wootton, Barbara. *Diminished Responsibility: A Layman's View*. *Law Quarterly Review*, Vol. 76, No. 302, April, 1960.
- ⁴ Unpublished transcript of evidence in the case named.
- ⁵ [1960] 44 Cr App. R. 246.
- ⁶ Snell, H. K., Director of Medical Services, Prison Commission, H.M. Prison, Grendon. *Brit. med. J.*, 1962, 2, 789.
- ⁷ [1962] *Criminal Law Review* 647.
- ⁸ *The Times Law Report*, August 15, 1961.
- ⁹ Snell, H. K., *Op cit.*
- ¹⁰ *Depression and Suicide*. Mental Health Research Fund Pamphlet No. 2.
- ¹¹ Leading article in the *Nursing Times*, summarized in *The Times*, April 5, 1963.
- ¹² House of Lords Debates, May 14, 1963, col. 1174.
- ¹³ Eatherly, Claude, and Anders, Gunther, *Burning Conscience*, 1962. Weidenfeld and Nicolson, London.
- ¹⁴ House of Lords Debates, December 19, 1961, col. 651.
- ¹⁵ Jones, Maxwell, *The Treatment of Character Disorders*, *Brit. J. Crim.*, 1963, 3, 267.
- ¹⁶ *The Times*, May 19, 1961.
- ¹⁷ Hart, H. L. A., *Punishment and the Elimination of Responsibility*, 1962. Athlone Press, London.
- ¹⁸ Szasz, Thomas S., *Moral Conflict and Psychiatry*, *Yale Review*, Summer, 1960.
- ¹⁹ Quoted by *The Observer*, February 3, 1963, from Benjamin Spock's *Problems of Parents*, 1963. Bodley Head, London.

INACTIVATION OF BENZYL PENICILLIN AND METHICILLIN BY HOSPITAL STAPHYLOCOCCI

BY

G. A. J. AYLIFFE, M.B., Ch.B., B.Sc.

MARY BARBER, M.D.

Department of Bacteriology, Postgraduate Medical School of London

Since the penicillin-resistant strains of *Staphylococcus aureus* isolated in clinical practice owe their resistance to the production of the enzyme penicillinase and remain sensitive to the unchanged antibiotic, their sensitivity *in vitro* varies many hundredfold with the size of the inoculum. This means that a figure for the minimum inhibitory concentration of penicillin for these strains, based on a single estimation of sensitivity, is misleading, and in most routine clinical laboratories the organisms are simply reported as penicillin-resistant.

Nevertheless the rate of production of penicillinase and the degree to which it is inducible by penicillin vary very much with different strains, and quantitative estimation of this is of practical value for two reasons. In the first place infections due to strains with weak penicillinase production may respond to benzylpenicillin alone, and, more important, it is only against strains of this type that there is an apparent synergy between benzylpenicillin and fucidin.

Secondly, this has an important bearing on the recent discussion in the *British Medical Journal* with regard to

methicillin inactivation by methicillin-resistant staphylococci. Methicillin is in fact slowly hydrolysed by staphylococcal penicillinase, so that with highly active penicillinase-producing strains detectable inactivation of methicillin occurs in 6 to 24 hours, whereas with weak penicillinase-producers this is not the case. Thus in comparing methicillin-inactivation by a methicillin-sensitive and a methicillin-resistant staphylococcus it is important to compare their activity against benzylpenicillin at the same time, and without such tests it cannot be assumed that methicillin-resistance in itself implies an increased capacity to inactivate methicillin.

A simple quantitative test of penicillinase activity would therefore be of great value. In the present investigation 276 penicillin-resistant strains of *Staph. aureus*, including 37 methicillin-resistant strains, have been graded with regard to penicillinase activity on the basis of the rapid screening test of Foley and Perret (1962), and the reliability of this grading has been assessed by iodometric estimations of penicillinase activity in a proportion of the strains.

Material and Methods

Source of Strains.—All strains tested were isolated from clinical material. Most of them were recently isolated in the routine bacteriological department of a single large London hospital. Nine (7 of which were methicillin-resistant) came from other laboratories in England (Messenger, Q.M.2, N.I.524, and three others), Paris (P6), and Copenhagen (Ca and Cb).

Screening Test for Penicillinase Activity.—This was based on the method described by Foley and Perret (1962). Single colonies from 18-hour cultures on unenriched lab-lemco agar plates were emulsified in 0.2 ml. of a freshly prepared solution of 10,000 units of penicillin/ml. in 0.05 M phosphate buffer at pH 6.5, and the mixtures left at room temperature for one hour. One drop (0.02 ml.) of a freshly prepared mixture of equal parts 1% starch solution and 0.04 N iodine was then added and the time of disappearance of the blue colour noted. Control cultures of a penicillin-sensitive *Staph. aureus* and a weak penicillinase-producing strain were put up with every series of tests. The penicillin solution had to be prepared freshly each day, as minor degrees of hydrolysis gave false-positive results. It was important for comparative purposes to standardize not only the size of colony used but also the culture medium and the time of incubation. It was found, for example, that colonies taken from blood-agar plates caused much more rapid hydrolysis than did those from plain lab-lemco agar, even if the former were no larger than the latter.

Iodometric Assay.—With a proportion of strains, estimations of penicillinase activity were made on six-hour shaken cultures in nutrient broth with or without 0.25 µg. of methicillin/ml., by the iodometric method described by Perret (1954).

Methicillin Inactivation.—Tests for methicillin inactivation were carried out by two methods. In method 1 broth containing 20 or 50 µg./ml. was inoculated with one drop of an 18- to 24-hour plain broth culture, and shaken at 37° C. for 24 hours. In method 2 the strains were shaken for six hours in broth containing 0.25 µg. of methicillin/ml., and then 2 ml. of these pre-induced cultures was mixed with 2 ml. of broth containing 40 or 100 µg. of methicillin/ml. (to give a final concentration of 20 and 50 µg./ml.); the mixtures were then incubated at 37° C., without shaking, for a further 24 hours. In both methods the remaining methicillin was assayed by the cup-plate method. The plates were seeded with the Oxford staphylococcus and after the antibiotic-containing mixtures had been added to the cups the plates were kept at 4° C. for two hours to allow prediffusion before incubating at 37° C. Controls consisting of uninoculated broth containing methicillin were used in all experiments.

Results

Screening Test for Penicillinase Activity

A total of 432 strains of *Staph. aureus* were tested. Of these, 156 were penicillin-sensitive and all gave negative results. The other 276 strains were all penicillin-resistant, and all but 21 decolorized the starch-iodine mixture in 60 minutes or less. The positive results were graded as follows:

Decolorization within 5 minutes ..	++
Decolorization within 5-30 minutes ..	+
Decolorization within 30-60 minutes ..	±

Table I analyses the results according to phage group and whether the strain was resistant to penicillin only or

resistant to penicillin and one or more other antibiotic. It will be seen that 56% of the multiple-resistant strains were highly active compared with only 34% of the strains resistant to penicillin only. Among the typable strains

TABLE I.—Results of Screening Test for Penicillinase Activity

Phage Group	Strains Resistant to Penicillin only (132)				Multiple-resistant Strains (144)			
	++	+	±	—	++	+	±	—
I ..	25	20	8	12	28	17	4	0
II ..	5	1	4	1	1	1	1	0
III ..	8	7	5	0	44 (23)	20 (11)	4 (3)	0
Miscellaneous	5	6	4	3	0	0	0	0
Non-typable	2	6	8	2	7	3	11	3
Total ..	45	40	29	18	80	41	20	3

Figures in parentheses represent methicillin-resistant strains.

there was no significant trend within the different phage groups, except for the fact that a high proportion of the multiple-resistant strains were phage group III. Among the latter were 37 methicillin-resistant strains, and the proportion of these which gave a ++ reaction was just over 60%. Of 120 phage-typable multiple-resistant strains, 73 (61%) gave strong positive reactions, whereas only nine gave weak reactions and none was negative. There was a significantly higher incidence of weak penicillinase-producers among the non-typable multiple-resistant strains and the strains resistant to penicillin only.

Repeat Tests.—With 74 strains the screening test was repeated on a second occasion. Minor variations in the results obtained each time (that is, from ++ to + or ± to —) were frequent, but a gross variation was noted with only three strains.

Iodometric Assay

Iodometric assay of penicillinase activity was carried out with 17 strains; with the screening test eight of these gave a ++ reaction, one a + reaction, and eight ± or —. The results are given in Table II. It will be seen that there is general agreement with the screening test. It will also be seen that there is considerable variation in activity in repeat tests on the same strains. This was probably due

TABLE II.—Results of Iodometric Assay

Strain	Result of Screening Test	Penicillinase Activity (µmol Penicilloic Acid ml. of Broth Culture hr.)			
		A. Non-induced	B. Induced		
			Test 1	Test 2	Test 3
H.1 ..	++	12.6	216		
H.18 ..	++	2.6	66		
H.30 ..	++	Not tested	168		
H.31 ..	++	9.6	110		
H.217 ..	++	12.8	64		214
Q.M.2 ..	++	6.2	60	158	
Ca ..	++	16.2	212		
N.I.524 ..	++	6.2	200	160	118
H.54 ..	+	1.6	78		
Messenger ..	±	1.8	18	13	
H.9 ..	±	3.6	34		
H.16 ..	±	3.6	31		
H.19 ..	—	Not tested	17		
H.22 ..	—	1.8	13	15	29
H.25 ..	—	7.6	31		
H.28 ..	—	1.6	46		
H.33 ..	—	14.8	48	31	14

A, 6-hour shaken cultures in digest broth.

B, 6-hour shaken cultures in digest broth plus 0.25 µg. methicillin/ml.

mainly to variations in degree and stage of growth of the cultures tested. The number of organisms present in the cultures, as measured by opacity, ranged from approximately 2×10^9 /ml. to 4×10^9 /ml. The two strains showing the most variable results, H.217 and Q.M.2, are both methicillin-resistant strains, and these tend to be very variable in cultural characteristics.

Inactivation of Methicillin

Methicillin inactivation was studied with three methicillin strains, from London (Q.M.2 and H.217) and Copenhagen (Ca), all of which were highly active penicillinase-producers; and with two methicillin-sensitive strains, one active penicillinase-producer (N.I.524) and one weak penicillinase-producer (Messenger). By the first method, in which one drop of a culture, shaken overnight, was added to broth containing 50 or 20 µg. of methicillin/ml., inactivation was related to growth. Thus the methicillin-sensitive strain, N.I.524, failed to grow, and caused no inactivation (Table III). On the other hand, the three methicillin-resistant strains grew well in the presence of 20 µg./ml. and inactivated most of it, whereas in the presence of 50 µg./ml. they grew slightly and caused only slight inactivation.

TABLE III.—*Inactivation of Methicillin*
Method 1

Organisms	Methicillin Sensitivity	Broth containing			
		50 µg. Methicillin ml.		20 µg. Methicillin/ml.	
		Diameter of Inhibition Zone (mm.)	Growth	Diameter of Inhibition Zone (mm.)	Growth
Q.M.2 ..	R	15	+	< 8	+++
H.217 ..	R	18	±	< 8	+++
Ca ..	R	18	±	< 8	+++
N.I.524 ..	S	20	—	14	—
Control ..		20	—	15	—

Method 2: Pre-induced Cultures

Organisms	Methicillin Sensitivity	Penicillinase Activity	Diameter of Inhibition Zone (mm.)	
			50 µg. Methicillin/ml.	20 µg. Methicillin/ml.
Q.M.2 ..	R	++	22	10
H.217 ..	R	++	21	10
Ca ..	R	++	19	< 8
N.I.524 ..	S	++	20	10
Messenger ..	S	±	22	17
Control ..	—	—	23	16

In method 2, 2 ml. of an overnight culture grown in broth containing 0.25 µg. of methicillin/ml. was added to an equal quantity of broth containing methicillin. In this test all the active penicillinase producers, whether methicillin-resistant or not, inactivated a considerable portion of the methicillin in the broth containing 20 µg./ml., whereas the weak penicillinase producer caused no inactivation. With the higher concentration little or no detectable inactivation occurred with any of the strains by this method.

Isoxazolyl Penicillins.—Similar comparative results with methicillin-sensitive and methicillin-resistant staphylococci were obtained with oxacillin and cloxacillin except that since they are less resistant to staphylococcal penicillinase they were more readily inactivated by all strains tested.

Discussion

The screening test for penicillinase activity described in this paper can only be regarded as a rough guide, but it does differentiate between highly active and poor penicillinase-producing strains of *Staph. aureus*. Of the 276 penicillin-resistant strains tested, 49 gave a very slow reaction and 21 were completely negative, although they were in fact penicillinase-producing strains. Foley and Perret (1962) got a rapid reaction with all penicillinase-producing strains tested. This discrepancy is possibly because they took colonies from blood-agar plates, whereas we used unenriched lab-lemco agar.

Since it can be shown that the penicillin-resistant staphylococci isolated in clinical practice owe their resistance almost solely to penicillinase-production, the same information could probably be obtained by direct sensitivity tests with a carefully controlled inoculum. In this laboratory routine sensitivity tests for staphylococci are carried out by streaking single colonies across a ditch-plate with 10 units of benzylpenicillin/ml. in the ditch. The Oxford staphylococcus is streaked across the centre of each plate. If the test strain grows to the edge of the ditch or across it it is recorded as resistant, and if it grows nearer to the ditch than the Oxford staphylococcus, but does not reach it, it is recorded as partially resistant. Nearly all the weak penicillinase-producing strains in this investigation were in fact recorded as partially resistant.

In the present investigation of 276 penicillin-resistant strains, 70 (25%) gave weak or negative reactions with the screening test used. Out of 120 multiple-resistant strains, which were typable by bacteriophage, only 9 (8%) gave weak reactions and none was negative. The typable multiple-resistant strains represent the hospital staphylococci and are responsible for nearly all cases of cross-infection in the surgical wards. They have probably, therefore, been in contact with penicillin for many years.

The strains showing weak penicillinase production were all inducible to some degree, in the sense that they all showed some increase in penicillinase activity after growth in low concentrations of methicillin. Whether or not prolonged passage in penicillin or methicillin would render them active penicillinase-producers remains to be seen, but this effect does not occur with a few passages. It is infection with this type of staphylococcus which is likely to respond to large doses of benzylpenicillin and for which benzylpenicillin and fucidin would be synergic.

As shown by Pollock (1962) and Novick (1962) methicillin is not completely resistant to staphylococcal penicillinase, although the rate of hydrolysis (V_{max}) is 30 times lower than with benzylpenicillin and, moreover, the affinity of the enzyme for methicillin is extremely low. The two factors together mean that the half-life of an antibiologically effective concentration of methicillin in the presence of staphylococcal penicillinase is 360,000 times longer than that of an antibiologically equivalent concentration of benzylpenicillin (Novick, 1962). In the present investigation it has been shown that active penicillinase-producing strains of *Staph. aureus* cause a similar order of methicillin inactivation whether they are methicillin-sensitive or methicillin-resistant, provided the test used does not involve growth in the presence of the antibiotic during the period of inactivation. If, however, tests are carried out as described in method 1, in such a way that penicillinase is produced only by the test strain if it can multiply in the presence of methicillin, then, of course, methicillin-resistant strains will be able to multiply in and inactivate a concentration of methicillin which will be totally inhibitory to the methicillin-sensitive organism. This is the type of test used by Eriksen and Erichsen (1963), and clearly it does not imply a changed penicillinase activity on the part of the methicillin-resistant strain. Moreover, from a practical point of view the degree of methicillin inactivation is infinitesimal compared with the inactivation of benzylpenicillin by these strains.

Finally, in view of the recent discussion on the subject in the *British Medical Journal* it is perhaps worth saying a word on the virulence of naturally occurring methicillin-resistant strains of *Staph. aureus*. The 37 strains tested in this investigation all shared the properties described by

other investigators (Jevons, 1961; Knox and Smith, 1961; Barber, 1961, 1962; Eriksen and Erichsen, 1963; Stewart and Holt, 1963; Rolinson, 1963). It is possible, therefore, that infections with these strains would respond to treatment with large doses of methicillin. In the presence of quite low concentrations of methicillin (5–10 µg./ml.) they grow poorly; not all the cells in a given inoculum grow, and on solid medium these tend to be massed together at the site of heavy inoculum, and individual colonies if present tend to be small and atypical. However, in the absence of methicillin these strains appear to have all the characteristics of fully virulent *Staph. aureus in vitro*. Moreover, the strains isolated in this hospital, like those of Stewart and Holt (1963), were able to cause cross-infection in the surgical wards, and in some instances the infection was of a severe generalized nature.

Summary

A simple screening test for penicillinase activity has been used to investigate 432 strains of *Staph. aureus* and its reliability assessed by iodometric estimations with a proportion of strains.

None of 156 penicillin-sensitive strains showed a positive reaction.

The results obtained with 276 penicillin-resistant strains were graded according to the rate of reaction; 70 (25%) were found to be weak penicillinase-producers, and 125 (45%) were highly active. Out of 120 phage-typable

multiple-resistant strains 73 (61%) were strongly positive, only 9 gave weak reactions, and none was negative. On the other hand, of 132 strains resistant to penicillin alone, whether phage-typable or not, only 45 (34%) were strongly positive and 47 gave weak or negative reactions.

Methicillin inactivation by methicillin-sensitive and methicillin-resistant strains has been studied, and it has been shown that the degree of inactivation is related to penicillinase activity rather than methicillin-resistance, unless tests are carried out in such a way that inactivation is dependent on growth in the presence of methicillin.

The clinical application of these findings is discussed.

Our thanks are due to Mrs. June Filbey for technical assistance. We are also indebted to Drs. Y. Chabbert, Nuala Crowley, K. R. Eriksen, Patricia Jevons, R. W. Riddell, and G. T. Stewart for sending us cultures of methicillin-resistant strains and to Dr. M. R. Pollock for strain N.I.524.

REFERENCES

- Barber, M. (1961). *J. clin. Path.*, **14**, 385.
 — (1962). In *Resistance of Bacteria to the Penicillins*. Ciba Foundation Study Group, No. 13, p. 89. Churchill, London.
 Eriksen, K. R., and Erichsen, I. (1963). *Brit. med. J.*, **1**, 746.
 Foley, J. M., and Perret, C. J. (1962). *Nature (Lond.)*, **195**, 287.
 Knox, R., and Smith, J. T. (1961). *Lancet*, **2**, 520.
 Jevons, M. P. (1961). *Brit. med. J.*, **1**, 124.
 Novick, R. P. (1962). *Biochem. J.*, **83**, 229.
 Pollock, M. R. (1962). In *Resistance of Bacteria to the Penicillins*. Ciba Foundation Study Group, No. 13, p. 56. Churchill, London.
 Perret, C. J. (1954). *Nature (Lond.)*, **174**, 1012.
 Rolinson, G. N. (1963). *Brit. med. J.*, **1**, 542.
 Stewart, G. T., and Holt, R. J. (1963). *Ibid.*, **1**, 308.

STABILITY OF METHICILLIN AND CLOXACILLIN TO STAPHYLOCOCCAL PENICILLINASE

BY

R. KNOX, M.D., F.R.C.P.

Professor of Bacteriology

Guy's Hospital Medical School, London

J. T. SMITH, Ph.D.

Lecturer in Bacteriology

Penicillinase-producing staphylococci are resistant to benzylpenicillin only because large inocula can destroy the drug before it destroys them; small inocula are almost as sensitive to benzylpenicillin as fully sensitive strains which do not possess penicillinase. This effect of inoculum size is characteristic of penicillinase-producing strains. However, a similar effect can also be observed in populations of bacteria which have a wide range of susceptibility to an antibiotic. Thus the effect of inoculum size must not be taken as *prima facie* evidence for the penicillinase type of resistance. For example, the 13137 strain of methicillin-resistant staphylococcus (Jevons, 1961; Knox, 1961; Rolinson, 1961) has been found to contain many organisms which were of the same order of sensitivity as methicillin-sensitive strains of *Staphylococcus aureus*, together with a minority of rather highly resistant organisms (Knox and Smith, 1961).

The resistance of this and other strains to methicillin has been ascribed not to destruction of methicillin but to a mechanism similar to the inherent resistance of penicillinase-negative staphylococci trained towards benzylpenicillin resistance *in vitro*. It has also been suggested that resistance to the newer isoxazolyl penicillins (usually associated with cross-resistance to methicillin) is similarly of the "inherent" type (Knox and Smith, 1963; Ayliffe, Barber, and Waterworth, 1963; Rolinson, 1963). On the other hand, there have also been suggestions that destruction of these penicillins partly accounts for the resistance of such strains, and it has been implied, though not explicitly

stated, that they possess some special ability to destroy methicillin or the isoxazolyl penicillins not possessed by other penicillinase-producing staphylococci (Stewart and Holt, 1963a, 1963b, 1963c; Eriksen and Erichsen, 1963). This communication shows that the destruction of these penicillins is due to penicillinase, whether the strains are sensitive or resistant to methicillin and the isoxazolyl penicillins.

Materials and Methods

Microbiological estimation of the penicillins was made by the cup-plate assay method with *Sarcina lutea* ATCC 9341 as test organism (Heatley, 1944).

Initial rates of destruction of methicillin, cloxacillin, and benzylpenicillin were estimated by the hydroxylamine method at an initial penicillin concentration of 2 mg./ml. in the presence of 10^{-3} M *p*-chloromercuribenzoate as described by Knox and Smith (1962).

Sensitivity tests were carried out by preparing inocula from overnight broth cultures by dilution in broth, and 0.02-ml. amounts were dropped on to the surface of agar plates containing penicillin. The results were read after nine hours' incubation.

The organisms used were penicillinase-producing coagulase-positive strains of *Staph. aureus*. Strain E3 has been described by Knox (1960) and is methicillin- and cloxacillin-sensitive. The following methicillin- and cloxacillin-resistant strains were used: 13137/1000 (Knox and Smith, 1961); 6367, which was kindly supplied by