It is clear that he had nothing to gain, and perhaps much to lose, by our paying more than passing attention to this aspect of his case. So far as is known, he was fairly well adjusted to the male sex, but that is not to say that he may not have suffered acutely throughout his life by possessing such a deformity. A genital abnormality of this nature

Fig. 3.—Section of the ovarian part of the gland showing an old corpus albicans. ($\times 75$.)

should nowadays call for full investigation within the first year of life; once such investigation is complete, a decision may be made concerning the sex in which it is better to bring the patient up and appropriate plastic surgical procedures can be planned (Wilkins, 1960; Gordon and Dewhurst, 1962).

Summary

The case is described of a 40-year-old hermaphrodite whose social sex was male. The external genitalia showed a moderate-sized penis, a perineal urethra, and a rudimentary vagina; within the sac of a right inguinal hernia were a single ovotestis, a uterus, and a Fallopian tube. Chromosome analysis showed him to have 46 chromosomes with an XX arrangement of sex chromosomes.

We wish to record our grateful thanks to Mr. Francis Masina, under whose care the patient was admitted and who performed the operation, for permission to study and to report this case.

REFERENCES

- de Assis, L. M., Epps, D. R., and Bottura, C. (1960). *Lancet*, **2**, 129.
 Ferguson-Smith, M. A., Johnston, A. W., and Weinberg, A. N. (1960). *Ibid.*, **2**, 126.
 Gordon, R. R., and Dewhurst, C. J. (1962). *Ibid.*, **2**, 872.
 ———, O'Gorman, F. J. P., Dewhurst, C. J., and Blank, C. E. (1960). *Ibid.*, **2**, 736.
 Grumbach, M. M., and Barr, M. L. (1958). *Recent Progr. Hormone Res.*, **14**, 255.
 Harnden, D. G. (1960). *Brit. J. exp. Path.*, **41**, 31.
 ——— and Armstrong, C. N. (1959). *Brit. med. J.*, **2**, 1287.
 Hirschorn, K., Decker, W. H., and Cooper, H. L. (1960). *New Engl. J. Med.*, **263**, 1044.
 Hungerford, D. A., Donnelley, A. J., Nowell, P. C., and Beck, S. (1959). *Amer. J. hum. Genet.*, **11**, 215.
 Miller, O. J. (1961). Quoted by S. Rappoport, and W. D. Kaplan, *J. Pediat.*, **1961**, **59**, 415.
 Moorhead, P. S., Nowell, P. C., Mellman, W. J., Battips, D. M., and Hungerford, D. A. (1960). *Exp. Cell Res.*, **20**, 613.
 Sandberg, A. A., Koepf, G. F., Crosswhite, L. H., and Hauschka, T. S. (1960). *Amer. J. hum. Genet.*, **12**, 231.
 Sasaki, M., and Makino, S. (1960). *Texas Rep. Biol. Med.*, **18**, 493.
 Waxman, S. H., Kelley, V. C., Gartler, S. M., and Burt, B. (1962). *Lancet*, **1**, 161.
 Wilkins, L. (1960). *Pediatrics*, **26**, 846.

Medical Memoranda

Heal Prolapse Through Ruptured Vagina : Complication of Manchester Operation

The following case is thought to be rare enough to warrant recording.

CASE REPORT

The patient, a para-8 aged 79, was first operated on in September, 1958, for a procidentia. The standard Manchester operation (anterior colporrhaphy with amputation of the cervix and precervical suture of the cardinal ligaments and posterior colpoperineorrhaphy) was performed. The pouch of Douglas was not opened during the operation. Post-operative progress was satisfactory, and at follow-up six weeks later the vaginal walls and cervical stump were well healed and supported.

She remained well subsequently until the afternoon of January 6, 1960, when, while waiting in a queue, she suddenly developed a severe pain accompanied by a sensation of "something coming down" and slight bleeding per vaginam. There was no history of recent strain or trauma. After a while the pain eased, and she was taken home by car. During that night, however, she experienced gradually worsening colicky lower-abdominal pain, and on getting out of bed the next morning noticed that something was protruding from her vulva. During the ensuing day the pain continued to increase in severity, and she vomited three times prior to sending for her practitioner, who arranged her immediate transfer to hospital.

On admission at 6 p.m. on January 7, more than 24 hours after the original incident, she was restless and slightly shocked. Abdominal examination showed slight distension with obstructive bowel sounds. Vaginal examination revealed an 18–24 in. (45–60 cm.) loop of gangrenous small intestine protruding through a clean-edged transverse tear about 1 in. (2.5 cm.) long in the posterior fornix, just behind the cervical stump. The vaginal epithelium and the cervical stump were healthy

apart from the tear, and there was no recurrent prolapse. An intravenous glucose-saline drip was started, a Ryle tube was passed, and the gastric contents were aspirated. It was decided to perform an immediate laparotomy under general anaesthesia.

At operation a loop of proximal ileum was seen to be disappearing through a transverse tear $1\frac{1}{2}$ in. (3.8 cm.) long in the floor of the pouch of Douglas. The pelvic organs were otherwise normal in every way for a woman of her age. The afferent and efferent loops of the strangulated ileum were divided between intestinal clamps through healthy tissue close to the point of disappearance through the vaginal vault, and the intervening mesentery was ligated and divided. End-to-end anastomosis was then performed and the freed loop of gangrenous gut removed per vaginam, after closing the ends with continuous sutures to prevent spill of its contents. The laceration through the peritoneum of the floor of the pouch of Douglas into the posterior vaginal fornix (to which it was firmly adherent at this point) was sutured from above in one layer and reinforced by oversewing the Fallopian tubes.

Post-operatively a severe paralytic ileus developed and the "drip" and gastric aspirations had to be continued for seven days. Penicillin and streptomycin were given intramuscularly. The patient responded to this treatment and appeared to be progressing satisfactorily when, on the 14th day after operation, she suddenly collapsed and died before medical aid could be summoned. Post-mortem examination was not allowed by the relatives.

COMMENT

Spontaneous rupture of the vagina with extrusion of bowel appears to be a rare late complication of vaginal surgery, and an extensive review of the literature of the past 40 years discloses only one such case following colporrhaphy (Thompson, 1948). His patient was aged 61, the accident occurring eight years after the operation. He did not state whether the full Manchester repair had been performed, or simply anterior and posterior colporrhaphy. Thomson's case differed from the one described here in that the large bowel (pelvic colon) was involved

and rupture occurred through a recurrent enterocele, whereas in my patient there was no recurrence of the prolapse. No cause was found for the sudden rupture of the vagina, but presumably it occurred at a point of weakness caused by scarring of the vaginal vault during the previous Manchester operation. This view is supported by the finding that the peritoneum of the pouch of Douglas was firmly adherent to the underlying vaginal vault. It seems possible that the poor post-operative union of tissues often associated with old age may have been a contributory factor to the vaginal weakness in this case. If so, it may be expected, in view of the increasing age at which colporrhaphy is now performed, that instances of this kind will occur more frequently in the future.

Perusal of the literature suggests that vaginal rupture with evisceration of bowel occurs more often after vaginal hysterectomy than after colporrhaphy; Fox (1949), Kayser *et al.* (1951), and Outerbridge (1953) all report cases of this type. The decreased incidence of this admittedly rare but often fatal complication would therefore seem to be a small point in favour of the Manchester operation as opposed to vaginal hysterectomy in the treatment of uncomplicated procidentia.

I would like to thank Dr. W. Calvert for permission to publish this report.

R. F. HEYS, M.B., M.R.C.O.G.,
Registrar, St. Mary's Hospital, Manchester;
formerly Registrar in Gynaecology, Stepping Hill
Hospital, Stockport.

REFERENCES

- Fox, P. C. (1949). *Illinois med. J.*, **96**, 32.
Kayser, F. W., Heaney, N. S., and Dahleen, H. C., jun. (1951).
Amer. J. Obstet. Gynec., **61**, 1393.
Outerbridge, G. W. (1953). *Ibid.*, **65**, 439.
Thompson, W. (1948). *J. Obstet. Gynaec. Brit. Emp.*, **55**, 664.

Resected Coarctation of the Aorta and Pregnancy

Coarctation of the aorta of the adult type is an unpredictable anomaly, usually symptomless and diagnosed accidentally; but the prognosis is so poor that only one-third of the sufferers reach the age of 40. Sudden haemorrhagic death may intervene at any time, for 40% of patients rupture either the aorta itself or an intercostal, intracranial, or dissecting aneurysm. The association of coarctation with pregnancy is very rare, and the uncertain risks involved may be assessed accurately only by recording individual cases. It is, however, confidently expected that prior operative resection of the coarctation may obviate many of these dangers.

CASE REPORT

The patient was a healthy 18-year-old girl whose routine miniature chest radiographs disclosed an enlarged heart shadow. No symptoms were complained of, but she did admit to slight shortness of breath when cycling fast. Her apex beat was forceful, and there was a grade 2 aortic systolic murmur, which was conducted to the right common carotid and followed by a loud aortic second sound. A grade 1 systolic murmur was heard in all other cardiac areas, and there was a harsh continuous murmur at the junction of the innominate and right subclavian arteries. The blood-pressure in both arms was 170/100. Over the scapulae was a well-marked collateral circulation. No femoral pulses could be felt. Chest radiographs showed an enlarged left ventricle, a dilated proximal aorta, and moderate rib-notching. The electrocardiogram was normal. The diagnosis was classical coarctation.

In September, 1955, the coarctation was excised and an end-to-end anastomosis performed without tension. It was found at the classical site in relation to the ductus arteriosus,

which was not patent. The stenosed segment was about 1 in. (2.5 cm.) in length, and the lumen was reduced to about 2 mm. The ductus arteriosus and two large intercostal arteries with aneurysmal dilatations were divided. Immediately after operation pulsation was felt in the lower limbs.

The operative results were good, and for the ensuing six years she remained in excellent health and was able to undertake any physical exertion. No collateral circulation could be felt, but the pulses were palpable in both femoral and both dorsalis pedis arteries. The blood-pressure in both arms was 130/80. The systolic murmur completely disappeared, but was replaced by a grade 2 early diastolic murmur, indicating residual aortic incompetence. Screening showed that some left ventricular hypertrophy persisted.

She was advised that pregnancy would now carry no risk so far as the coarctation was concerned, and she conceived in October, 1961. Close antenatal supervision throughout revealed no rise in blood-pressure, no toxæmia, and no changes in her cardiac condition. She continued in good health until the 34th week of pregnancy, when she was awakened at 3 a.m. by a feeling of suffocation, and she collapsed. She became pulseless and cyanosed, and the blood-pressure could not be recorded. Despite blood transfusion and oxygen therapy she died eight hours later.

Necropsy showed rupture of a dissecting aneurysm of the ascending arch of the aorta, with haemorrhage into the pericardial sac and the pleural cavity. The site of operative anastomosis was undisturbed. The foetus, placenta, and gravid uterus were all normal.

COMMENT

Goodwin (1961) reviewed the literature on coarctation and pregnancy and collected reports of 147 cases with 13 deaths, a mortality of 9%. Eight of the deaths were caused by vascular rupture—three of the aorta, three of a dissecting aneurysm, and two of an intracranial aneurysm. All the deaths were in cases of unresected coarctation, and 10 were instances of undiagnosed coarctation, although proper medical examination at the initial antenatal visit should enable the diagnosis to be made (Farrelly and Thompson, 1961). However, even without operation some patients may survive repeated pregnancies without any deleterious effect (Jordan, 1958). The divergent fortunes of different patients have led to conflicting views on their management.

Dissecting aneurysm of the ascending aorta is not very rare in the young and healthy, and is found in 1 in 70 of coroners' necropsies (Jordan, 1954). In women it occurs with special frequency in association with pregnancy (Schnitker and Bayer, 1944). It is also a common cause of death in unresected coarctation even without pregnancy. So there exists a double predisposition of both pregnancy and coarctation towards developing a dissecting aneurysm.

Only recently has pregnancy been occurring in patients whose coarctation has been previously diagnosed and excised. Naturally it has been assumed that successful prior operation and careful antenatal supervision would remove the pre-existing risk of aortic rupture and dissecting aneurysm. Unfortunately that assumption proved false in the present tragic case. The risk of dissecting aneurysm may remain even in patients with resected coarctation, and the advisability of pregnancy must be regarded as doubtful.

W. M. JORDAN, M.A., B.M., B.Ch.,
D.Obst.R.C.O.G.

Bungay, Suffolk.

REFERENCES

- Farrelly, P. A., and Thompson, J. W. (1961). *J. Obstet. Gynaec. Brit. Cwlth.*, **68**, 961.
Goodwin, J. F. (1961). *Clin. Obstet. Gynec.*, **4**, 645.
Jordan, W. M. (1954). *Brit. med. J.*, **2**, 131.
— (1958). *Lancet*, **1**, 163.
Schnitker, M. A., and Bayer, C. A. (1944). *Ann. intern. Med.*, **20**, 486.