satisfactory by patient-centred students. Medical students, therefore, in addition to being psychological or organic in orientation, also differ in a dimension which seems a separate one—whether they are primarily interested in people or more concerned about, for example, neurophysiology and biochemical matters; and the two dimensions together measure whether a student approves of the course or not. The course provided was liked by psychologically oriented students who have a scientific bias. But the course is not liked so much by students who care primarily about patients, whether they are psychologically or organically oriented.

4. There are factors which do not indicate at all whether students are affective or physical. In general, these are attributes which are linked with technical learning, are not dependent on individual personal choices of students, but are taught to all students as an aspect of medical role-definition during professional training. These factors, which seem an aspect of doctoring in general and not affected by organic orientation on the one hand, or psychological orientation on the other, are: general medical ability (that is, capacity to graduate in medicine); ability to acquire factual psychiatric information; the student's confidence in his ability to diagnose and treat psychiatric disorders; extent of examination anxiety; uncertainty about having chosen medicine as a career; having difficulty in relationships with patients; and desire in later practice to obtain gratitude of patients or a good standing in the community. These attributes were found to be unrelated to psychological orientation.

We wish to express our thanks to Dr. F. M. Martin and Dr. F. A. Boddy, of the Department of Public Health and Social Medicine, University of Edinburgh, for making available the responses obtained from this student class in the Medical Student Survey of the Association for the Study of Medical Education.

REFERENCES

- Aldrich, C. K., and Bernhardt, H. (1961). Evaluation of a Change in Teaching Psychiatry to Medical Students. Unpublished report.
- Becker, H. S., Hughes, E. C., Geer, B., and Strauss, A. L. (1961). *Boys in White—Student Culture in Medical School*. University of Chicago Press.
- Caplovitz, D. (1960). Student-Faculty Relations in Medical School. Unpublished Ph.D. Thesis, Faculty of Political Science, Columbia University.
- College of General Practitioners, Working Party Report (1958). *Brit. med. J.*, **2**, 585.
- Eron, L. D. (1955). *J. med. Educ.*, **30**, 559.
- Ham, T. H. (1959). *Ibid.*, **34**, 1163.
- Hammond, K. R., and Kern, F. (1959). *Teaching Comprehensive Medical Care*. Harvard Univ. Press, Cambridge, Mass.
- Hill, D. (1960). *Brit. med. J.*, **1**, 917.
- Hubbard, J. P. (1960). *Nat. Bd Examr.*, **8**, 1.
- Langsley, D. G. (1962). *J. med. Educ.*, **37**, 429.
- Levine, M., and Lederer, H. D. (1959). In *American Handbook of Psychiatry*, edited by S. Arieti. Basic Books, New York.
- Martin, F. M., and Boddy, F. A. (1962). *Patterns of Career Choice and Professional Attitudes Among Medical Students*. In *Sociological Review Monograph No. 5*, edited by P. Halmos. University of Keele, Keele.
- Martin, W. (1957). In *The Student-Physician*, edited by R. K. Merton, G. G. Reader, and P. L. Kendall. Harvard Univ. Press, Cambridge, Mass.
- Merton, R. K., Bloom, S., and Rogoff, N. (1956). *J. med. Educ.*, **31**, 556.
- Reader, G. (1958). *Ibid.*, **33**, Part 2, 164.
- Richardson, I. M. (1962). *Brit. med. J.*, **2**, 582.
- Schumacher, C. F., and Gee, H. H. (1961). *J. med. Educ.*, **36**, Part 2, 155.

EXAMINATION PERFORMANCE OF MEDICAL AND NON-MEDICAL STUDENTS AT CAMBRIDGE

BY

GABRIEL HORN, M.A., M.B., Ch.B., B.Sc.

Department of Anatomy, University of Cambridge

Many opinions have been expressed both by headmasters and by deans of medical schools concerning the academic standards of entrants to the medical profession (B.M.A., 1962). For the most part these opinions are based on subjective experience and not infrequently suggest that the number of able people entering the profession is declining (B.M.J., 1962). The present survey was therefore undertaken to see whether there is any support for the view that the performance of medical students in certain university examinations at Cambridge is less satisfactory than that of a group of non-medical students.

The system of university examinations at Cambridge is complicated, and any generalization about it is apt to be misleading in many special cases. Bearing in mind these reservations, however, certain generalizations can be made.

At the end of their first year at the university students of the Natural Sciences (including medical students) sit for the Preliminary Examination for Part I of the Natural Sciences Tripos. Some students then proceed to the examination for Part I of the Tripos at the end of the second year, while others take this examination at the end of their third year. The majority of non-medical students sit for the Part I Examination after two years spent at the university, and the majority of medical students sit for that examination after three years.

The Natural Sciences Tripos may be taken in different combinations of subjects. Medical students are required to read anatomy and physiology as "whole" subjects

together with at least two "half" subjects, frequently pathology and biochemistry. Non-medical students, of course, take other combinations of subjects. Placing in the class lists—that is, class I, II 1, II 2, III, or failed the examination—for Part I of the Tripos is determined by the overall performance in the various subjects taken by the candidate.

There is a suggestion (Horn, 1963) that medical students are awarded fewer first-class honours in the examination for Part I of the Natural Sciences Tripos than students who read other combinations of subjects. This may happen because students who come up intending to read medicine have fewer of the more able individuals among them than non-medical students. Another explanation for this difference, however, is that it may be easier to obtain high marks in subjects which tend not to be read by medical students—for example, physics, chemistry, mathematics. An explanation of this kind could account for the trend toward a higher proportion of firsts among non-medical natural scientists. It was therefore considered that if the performance of medical students could be compared with that of non-medical students in the same examination this particular variable would be taken into account.

Material

The performance of the two groups in two sets of examinations was analysed. One of the examinations was the Preliminary Examination in Physiology, which is taken

when both groups of students have attended the same course for the same length of time. The other examination was Part I of the Natural Sciences Tripos in Physiology, which is taken by the majority of non-medical students at the end of their second year and by the majority of medical students at the end of their third year at the university. Both medical and non-medical students take the same examination papers and practicals. Performance in the Preliminary Examination in Physiology was analysed for the years 1954 to 1962 inclusive. For the Part I Examination the years 1955, 1956, 1957, 1958, 1959, 1960, and 1962 were analysed. The choice of these years was determined solely by the availability of the relevant mark-books from which the data had to be extracted.

In the nine-year period over which the results of the Preliminary Examination were analysed, 1,688 medical and 452 non-medical students sat for the examination. Of the non-medical students 321 read, in addition to physiology, at least one of the subjects—physics, chemistry, mathematics. Of these 321 candidates, 312 read chemistry.

Results

Preliminary Examination in Physiology

The percentages of students listed in each class for each of the nine years analysed are given in Table I. A chi-square test on the numerical data summed for the whole period (Table I, bottom row) was performed. $\chi^2=11.73$, and, with four degrees of freedom, $P<0.02$. This probability level suggests that the distribution of candidates among the various classes is different in the two groups. A number of *t* tests were therefore performed on the columns of percentages in Table I. The percentages of medical students listed in class I over the years were compared with those of non-medical students listed in the same class. The value of *t* was 0.76 and $0.4<P<0.5$. There are thus no significant differences between the proportions in each group awarded first-class honours. The value of *t* for the comparison of the percentages in the II 1 classes is 1.19 and $0.2<P<0.3$. For the II 2 classes $t=1.53$ and 0.1

$<P<0.2$. There are no significant differences between the proportions of the two groups listed as having failed the examination ($t=0.90$, $0.3<P<0.4$). The proportion of

Percentages from Table I (Preliminary Examination in Physiology) plotted to show variations in the percentage of students in each class during 1954 to 1962 inclusive. Curves for medical students are shown as continuous lines, those for non-medical students as broken lines.

medical students placed in class III was higher (6% on average) than the proportion of non-medical candidates placed in this class. This was the only class where the differences achieved statistical significance ($t=3.21$, $P<0.02$).

TABLE I.—Results of the Preliminary Examination in Physiology Taken at the End of the First Year of the Course

Year	Medical Group							Non-medical Group						
	Class (Data expressed as percentages)					Totals		Class (Data expressed as percentages)					Totals	
	I	II 1	II 2	III	Fail	%	No.	I	II 1	II 2	III	Fail	%	No.
1954 ..	4.9	8.8	31.4	48.0	6.9	100	204	0	7.3	46.3	41.5	4.9	100	41
1955 ..	4.0	11.5	48.3	34.9	1.3	100	226	6.3	9.4	53.0	28.2	3.1	100	32
1956 ..	6.8	14.6	40.2	36.9	1.5	100	206	5.1	10.2	46.3	38.4	0	100	39
1957 ..	5.9	13.9	51.0	25.2	4.0	100	202	13.3	20.0	51.1	15.6	0	100	45
1958 ..	1.9	11.4	40.6	39.0	7.1	100	210	1.9	25.0	44.3	26.9	1.9	100	52
1959 ..	5.2	20.8	37.6	34.7	1.7	100	173	6.6	21.3	45.8	23.0	3.3	100	61
1960 ..	4.6	11.2	47.4	34.2	2.6	100	152	3.6	9.1	45.5	38.2	3.6	100	55
1961 ..	3.7	19.7	46.9	27.8	1.9	100	162	4.4	23.2	53.5	18.9	0	100	69
1962 ..	7.2	11.8	50.2	30.1	0.7	100	153	12.0	18.9	39.9	25.8	3.4	100	58
Mean % ..	4.9	13.7	43.8	34.5	3.1	100	—	5.9	16.0	47.4	28.5	2.2	100	—
Total Nos.	82	229	734	589	54	—	1,688	27	77	213	125	10	—	452

TABLE II.—Results of the Examination in Physiology for Part I of the Natural Sciences Tripos

Year	Medical Group							Non-medical Group						
	Class (Data expressed as percentages)					Totals		Class (Data expressed as percentages)					Totals	
	I	II 1	II 2	III	Fail	%	No.	I	II 1	II 2	III	Fail	%	No.
1955 ..	7.3	13.7	35.6	38.5	4.9	100	205	2.9	8.9	52.9	32.4	2.9	100	34
1956 ..	7.0	20.1	46.7	25.7	0.5	100	214	9.7	9.7	48.4	29.0	3.2	100	31
1957 ..	5.7	13.5	53.8	27.0	0	100	230	3.0	3.0	66.7	27.3	0	100	33
1959 ..	7.9	22.3	52.0	17.8	0	100	202	5.7	30.2	45.4	18.7	0	100	53
1960 ..	5.4	19.9	52.7	22.0	0	100	186	14.3	25.0	46.4	12.5	1.8	100	56
1962 ..	4.1	27.4	56.9	11.6	0	100	146	7.7	23.1	46.1	20.5	2.6	100	78
Mean % ..	6.2	19.5	49.6	23.8	0.9	100	—	7.2	16.7	50.9	23.4	1.8	100	—
Total Nos.	75	224	583	290	11	—	1,183	22	55	141	62	5	—	285

The percentages in the various classes are plotted in the Chart. The curves illustrate the extent of overlap between the two groups. They also provide some indication of trends over the period studied.

There is no sign of any fall over the years in the percentage of medical students listed in the first or upper second classes. There is a slight trend for the percentage of medical students listed in the lower second class to increase, and a suggestion that the proportion who fail the examination and the proportion who fall into class III are gradually declining. The distribution of non-medical students among the various classes appears to have remained more or less stationary.

The overall picture suggests that, over the period studied, proportionately fewer medical candidates for this examination fail or are placed in class III. These changes may be brought about because the standard of academic attainment of the candidates is improving or because the policy of the examiners is changing. The latter explanation seems an unlikely one, since there is very considerable fluctuation from year to year in the proportions placed in each class, and because the results for the non-medical students appear to have changed little over the nine years analysed.

Part I Examination in Physiology

The results of this examination were analysed over a six-year period. The distribution of each group of students among the various classes is given, for each year, in Table II, together with the total number in each class. A chi-square test performed on the latter data (Table II, bottom row) showed that the distribution of medical students among the various classes does not differ significantly from that of non-medical students ($\chi^2=2.81$, and, with four degrees of freedom, $0.5 < P < 0.7$). The data were not plotted graphically since there are two years (1958 and 1961) for which no data were available. However, if the percentages in Table II are examined it will be found that a trend similar to that described for the Preliminary Examination (see Chart) appears to exist. In addition there seems to be a notable drop in the proportion of both groups placed in class III and in the proportion of medical students listed as having failed the examination.

Discussion

The fact that the performance of medical students in the Examination in Physiology for Part I of the Natural Sciences Tripos does not differ significantly from that of non-medical students may not be regarded as remarkable, since the majority of medical students spend three years at the university before sitting the Part I Examination in Physiology, while the majority of non-medical students spend only two. It could legitimately be argued that the former group require the extra year to reach the same standard as the latter group. This argument loses much of its force with regard to the results of the Preliminary Examination in Physiology. Both groups of students spend the same length of time reading the subject before taking the examination, and the only statistically significant difference that emerges is that medical students are listed slightly more frequently (6% on average) in class III than the non-medical group. Even this observation is difficult to interpret. It may be that a few more weak men read medicine than read other subjects. Alternatively, these results may only reflect the examination strategy of some of the medical candidates. A less able medical student will almost certainly spend three years over the Tripos Part I and so has longer to revise than the majority of

non-medical students. For this reason such a medical candidate may not have as much incentive to work industriously during his first year at the university as the non-medical student might have.

If the performance in the Physiology Examination is any guide to the academic ability of medical students, then there is no reason to suppose either that this group of students has less than its share of the more able men (see percentages in classes I and II 1) or that the academic quality of these students is declining, at Cambridge at least. If anything, it may be improving.

Summary

The performance of medical and non-medical students in certain university examinations at Cambridge has been examined over a period of several years.

In the Physiology Examination for Part I of the Natural Sciences Tripos there are no statistically significant differences between the two groups of students listed in the various classes.

In the Physiology Examination (first-year Preliminary) taken by both groups at the end of the first year of the course in Physiology, the only difference, significant at the 2% level, is that the percentage of medical students listed in class III is higher (6% on average) than the percentage of non-medical students listed in this class. The possible reasons for these differences are discussed.

There is no evidence to support the view that fewer more able men read medicine than other subjects.

Over the period examined the percentage of medical students listed in cases III or as having failed the Preliminary Examination appears to have declined slightly.

It is a pleasure to thank Mrs. S. Traut for her invaluable assistance in all stages of this study, and the Medical Research Council for financial assistance.

REFERENCES

- British Medical Association (1962). *Recruitment to the Medical Profession*, B.M.A., London.
Brit. med. J., 1962, 1, 1325.
 Horn, G. (1963). *Camb. Rev.*, 84, 257.

"The Laboratory Animals Centre's new animal house at Carshalton was put into use on April 30 this year, after appropriate sterilization procedures. The animal house has been designed so that mice and rats can be bred under conditions of rigid isolation. The foundation stocks of mice and rats are derived by caesarean section from strains already maintained under conventional conditions at the centre. The caesarean sections are carried out under aseptic conditions outside the building and the intact uteri are introduced into the building through a dunk tank of disinfectant. The foetuses are removed from the uteri and reared by fostering them on to mice and rats derived from germ-free stocks obtained from two laboratories in the United States. These stocks were imported from the U.S.A. in special containers. The animals were transferred directly into the new animal house and allowed to acquire a non-pathogenic flora within these isolated conditions. This process was assisted by deliberately contaminating some of the animals with *Lactobacillus*. The stocks introduced into the building by caesarean section and fostering will, therefore, acquire a similar non-pathogenic flora. Details of the gut flora may be obtained from the L.A.C. In order that Carshalton mice shall deserve a reputation for a high standard of health the stocks will be tested regularly for the presence of a number of the more important pathogens. As a further measure, designed to reduce the chances of a build-up of undesirable parasites, each strain will be refounded from time to time, probably via a germ-free cycle." (Pamphlet, Laboratory Animals Centre, August, 1963.)