BLOOD-PRESSURE IN UNDERNOURISHED JAVANESE

BY

K. V. BAILEY, M.D.

Indonesian Nutrition Institute, Wonosari, Gunung Kidul, Jogjakarta

Many observations have been made indicating that arterial blood-pressure, especially systolic, tends to be lower in undernourished subjects (Keys *et al.*, 1950). During studies of nutrition made for the Indonesian Nutrition Institute in Java in 1957-9 blood-pressures were measured with a view to comparing the relative effects of low-calorie and low-protein diets on pressures. The geographical and dietary circumstances are described elsewhere (Bailey, 1960, 1961).

Materials and Methods

Blood-pressure was studied in three villages. One, Pakem, was in the densely populated irrigated ricefield area near Jogjakarta. The other two, Tjuwelo and Kedungpoh, were in the unfertile limestone hill region of Gunung Kidul, further east of Jogjakarta, where cassava (tapioca) was the staple foodstuff.

In these villages all available people were mustered at the headman's home or office where I measured the blood-pressures. About half the adult population in each village were measured. Factors determining the composition of the sample in each village were similar, absentees being chiefly men and women under 30 who were out in the fields working, and some decrepit old people who stayed at home.

Casual blood-pressures were measured in the same way as those in Pacific Islanders reported by Lovell *et al.* (1960) except that diastolic pressure was taken at the point of disappearance of sounds. Weights, heights, and arm circumferences midway between acromion and olecranon were also measured. The ages recorded were guesses based on the knowledge of the village headman of certain dates, such as the two world wars, and on my own impression based on four years' residence in Java.

Results

Nutrition.—Estimates of calorie and protein intake are summarized in Table I. Calorie intakes were low and fairly similar in all the villages, but lowest in Kedungpoh. Protein intakes were distinctly better in Pakem than in the other villages because rice contains 7½% protein whereas cassava (equicaloric with rice) contains only 1½%.

TABLE I.—Diets in Javanese Villages Surveyed

Village	Staple Foodstuff	Daily Intakes per Adult Man	
		Calories	Protein (g.)
Pakem	Rice (+cassava)	1,500-1,700	30-40
Tjuwelo	Cassava (+maize)	1,500-1,700	18-20
Kedungpoh	Cassava	1,400-1,600	15-18

This pattern of low-calorie intake, with protein varying according to staple foodstuff, was noted elsewhere in the poor regions of Java. Salt intake was about 8-10 g. daily in all the villages. In Pakem there was no nutritional oedema, but in Tjuwelo and Kedungpoh about 5% of the population aged 20-40 and 18% of those aged over 40 had nutritional oedema, due usually to calorie deficiency superimposed on severe protein deficiency.

Blood-pressure.—Fig. 1 relates mean systolic and diastolic pressures to age and sex in each village. The blood-pressures and age structure were similar in each village and so the findings are combined in Table II.

FIG. 1.—Mean systolic and diastolic pressures relative to age and sex.

FIG. 2.—Weight/height ratios relative to age and sex.

TABLE II.—Means and Standard Deviations of Systolic and Diastolic Blood-pressures in 3 Javanese Villages (Combined)

Age-group	No. of Subjects	Males								Females							
		Age		Systolic		Diastolic		No. of Subjects		Age		Systolic		Diastolic			
		Mean	S.D.	Mean	S.D.	Mean	S.D.			Mean	S.D.	Mean	S.D.	Mean	S.D.		
17-19	90	18.0	0.8	110.02	15.06	67.98	9.27	61		17.9	0.5	118.56	16.01	75.95	11.07		
20-29	129	23.7	3.0	112.21	14.73	66.84	10.12	146		23.3	3.0	117.40	13.25	74.53	11.49		
30-39	123	33.7	3.0	112.72	15.81	69.99	8.27	197		33.2	2.8	114.65	15.31	72.08	10.62		
40-49	139	43.2	2.8	115.15	14.84	71.27	8.86	181		42.6	2.8	123.44	18.75	75.28	9.50		
50-59	123	52.0	2.7	123.76	21.25	73.03	11.43	189		51.9	2.6	127.88	19.58	74.82	10.20		
60+	83	62.5	4.3	128.76	20.45	74.90	11.59	96		63.3	4.9	143.29	24.69	77.23	11.98		

Body Build.—The better-nourished Pakem villagers had greater mean weight, height, weight/height ratios (W/H), and arm circumferences than the other villagers. The differences between them in body build are illustrated in Fig. 2, which shows the mean W/H ratios in relation to age and sex in each village.*

Discussion

The mean systolic, but not diastolic, pressures recorded in this study are lower than those found in Western peoples—Hamilton *et al.* (1954)—and in several Asian and Pacific peoples (Lovell *et al.*, 1960). Low blood-pressure has been recorded in semi-starvation (Keys *et al.*, 1950) and other poorly nourished groups—for example, Chinese rickshaw men (Lowenstein, 1954)—but whether this is due to lowered calorie intakes only or in part to lowered protein intakes is not known. In these Javanese villagers, with very low intakes of both, differences in protein intake do not seem to influence blood-pressure.

The relation of blood-pressure to age varies in different peoples. In Western peoples the mean systolic and diastolic pressures rise with age, while in several other populations rises have been found to be slight or absent; these include poor Chinese (Kean and Hammill, 1949), East Africans (Donnison, 1929), Gilbertese (Maddocks, 1961), and New Guinea Highlanders (Whyte, 1958). In these Javanese villagers systolic pressures rise with age. It seems, therefore, that while malnutrition may be associated with relatively low systolic pressure, it does not prevent the rise with age which is found in better-nourished peoples. However, renal disease was not eliminated in the Javanese as a possible cause of this rise (cf. Stirling, 1960).

*Detailed measurements available on request. (Author's present address, 5 Lamington Street, Deakin, Canberra, A.C.T., Australia.)

Summary

Blood-pressure and measures of body build were recorded from three Javanese villages where there was general undernutrition. Rice was the staple foodstuff in one, and cassava in two; protein intakes were appreciably better in the former.

Blood-pressure levels were similar in each village. Systolic pressures (but not diastolic) were low compared with other populations and systolic pressures rose with age.

In these undernourished subjects, protein deficiency does not seem to be the cause of lowered systolic pressure, and low calorie and protein intakes do not prevent rising systolic pressures with age.

I am indebted to Professor R. R. H. Lovell, University of Melbourne Department of Medicine, and his staff, and to Miss B. Laby, University of Melbourne Department of Statistics, for guidance and assistance in the preparation of this report.

REFERENCES

- Bailey, K. V. (1960). *Rural Nutrition Studies in Indonesia* (Thesis), University of Melbourne.
 — (1961). *Trop. geogr. Med.*, **13**, 216, 233.
 Donnison, C. P. (1929). *Lancet*, **1**, 6.
 Hamilton, M., Pickering, G. W., Roberts, J. A. F., and Sowry, G. S. C. (1954). *Clin. Sci.*, **13**, 11.
 Kean, B. H., and Hammill, J. F. (1949). *Arch. intern. Med.*, **83**, 355.
 Keys, A., Brozek, J., Henschel, A., Mickelsen, O., Taylor, H. L., *et al.* (1950). *The Biology of Human Starvation*. University of Minnesota Press, Minneapolis.
 Lowenstein, F. W. (1954). *Amer. Heart J.*, **47**, 874.
 Lovell, R. R. H., Maddocks, I., and Rogerson, G. W. (1960). *Aust. Ann. Med.*, **9**, 4.
 Maddocks, I. (1961). *Lancet*, **2**, 396.
 Stirling, G. A. (1960). *Brit. med. J.*, **1**, 1173.
 Whyte, H. M. (1958). *Aust. Ann. Med.*, **7**, 36.

GUANETHIDINE IN THE TREATMENT OF HYPERTENSION

BY

CLIFTON P. LOWTHER, M.B., M.R.C.P.G.
M.R.C.P.Ed.

Senior Medical Registrar, Western General Hospital,
Edinburgh

RICHARD W. D. TURNER, O.B.E., M.A., M.D.
F.R.C.P., F.R.C.P.Ed.

Senior Lecturer, Department of Medicine, University of
Edinburgh; Physician, Western General Hospital, Edinburgh

This paper describes the treatment of 75 patients with guanethidine over periods ranging from 3 to 36 months, with an average duration of 24 months. Sixty-seven have been treated for more than one year, 58 for more than 18 months, and 40 for more than two years. The indications for treatment, selection of patients, measurement of blood-pressure, and classification of results have been similar to those previously described in relation to other drugs (Lowther and Turner, 1960). Forty-one of these patients had been treated with various other preparations before guanethidine was given; the remaining 34 were treated with guanethidine from the outset.

The assessment of the drug has been based solely on its effect on blood-pressure and on the incidence of tolerance and side-effects. Benefits such as the regression of retinopathy or of ventricular hypertrophy are secondary to the fall in blood-pressure, however produced, and not directly attributable to any particular drug.

Material and Methods

Clinical Features.—Only those patients have been treated who, in addition to high resting diastolic pressures, showed

objective evidence of secondary changes in the fundi or heart.

Forty patients were male and 35 female. The average age was 54 years: 10 were under 40 and 14 more than 60. Clinical details are given in Table I, which shows that these features indicate a fair cross-section of patients with severe hypertensive disease. Retinal changes were classified for convenience according to the criteria of Keith, Wagener,

TABLE I.—Clinical Details (75 Cases)

	No. of Cases
Method of presentation	
Left ventricular failure	17 (23%)
Cerebrovascular accident	14 (19%)
Headache	17 (23%)
Disturbance of vision	10 (13%)
Routine examination	11 (15%)
Miscellaneous	6 (7%)
Resting diastolic blood-pressure (mm. Hg)	
150+	11 (15%)
130-149	39 (52%)
110-129	25 (33%)
Retinal changes	
Grade 4	8 (11%)
" 3	27 (36%)
" 2	38 (51%)
" 1	2 (2%)
Left ventricular hypertrophy (E.C.G.)	
Present	72 (96%)
Absent	3 (4%)
Renal function	
Impaired	13 (17%)
Renal failure	4 (5%)
Normal	58 (78%)