NALIDIXIC ACID IN URINARY INFECTIONS

BY

J. F. N. C. WARD-McQUAID, M.S., F.R.C.S.

Consultant Surgeon

D. JICHLINSKI, M.D.

Surgical Registrar

R. MACIS, A.I.M.L.T.

Laboratory Technician

Mansfield Group of Hospitals

Nalidixic acid ("negram") is one of a new series of 1,8-naphthyridine derivatives with antibacterial properties which have been synthesized by Leshner *et al.* (1962). Chemically it is 1-ethyl-7-methyl-1,8-naphthyridine-4-one-3-carboxylic acid.

McChesney *et al.* (1963) have demonstrated *in vitro* activity against pathogenic Gram-negative bacteria, and studied human and animal urine after oral administration. Lishman and Swinney (1963) have carried out a clinical trial with 60 patients and summarized the results of data obtained at the Sterling-Winthrop Research Institute. The present paper describes a short clinical trial of this compound in the urological out-patient department and in the wards of a general hospital. The range of activity *in vitro* and the results of treatment in 52 patients with acute or recurrent urinary infection are assessed. This new oral antibacterial agent is henceforth referred to as nalidixic acid.

Pharmacology

The data in this section were obtained from the files of the Sterling-Winthrop Research Institute.

Laboratory Animals

Nalidixic acid is well tolerated in mice, in which an oral LD_{50} of approximately 4,000 mg./kg. has been shown. Chronic toxicity studies were carried out on rats, dogs, and monkeys for over a year. The drug was well tolerated in monkeys at the three dose levels of 25, 75, and 225 mg./kg./day. Single oral doses as high as 900 mg./kg. produced no evidence of toxicity in monkeys. Haematological studies, including total blood-cell count, haemoglobin concentration, and haematocrit values, were performed monthly. All results were within normal limits. Other analyses carried out at the end of the study showed no deviations from normal limits. These were: serum glutamic oxalo-acetic transaminase, serum glutamic pyruvic transaminase, clotting-time, prothrombin time, serum alkaline phosphatase, cephalin-cholesterol flocculation, urea nitrogen, and total cholesterol. Routine urinalyses before and after the medication period showed all values to be within normal limits. No generalized sedative or stimulant effect on the central nervous system can be ascribed to nalidixic acid, and no effects on the autonomic nervous system, renal function, or gastro-intestinal tract have been noted.

Pregnancy Tests.—The drug was given orally to pregnant rats and monkeys. Rats were given doses of 75 and 150 mg./kg. throughout pregnancy and for one week afterwards. On these dose levels there were no adverse effects

on the course of pregnancy or on the growth and survival capacities of the offspring. At the highest dose level of 300 mg./kg., which is that approaching the limit of rat tolerance, there was maternal mortality and interference with the course of pregnancy of the survivors. At the end of the study there was no evidence of gross disease among survivors at all dose levels. Doses of 200 mg./kg./day were given to three pregnant monkeys. One animal was medicated for two months and others for one month prior to delivery. The drug was given to all animals during the first two weeks post partum. No untoward reactions were noted. Each mother delivered one normal offspring, and mothers and offspring were observed for three months after stopping medication and appeared normal during this time. The body weight of the mothers was normal throughout the study. The Somers test has been repeated using pregnant rabbits; no teratogenic changes or other congenital malformations were observed.

Human Volunteers

The drug was given to 13 men and 9 women four times a day for up to 180 successive days. The maximum sustained dose was 8 g./day for 56 days. An initial single 4-g. dose was well tolerated by most volunteers.

Clinical Trial

Mid-stream specimens of urine were collected in both sexes except when a specimen was taken at the time of cystoscopy. The urines were refrigerated and cultured within an hour of arrival in the laboratory.

Range of Activity *In Vitro*

A simple disk technique was used to assess sensitivity to nalidixic acid and eight other commonly used antibacterial agents (Table I). Not all organisms were tested

TABLE I.—Disk Sensitivity of 133 Organisms

Organism	Total	Nalidixic Acid	Sulphafurazole	Nitrofurantoin	Chloramphenicol	Tetracycline	Penicillin	Streptomycin	*Kanamycin	*Mandelamine
<i>E. coli</i> ..	47	41	31	45	46	34	1	39	7/7	6/7
Coliform ..	6	5	4	4	6	5	0	5	2/2	2/2
<i>Proteus</i> ..	20	15	8	15	15	3	1	9	6/6	6/6
<i>K. aerogenes</i> ..	24	8	5	12	14	11	0	8	13/13	13/13
<i>Ps. pyocyanea</i> ..	3	1	0	3	1	2	0	1	0/3	1/3
<i>Str. faecalis</i> ..	26	6	4	24	25	16	19	7	1/2	2/2
<i>Str. viridans</i> ..	1	0	1	1	1	1	1	0	0/0	0/0
<i>Staph. pyogenes</i> ..	6	3	1	1	5	3	2	3	0/0	0/0
Total ..	133	79	54	105	113	75	24	72	29/33	30/33

against mandelamine and kanamycin.* A 30 μ g. disk of nalidixic acid and standard disks of the other agents were used in this study.

The 133 organisms isolated include the 63 organisms treated in 52 patients, described in more detail in the next section and summarized in Table II. Nalidixic acid compares favourably with other antibacterial drugs commonly

used against Gram-negative organisms. Of the 53 *Escherichia coli* and other coliform organisms 46 (86%) were sensitive to nalidixic acid, while 15 out of 20 (75%) *Proteus* organisms were also sensitive. *Klebsiella aerogenes* (8 of 24) and *Pseudomonas pyocyanea* (1 of 3) were less sensitive. A surprising finding was the sensitivity of 6 of 26 *Streptococcus faecalis*, and 3 of 6 *Staphylococcus pyogenes*. The sensitive *Str. faecalis* organisms were found early in the series, when primary plating was used for sensitivity tests in mixed cultures.

Treatment

All the 52 patients had clinical and bacteriological evidence of infection, with excess of leucocytes and a positive culture (Table II); follow-up urine specimens were taken and appropriate sensitivity studies were made. The ages ranged widely, from 10 to 89. There were 26 patients with acute infections, previously untreated with antibacterial agents, and 26 with recurrent or chronic infections, many

TABLE II.—Disk Sensitivity of 63 Organisms (52 Patients)

Organism	Total	Nalidixic Acid	Sulpha-furazole	Nitro-furantoin	Chloram-phenicol	Tetra-cycline
<i>E. coli</i> ..	31	29	21	31	31	27
Coliform ..	3	3	3	3	3	3
<i>Proteus</i> ..	10	7	4	8	8	1
<i>K. aerogenes</i> ..	7	4	2	7	7	6
<i>Ps. pyocyanea</i> ..	1	1	1	0	0	1
<i>Str. faecalis</i> ..	9	5	1	9	9	8
<i>Staph. pyogenes</i> ..	2	1	1	1	1	1
Total ..	63	50	33	59	59	47

with an acute exacerbation, who had received previous treatment, but not with nalidixic acid. The patients with acute infections, and some of those with recurrent infections, were started on nalidixic acid after the urine specimen had been taken but before the result of the culture and sensitivity were known. The underlying causes are summarized in Table III, and the bacteriological results in Table IV.

TABLE III.—Underlying Causes in Patients Treated with Nalidixic Acid (Failures in Parentheses)

	Total	New	Recurrent
Pyelocystitis ..	34	19 (2)	15 (6)
Epididymitis and prostatitis ..	8	5	3
Bladder tumour ..	4	0	4 (3)
Urolithiasis (operated) ..	4	1	3 (2)
Urethral stricture ..	1	0	1 (1)
Renal injury ..	1	1 (1)	0
Total ..	52	26	26

TABLE IV.—Bacteriological Results of Treatments with Nalidixic Acid in 63 Organisms (52 Patients)

	Total	Success	Failure
<i>E. coli</i> ..	31	24	7
Coliform ..	3	3	0
<i>Proteus</i> ..	10	7	3
<i>K. aerogenes</i> ..	7	4	3
<i>Ps. pyocyanea</i> ..	1	0	1
<i>Str. faecalis</i> ..	9	7	2
<i>Staph. pyogenes</i> ..	2	1	1
Total ..	63	46	17

Treatment was by 1 g. (2 tablets) of nalidixic acid four times a day for 14 days. A mid-stream specimen of urine was collected immediately before treatment was started and a further specimen 10 days after. If the second specimen was sterile a third was taken 14 days after completion of treatment. Where the second specimen was not satisfactory nalidixic acid therapy was stopped and another

drug substituted. Alkalis were not used, nor were any special precautions taken with fluid intake.

Results in Acute Group

This group of 26 patients consisted of 11 males and 15 females, with 32 infecting organisms, there being six double infections. Clinical cure was obtained in 24 of the 26 and a bacteriological cure in 23. In two of the three failures the drug was given before the sensitivity results were available and the organisms were found to be resistant. On the other hand, sensitivity tests suggested that 3 out of 29 successfully treated organisms would prove to be resistant.

Analysis of Results: Infecting Organisms

E. coli and Other Coliforms.—17 infections; 16 sensitive. There were two failures—one a sensitive coli organism becoming resistant in cystitis; the other a sensitive coli organism replaced by resistant *K. aerogenes* in severe renal injury.

Proteus.—Five infections: two sensitive. There was one failure—a sensitive organism replaced by resistant *K. aerogenes* in the patient with severe renal injury mentioned above.

K. aerogenes.—Two infections; both sensitive. Two clinical cures.

Str. faecalis.—Eight infections; five sensitive. One failure—a resistant organism in cystitis.

Results in Recurrent or Chronic Group

All these 26 patients had had other previous antibacterial treatment. There were 15 males and 11 females, and 31 infecting organisms (five double infections). A clinical cure was obtained in 20 patients, but in only 14 of the 26 was there bacteriological evidence of cure.

Had we waited for and relied upon the sensitivity results before starting treatment, nalidixic acid would not have been prescribed in 6 of the 12 failures, as the organisms were resistant.

Analysis of Results: Infecting Organisms

E. coli and Other Coliforms.—17 infections; 16 sensitive. There were five failures: due to a resistant organism in cystitis, and to bladder carcinoma; to a sensitive organism becoming resistant to cystitis; to a sensitive organism replaced by resistant *Str. faecalis* in cystitis and by a resistant *K. aerogenes* in bladder tumours.

Proteus.—Five infections; five sensitive. The two failures were in urethral stricture replaced by resistant *Str. faecalis*; and persistence of sensitive organisms in acid urine after pyelolithotomy.

K. aerogenes.—Five infections; two sensitive. Four failures in pyelonephritis; cystitis and bladder carcinoma, and replacement of a sensitive organism in pyelitis.

Str. faecalis.—One infection; insensitive. One failure.

Staph. pyogenes.—Two infections; one sensitive. This sensitive organism became resistant in cystitis.

Side-effects

A transient rash occurred in 1 of these 52 patients. In treating 120 patients with nalidixic acid the only other side-effect was nausea in a patient with a duodenal ulcer.

Discussion

Range of Activity in Vitro

The range of activity of nalidixic acid *in vitro* as shown by our disk-sensitivity tests broadly resembles that by disk

and tube-dilution techniques in the few relevant papers published. When compared with other commonly used antibacterial agents, Barlow (1963), Lishman and Swinney (1963), Slade (1963), Thompson and Rae (1963), and Buchbinder *et al.* (1963) found nalidixic acid more effective than sulphafurazole, tetracycline, or nitrofurantoin against *E. coli*. The first three of these authors also found this true for *Proteus* organisms, and Thompson and Rae (1963) stated that only chloramphenicol was more effective. Lishman and Swinney (1963) thought nalidixic acid the more effective agent against *K. aerogenes*. We found it as effective against *Proteus* organisms as chloramphenicol and nitrofurantoin but less so than kanamycin, and rather less so than these antibacterial agents against *E. coli*, other coliforms, and *K. aerogenes* organisms. Nalidixic acid was slightly more effective than tetracycline and streptomycin, and considerably more than sulphafurazole against the common Gram-negative organisms, *E. coli*, other coliforms, *Proteus*, and *K. aerogenes*. We found a higher proportion of *Str. faecalis* organisms sensitive to nalidixic acid than did the other authors, but this may have been due to primary plating in mixed infections. The interpretation of mandelamine disk sensitivity has recently again been questioned by Waterworth (1962).

Results of Treatment

Our course of treatment of 14 days was perhaps longer than necessary in some patients and longer than that used by other authors. Our results, both in the whole series and in infections by individual organisms, agree broadly with those of Carroll (1963) and those of the authors already quoted. The cure rate in acute infections (23 out of 26) and in chronic infections (14 out of 26) is similar to, and in chronic infections better than, results reported by Barlow (1963) with nalidixic acid and previous authors using various agents (Rhoads *et al.*, 1952; Garrod *et al.*, 1954; Kass, 1955; Turck *et al.*, 1962). Our figures would have been better had we treated only sensitive organisms and omitted the eight failed cases due to insensitive organisms. Development of resistance during treatment with nalidixic acid occurred in 2 of the 34 *E. coli* and other coliforms but in none of the 10 *Proteus* organisms.

Nalidixic acid has been given safely to toxic patients with renal failure. There is doubt whether the administration of alkalis greatly affects the success of treatment (Slade, 1963). Most authors did not use them.

Assessment of urinary antibacterial drugs is notoriously difficult because of the many variable factors involved, especially the wide variety of underlying causes, and in the case of nalidixic acid the relatively small samples available for study.

Therapeutic Use of Nalidixic Acid

In acute urinary infections, where treatment is started before urine culture and sensitivity results are known, or the cause is diagnosed, a wide range of different antibacterial agents are used. The newer preparations with a wider spectrum are often preferred to the older sulphonamides. Nitrofurantoin has recently been used increasingly without preliminary investigations. There are well-known disadvantages to the use of chloramphenicol, tetracycline, and kanamycin, which at present do not seem to justify their routine use in uninvestigated acute cases.

Our own practice has been to use sulphonamides first, unless the infection is fulminating, and to reserve more powerful drugs until the sensitivity of the organisms and the underlying causes of infection are known. Develop-

ment of resistance means loss of a valuable antibacterial agent to the patient under treatment, and also, in time, to other patients. The range of nalidixic acid is wider than that of the sulphonamides, but less than that of nitrofurantoin. It may well prove to be a satisfactory drug to use empirically in acute infections while awaiting the sensitivity results. Although about a fifth of acute urinary infections are due to *Str. faecalis*, nalidixic acid may deal with many of these.

In recurrent or chronic infections it is usual to wait for the result of sensitivity tests unless an acute exacerbation demands urgent treatment. One then often acts on the last laboratory report of culture and sensitivity. Present evidence suggests that nalidixic acid is as effective as nitrofurantoin in infection due to *E. coli*, other coliforms, and *Proteus* organisms. It may also act faster (Jameson and Swinney, 1963). It should prove valuable in these infections, at the very least in patients with toxic reactions to nitrofurantoin.

Some urologists use antibacterial agents as a prophylactic against infection. We have observations (unpublished) on 30 patients undergoing cystoscopy (10 with an indwelling catheter and 20 after prostatectomy, some already infected) treated with nalidixic acid. These do not justify comment except to say that when the urine was infected or became infected by an organism sensitive to nalidixic acid this organism was usually replaced by a resistant *Str. faecalis*, *K. aerogenes*, or staphylococcus. Urinary infection in these cases, as in those reported earlier in this paper, may be due to a cross-infection with a resistant organism, especially in the case of a *K. aerogenes* found in one particular ward.

Results of the long-term use of nalidixic acid in recurrent infection and pyelonephritis are not yet available.

Summary

Nalidixic acid ("negram") (1-ethyl-7-methyl-1,8-naphthyridine-4-one-3-carboxylic acid) is one of a new series of recently synthesized 1,8-naphthyridine derivatives with antibacterial properties.

It has considerable *in vitro* activity against pathogenic Gram-negative bacteria and is a valuable therapeutic agent in urinary infections, particularly those due to *E. coli*, other coliforms, and *Proteus* organisms.

The range of activity *in vitro* assessed by a disk technique against 133 organisms compares favourably with other drugs commonly used. Forty-six out of 53 *E. coli* and other coliforms and 15 out of 20 *Proteus* organisms were found to be sensitive, while only 8 out of 24 *K. aerogenes* were sensitive.

The only toxic reactions in 120 patients treated with the drug were a transient skin rash in one, and nausea in a patient with a duodenal ulcer.

A short clinical trial was done in 52 patients with 63 infecting urinary organisms. In 26 acute infections which had not previously been treated there were three failures, two due to insensitive organisms. In 26 other patients, some with an underlying cause for the infection and all previously treated with other drugs, there was a clinical cure in 20, but a bacteriological cure in only 14.

Nalidixic acid should prove of particular value against *Proteus* organisms.

We are grateful to Dr. Andrew MacFarlane, of the Mansfield Area Pathology Service, for his help. We wish to thank the Clinical Evaluation Unit of the Bayer Products Co. for supplies

of nalidixic acid, and Mr. P. J. O'Donnell of that unit for information.

REFERENCES

- Barlow, A. M. (1963). *Brit. med. J.*, **2**, 1308.
 Buchbinder, M., Webb, J. C., Anderson, L. V., and McCabe, W. R. (1963). *Antimicrobial Agents and Chemotherapy—1962*, p. 308. American Society for Microbiology, Michigan.
 Carroll, G. (1963). *J. Urol. (Baltimore)*, **90**, 476.
 Garrod, L. P., Shooter, R. A., and Curwen, M. P. (1954). *Brit. med. J.*, **2**, 1003.
 Jameson, R. M., and Swinney, J. (1963). *Brit. J. Urol.*, **35**, 122.
 Kass, E. H. (1955). *Amer. J. Med.*, **18**, 764.
 Leshner, G. Y., Froelich, E. J., Gruett, M. D., Bailey, J. H., and Brundage, R. P. (1962). *J. med. pharm. Chem.*, **5**, 1063.
 Lishman, I. V., and Swinney, J. (1963). *Brit. J. Urol.*, **35**, 116.
 McChesney, E. W., Froelich, E. J., Leshner, G. Y., Crain, A. V. R., and Rosi, D. (1963). *Toxicol. appl. Pharmacol.* In press.
 Rhoads, P. S., Billings, C. E., and O'Connor, V. J. (1952). *J. Amer. med. Ass.*, **148**, 165.
 Slade, D. A. (1963). *Brit. J. Urol.* In press.
 Thompson, R. E. M., and Rae, J. (1963). *Ibid.* In press.
 Turck, M., Browder, A. A., Lindemeyer, R. I., Brown, N. K., Anderson, K. N., and Petersdorf, R. G. (1962). *New Engl. J. Med.*, **267**, 999.
 Waterworth, P. (1962). *J. med. Lab. Technol.*, **19**, 163.

Preliminary Communications

Deoxyribonucleic Acid Content of Epidermal In-situ Carcinomas

Previous observations indicated that most carcinomas and sarcomas of man were composed of cells characterized by abnormal deoxyribonucleic acid (D.N.A.) values (Bader, 1959; Richards and Atkin, 1959; Stich *et al.*, 1960; Stich and Steele, 1962), whereas cells of several benign tumours had a normal diploid D.N.A. content (Stich *et al.*, 1960; Stich, 1962). Apart from these two types of neoplasm there is an epithelial proliferative lesion known as in-situ carcinoma, the true nature and eventual development of which is still ill-defined. It appeared, therefore, of interest to study the D.N.A. content in the neoplastic cells of these lesions in man, and to determine whether their D.N.A. content resembles that of benign or malignant cells.

The surgically excised lesions were fixed in 10% neutral formalin, embedded in paraffin, sectioned at 8 μ , and stained with Feulgen's technique. Patau's (1952) two wavelength method with Mendelsohn's (1958) calculation tables was used to estimate the relative amount of D.N.A. About 75 metaphase plates and 30 lymphocytes were measured per sample. The D.N.A. content of lymphocytes was used to give the diploid D.N.A. value of a normal cell. The symbol C denotes the amount of D.N.A. corresponding to a nucleus with haploid chromosome number. Thus a nucleus belonging to a diploid class has a D.N.A. amount of 2C in telophase nuclei and 4C in metaphase plates (Stich and Steele, 1962). The incidence of mitotic irregularities was estimated on Feulgen stained sections.

Metaphase plates of normal epidermis and of in-situ carcinomas were used for the D.N.A. measurements because at this particular stage of the mitotic cycle no D.N.A.

Distribution of D.N.A. values in normal skin and in-situ carcinomas.

synthesis occurs. Thus deviations in chromosome number from a diploid complement will be reflected in either an increased or a decreased D.N.A. content. All normal epidermal cells examined showed D.N.A. values which fell into the expected D.N.A. class for diploid cells (see Fig.). In contrast, the cells of in-situ carcinomas showed a great variability in the D.N.A. values. In eight cases they extended from the diploid to a hypertetraploid range (Cases 418, 2446, 6496, 5552, 57, 1859, 495, and 4778), in three cases (1407, 5669, and 3990) to hyperoctoploid range, and in one case (2224) to a sub-sixteen ploid range. These findings show that the lesions examined consist of heterogeneous cell populations in regard to their D.N.A. values. A dominant cell type characterized by a particular aneuploid D.N.A. value was not observed. In this regard, the lesion examined seems to differ from most invasive carcinomas, which, while showing a heterogeneous composition, have usually a particular stemline (Bader, 1959; Stich, 1960; Stich *et al.*, 1960; Stich and Steele, 1962).

Another consistent finding in the cases of in-situ carcinoma examined was a relatively high incidence of mitotic irregularities, including chromosome bridges, fragments, and multipolar mitosis (see Table). It is very likely

Incidence of Mitotic Irregularities in Normal Skin and Carcinomas In-situ

Specimens	Case No.	No. of Cells Counted	Mitotic Irregularities (%)			
			Bridges	Frag-ments	Multipolar Mitosis	Total
Normal skin	236-55	250				2.1
	2092-55	265				3.2
	2133-55	260				1.9
	2224-56	255				2.6
	5552-58	275				1.4
	4940-62	66				3.0
In-situ carcinomas	3990-56	85	14.4	5.8	1.1	21.3
	2224-57	91	23.0	3.2	0.0	26.2
	495-57	109	22.0	1.8	0.9	24.7
	4778-58	66	19.6	7.5	0.0	27.1
	418-58	84	13.0	4.7	1.1	18.8
	1859-58	65	6.1	4.6	0.0	10.7
	5552-59	110	16.3	3.6	0.0	19.9
	1407-61	108	34.2	2.7	0.0	36.9
	5669-62	72	16.6	2.7	1.3	20.6
	2446-62	93	16.1	6.4	0.0	22.5
	6496-62	84	15.4	5.9	1.2	22.5
	57-63	80	10.0	7.5	2.5	20.0

that these mitotic abnormalities led to the formation of the observed variation in D.N.A. values.

The occurrence of a genetically heterogeneous population and the absence of a dominant aneuploid stemline in the in-situ carcinomas resembles the pattern found in pre-neoplastic livers of rats fed various liver carcinogens (Stich, 1960, 1962) and to the pre-leukaemic lesions in mice injected with 7,12-dimethylbenz(α)anthracene (D.M.B.A.) (Joneja and Stich, 1963). The hepatomas and thymic lymphosarcomas consisting of cells with abnormal chromosome complement originated from this genetically heterogeneous population. It seems possible, therefore, that the heterogeneous cell population of the in-situ