Preliminary Communications

Some Endocrinological Effects of a Dithiocarbamoylhydrazine

Paget *et al.* (1961) described some effects of dithiocarbamoylhydrazines on various animals and suggested that these drugs inhibited pituitary gonadotrophic function. Among the most potent was 1- α -methylallylthiocarbamoyl-2-methylthiocarbamoylhydrazine (ICI 33828). The anti-gonadotrophic action of this drug on animals has been confirmed (Brown, 1963a), while experiments on rats and mice have suggested that it may also act as an inhibitor of pituitary thyrotrophic function (Tulloch *et al.*, 1963). If this drug does inhibit pituitary trophic hormones it may be clinically useful in situations where it is desirable to reduce the activity of the anterior pituitary gland. This paper is a report of our experience of the effects of this drug on the endocrine function of post-menopausal women, some of whom had metastatic breast cancer, and of male and female thyrotoxic patients with varying degrees of exophthalmos.

SUBJECTS AND METHODS

Clinical details of the 10 patients selected for treatment with ICI 33828 are shown in Tables I and II. The co-operation of the patients was obtained after the nature and objects of the investigation had been explained to them.

TABLE I.—Clinical Features and Treatment of Patients with Normal Endocrine Function

Case No.	Sex	Age in Years	Disease	ICI 33828	
				Daily Dose (mg.)	Days of Treatment
1	F	50	Carcinoma of breast	300	10
2	F	69	Uterine prolapse	300	10
3	F	56	Carcinoma of breast	300	5
4	F	68	" " "	100	5

TABLE II.—Clinical Features and Treatment of Thyrotoxic Patients

Case No.	Sex	Age in Years	Diagnostic Index	Exophthalmos	ICI 33828	
					Daily Dose (mg.)	Days of Treatment
5	F	62	+11	None	300	4
6	M	58	+39	Moderate	300	5
7	F	31	+24	"	300	5
8	M	40	+36	Severe, with ophthalmoplegia	300	10
9	F	27	+25	Mild	100	5
10	F	74	+21	None	100	10

Four of the patients (all female) had normal endocrine function, and six (four females and two males) were thyrotoxic. This diagnosis was made on the basis of the clinical diagnostic index of Crooks *et al.* (1959), radioactive iodine studies, and measurement of the serum protein-bound iodine (P.B.I.). The drug was given orally in the form of film-coated tablets in doses of 50 mg. twice or 100 mg. three times a day (see Tables I and II). The patients were in hospital, and inquiry was made daily for any symptoms of side-effects. In most cases the blood-pressure was measured daily in the supine and erect positions, and white blood counts together with a number of standard liver-function tests, usually including estimation of the bromsulphalein clearance rate, were done weekly.

Gonadotrophins were measured as follicle-stimulating hormone (F.S.H.) and total gonadotrophin by two different

methods of assay (Brown, 1963b), and the results were expressed as milligram equivalents of the international reference preparation for human menopausal gonadotrophin (IRP). Urinary oestriol was measured as described by Kloppe and Wilson (1962), and pregnanediol by the method of Kloppe *et al.* (1955).

Thyroid function was assessed by measuring the uptake of ^{131}I by the thyroid gland four hours after an oral tracer dose of carrier-free Na^{131}I . The short half-life of this isotope (2.3 hours) made it possible to repeat the test at daily intervals. Serum protein-bound iodine was measured by the method of Farrell and Richmond (1961) and plasma corticosteroids by that of Petersen *et al.* (1957).

RESULTS

Three post-menopausal women with normal endocrine function (Cases 1 to 3) were given 100 mg. of ICI 33828 three times a day, the results being summarized in Figs 1 to 3. One showed a definite fall in gonadotrophin excretion as early as the second 48 hours of treatment, and this was accompanied by a reduction in urinary oestriol excretion from a high initial level (10.8 and 21.9 $\mu\text{g./day}$) to less than 4 $\mu\text{g./day}$ (Fig. 1). The other two patients

FIG. 1.—Results in Case 1.

however, showed no evidence of any real change in gonadotrophin excretion during courses of treatment lasting for 10 and 5 days respectively (Figs. 2 and 3). Pretreatment levels of urinary steroids in these patients were not high, and there was only a doubtful suggestion of a fall in the level of excretion in the patient on the longer course of treatment (Fig. 2). All three patients showed a fall in the four-hour uptake of ^{131}I by the thyroid gland during the period of administration of the drug, with subsequent recovery to control levels or higher on stopping the treatment, and in all cases the serum P.B.I. level fell.

The fourth post-menopausal woman with normal endocrine function (Case 4) received 50 mg. of ICI 33828 twice daily for a period of five days and showed a fall in total gonadotrophin excretion from initial values of 36 and 44 mg. IRP/day to 6.5 mg./day by the fifth day of treatment. Measurement of F.S.H. was not made.

Urinary gonadotrophins were measured in one thyrotoxic patient (Case 5) and the results are shown in Fig. 4. There was a fall in gonadotrophin excretion on the third

measurements. Two other thyrotoxic patients (whose results are not illustrated) were treated with 100 mg. daily for periods of 5 and 10 days and showed no change in iodine uptake. In none of the thyrotoxic patients was there any convincing fall in serum P.B.I. level.

Plasma corticosteroids did not show a consistent response in those patients, both euthyroid and thyrotoxic, in whom measurements were made.

Side-effects.—Four patients complained of drowsiness, including one patient who received only 100 mg./day. Several patients also complained of nausea, and this was most severe and persistent in the man with exophthalmic ophthalmoplegia and in one thyrotoxic woman. In the latter patient it was associated with abdominal pain and vomiting, which suggested a recurrence of symptoms from

FIG. 2.—Results in Case 2.

FIG. 3.—Results in Case 3.

day of treatment with 300 mg. of ICI 33828, and it returned to pretreatment levels within 48 hours of stopping the drug, as was the case also in the euthyroid patients.

The effect of 300 mg. of ICI 33828 daily on the thyroid function of the thyrotoxic patients is illustrated in Figs. 4 and 5. There was a reduction in iodine uptake in three of the four thyrotoxic patients treated with this dose. The fourth patient, however—a man with exophthalmic ophthalmoplegia—received this dose for 10 days without any depression of uptake or change in exophthalmometric

FIG. 4.—Results in Case 5.

FIG. 5.—Radioiodine uptake by the thyroid in thyrotoxic patients (Cases 6-8) treated with 300 mg. ICI 33828 daily. Results in Case 5 are shown in Fig. 4.

which she had suffered previously and which had been shown to be associated with a duodenal ulcer. Treatment with ICI 33828 was stopped after only four days and her dyspepsia subsided. In most cases the blood-pressure was recorded daily, since a patient not included in the present series developed postural hypotension during treatment with the drug. Later, however, this patient was found to suffer from similar attacks without treatment, and it seems unlikely that the treatment was responsible. Postural hypotension was not found in any of the patients in the present series, and no signs of leucopenia or disturbed liver function were detected.

DISCUSSION

The administration of ICI 33828 in doses of 100 or 300 mg./day caused a considerable reduction in the excretion of gonadotrophin in three out of five post-menopausal women. This finding is in agreement with that of Bell *et al.* (1962a), who showed a reduction in total gonadotrophin excretion in two post-menopausal women treated with doses of 125 and 600 mg./day. In our patients two contrasting methods of assay demonstrated the same pattern of fall, and suggested that the gonadotrophic hormones were equally affected. The concentration of these hormones in the pituitary gland of male rats is believed to fall after treatment with ICI 33828 (Brown, 1963a), and if the mechanism of action is similar in man and the rat the combined evidence suggests that this drug can reduce the formation of pituitary gonadotrophins. The present data indicate, however, that reduction in urinary gonadotrophin in post-menopausal women treated with doses of 300 mg. daily is not invariable. It is also interesting to note that Bell *et al.* (1962b) have shown that the drug did not appear to affect gonadotrophin excretion in two premenopausal women, at least as this was measured by a single non-specific method of assay, although it suppressed ovulation and ovarian steroid secretion.

The fall in urinary oestriol excretion occurring in our post-menopausal patient who started with uncharacteristically high oestriol levels was of the type demonstrated in premenopausal women (Bell *et al.*, 1962b). Since her gonadotrophin excretion was also reduced, it would be reasonable to suggest that these two responses were causally related, but it should be noted that, while the excretion of gonadotrophin recovered quickly after stopping the drug, the oestriol level had not risen again after eight days.

The effect of the drug on thyroid function in the euthyroid patients was unequivocal in that it reduced both radioiodine uptake by the thyroid and serum P.B.I. These changes might be due either to interference with the production or metabolism of thyroid hormone or to a direct effect on pituitary thyrotrophic function. There is some evidence to suggest that the drug acts at the pituitary level in animals (Tulloch *et al.*, 1963), and we are attempting to test the various possibilities in man.

All but one of the thyrotoxic patients on 300 mg. of ICI 33828 daily showed a similar reduction in thyroid iodine uptake. The male patient with malignant exophthalmos (Case 8) showed no fall in uptake on this dose, and there was a similar lack of response in two women who were given the lower dose (Cases 9 and 10). There was no obvious correlation between any of the features of the thyrotoxic state and the presence or absence of a response

to ICI 33828. Since the number of patients treated is small, it is not yet possible to say whether there is any real difference in the effect of the drug on thyrotoxic and euthyroid patients.

The drug appears to reduce certain anterior pituitary functions, and consequently is a pharmacological tool of great interest. At present it has no place in practical therapeutics, but its properties should be explored with possible applications in mind.

SUMMARY

ICI 33828 caused a distinct fall in urinary gonadotrophin excretion in three out of five post-menopausal patients. There was no evidence of a qualitative change in the gonadotrophin excreted.

Oestriol excretion was unusually high in one of the post-menopausal women with breast cancer. It fell to normal post-menopausal levels during treatment.

Plasma corticoids were measured in four patients receiving the drug and showed no consistent effect.

Oral doses of 300 mg. daily reduced thyroid iodine uptake and serum P.B.I. in three euthyroid patients. The same dose reduced iodine uptake in three out of four thyrotoxic patients, while 100 mg. daily failed to suppress uptake in two further thyrotoxic subjects.

The most frequent side-effect was nausea, and in one case this was severe and associated with abdominal discomfort and vomiting.

We wish to thank Professor Sir Dugald Baird and Professor A. G. Macgregor, in whose wards the patients were studied. The work was supported by a grant to one of us (A. R. T.) from the Secretary of State for Scotland on the recommendation of the Advisory Committee on Medical Research.

P. S. BROWN, B.A., B.M., M.R.C.P.,
Scientific Staff, M.R.C.

J. CROOKS, M.D., F.R.C.P.Ed., M.R.C.P., M.R.C.P.Glasg.,
Senior Lecturer.

A. I. KLOPPER, M.B., Ch.B., B.Sc., Ph.D., M.R.C.O.G.,
Scientific Staff, M.R.C.

A. R. THORBURN, B.Sc.,
Research Assistant.

M. I. TULLOCH, B.Sc.,
Research Assistant.

Departments of Materia Medica and Therapeutics,
and Medical Physics, University of Aberdeen.
Obstetric Medicine Research Unit (M.R.C.),
University of Aberdeen.

REFERENCES

- Bell, E. T., Brown, J. B., Fotherby, K., and Loraine, J. A. (1962a). *Lancet*, **2**, 528.
— and Robson, J. S. (1962b). *J. Endocr.*, **25**, 221.
Brown, P. S. (1963a). *Ibid.*, **26**, 425.
— (1963b). *Ibid.*, **25**, 427.
Crooks, J., Murray, I. P. C., and Wayne, E. J. (1959). *Quart. J. Med.*, **28**, 211.
Farrell, L. P., and Richmond, M. H. (1961). *Clin. chim. Acta*, **6**, 620.
Klopper, A., Michie, E. A., and Brown, J. B. (1955). *J. Endocr.*, **12**, 209.
— and Wilson, G. R. (1962). *J. Obstet. Gynaec. Brit. Cwlth.*, **69**, 533.
Paget, G. E., Walpole, A. L., and Richardson, D. N. (1961). *Nature (Lond.)*, **192**, 1191.
Petersen, R. E., Karrer, A., and Guerra, S. L. (1957). *Analyt. Chem.*, **29**, 144.
Tulloch, M. I., Crooks, J., and Brown, P. S. (1963). *Nature (Lond.)*, **199**, 288.