Papers and Originals**ROLE OF THE THYMUS IN IMMUNITY***

BY

J. F. A. P. MILLER, M.B., B.S., B.Sc.(Med.), Ph.D.*Chester Beatty Research Institute, Institute of Cancer Research: Royal Cancer Hospital, London*

[WITH SPECIAL PLATE]

The thymus has been an enigmatic organ for centuries. Some of the ancient Greeks considered it to be the seat of the soul. Galen, on the other hand, thought that it acted as a cushion to protect the great vessels from the bony sternum. Since then, contentions and controversies have involved most of the questions regarding its embryology, anatomy, physiology, and pathology. Indeed, "has it yet fallen to the lot of any writer upon the thymus to write the truth and be believed?" (Beard, 1902).

Development and Structure

The thymus in mammals arises as paired structures from the endoderm of the third and fourth branchial clefts. Its epithelial character is plainly evident until the end of the second month of intrauterine life in the human. At that time the small thymic cells begin to appear.

There has been much controversy concerning the manner in which the embryonic epithelial thymus structure comes to assume its lymphoid appearance. Some, following Hammar (1908) and Maximow (1909), contended that the small thymic cells were of mesodermal origin, having been derived from mesenchymal cells which had migrated into the epithelial rudiment of the thymus. Others, following Maurer (1885) and Stöhr (1906), maintained that the thymic round cells arose directly from the reticular epithelium and believed also that they had the potentiality of reversion to their original epithelial form, a view also adopted by Burnet and MacKay (1962). The controversy about the origin of thymic lymphocytes appears to have been settled by Auerbach (1961), who used elegant tissue-culture techniques to analyse the morphogenetic events in thymus development. Thymic rudiments obtained from 12-day mouse embryos were separated into epithelial and mesenchymal components by the use of trypsin. It was shown that the epithelial component of the rudiment was capable of forming the lymphocytes of the thymus, the mesenchyme providing the initial inductive stimulus and serving to furnish the stromal elements of the gland. Neither host migrating cells, nor thymic mesoderm, nor generalized mesenchyme appeared to contribute significantly to the initial lymphoid cell population of the thymus.

In size the thymus increases more or less in step with the general growth until about puberty, when it slowly begins to atrophy. Its phase of most rapid growth therefore occurs in early life.

The fully developed thymus is incompletely divided into lobules composed of a central core or medulla and an external part or cortex. It has two main types of cells: the lymphocytes and the reticular cells. The lymphocytes are very closely aggregated in the cortex, where they overshadow the reticular cells. Most of the cortical lymphocytes are small round cells with a relatively great nuclear/cytoplasmic ratio. Their mitotic rate is high. On morphological grounds alone, it is not possible to find any

significant differences between lymphocytes found in the thymus and those present elsewhere. The reticular cells are most prominent in the medulla. They are large cells with ill-defined outlines and a large vesicular nucleus poor in chromatin content. Most of them are epithelial in nature, having originated from the endoderm of the branchial clefts. Some of them are typical reticulo-endothelial cells or macrophages.

The medulla is noteworthy for the presence of Hassall's corpuscles. These may consist of single hypertrophic reticular cells or groups of reticular cells which may appear active, irregularly polygonal or flattened, hyalinized and concentrically arranged about a central core of nuclear debris (Fig. 1, Special Plate). They have been variously interpreted as remnants of the original epithelial anlage, as structures derived from connective tissue, from the endothelium of blood-vessels, or as structures developed from reticular cells. Modern opinion favours the latter view (Sainte-Marie and Leblond, 1958).

The main morphological differences between the thymus cortex and medulla are summarized in Table I. Much work is required to appreciate fully the functional significance of these differences.

TABLE I.—*Morphological Differences Between Thymus Cortex and Medulla*

	Cortex	Medulla
Lymphocyte reticular-cell ratio	100	20
Lymphocyte mitosis	1.3-2.4 per high-power field	0.4 per high-power field
Hassall's corpuscles	Absent	Present
Main blood-vessels	Capillary	Venous spaces

The Thymus as a Lymphoid Organ

Because of its high content of lymphocytes the thymus has for long been regarded as part of the lymphoid system. It differs, however, from other lymphoid structures in several outstanding respects (Table II).

TABLE II.—*Morphological Differences Between Thymus and Other Lymphoid Organs*

	Thymus	Non-thymic Lymphoid Organs
Epithelial-reticular cells	Present	Absent
Hassall's corpuscles	Present	Absent
Mitotic activity	0.7%	0.05-0.1%
Germinal centres	Absent	Present
Plasma cells	Few (do not increase after antigenic stimulation)	Present (increase after antigenic stimulation)

Mitotic indices of thymic lymphoid cells are about seven times higher than those of other lymphoid cells. Even with advancing age, lymphocyte production has been shown to take place with almost the same intensity as in the young thymus and is always greater than it is in lymph nodes (Andreassen and Christensen, 1949). Thymus lymphopoiesis is not affected by antigenic stimulation, resection of other lymphoid organs, partial thymectomy, or the presence of

*Based on a lecture delivered to the University of Aberdeen on May 16, 1963.

one or more thymus grafts. Yet thymus lymphocytes are subjected to the same general hormonal regulators as other lymphocytes. These facts therefore suggest that the strong proliferative stimulus for thymic lymphoid cells must arise intrinsically in the thymus itself. Further, the fact that there is no regeneration of a thymus fragment left after partial thymectomy and that there is no increase in mitotic activity in the fragment suggests that the mitotic stimulus must be associated with certain structures widely distributed or scattered throughout the thymus. Indeed, there seems to be a physical association between mitotic activity in the cortex and specialized periodic-acid-Schiff (P.A.S.)-staining reticular cells (Metcalf and Ishidate, 1962). Mitoses are significantly more frequent in lymphocytes adjacent to P.A.S. cells than in lymphocytes elsewhere, and this association is even more obvious when mitotic activity is increased as in pre-leukaemic thymuses, thymuses regenerating after cortisone-induced atrophy, or thymuses in adrenalectomized animals. The uniqueness of thymus tissue as regards intense lymphopoietic activity may then rest on the presence of such P.A.S. cells in the thymus cortex.

What is the fate of lymphocytes produced in the thymus cortex? According to some (Saint-Marie and Leblond, 1958), cortex-formed small lymphocytes would migrate through the medulla in order to enter the circulation. Do thymus lymphocytes in fact reach the circulation? This is another controversy that remains unsettled. Much circumstantial evidence makes it appear that lymphocytes are released from the thymus and destined for other places. On the other hand, there is no *direct*, unequivocal evidence to establish this. In normal animals, even when the thymus is growing, there are many pycnotic lymphocytes in the organ. Theoretically, all the lymphocytes produced there could die locally without ever entering the circulation. Thus, while there is unequivocal evidence that the thymus produces lymphocytes, there is as yet no direct evidence to show that such lymphocytes are destined for other places.

Immunological Reactivity

Immunological reactivity is usually associated with lymphoid organs such as spleen and lymph nodes. The small lymphocyte is the commonest type of lymphoid cell present in such organs and in the blood and lymph, and it has recently been proved to be an immunologically competent cell—a cell which can initiate an immunological reaction when given the appropriate stimulus (Gowans *et al.*, 1962). Although the thymus is rich in small lymphocytes it does not show production of antibody, it does not respond to antigenic stimuli by producing plasma cells or germinal centres. It therefore behaves completely differently from other lymphoid structures in an immunized animal. Does this imply that there are no cells in the thymus capable of taking part in immune reactions? Whether or not thymus lymphocytes are immunologically

competent is the subject of yet another controversy. Some of the experiments which have been devised to settle this question are briefly described below.

1. Direct injection of antigen into the thymus is associated with a plasmacellular reaction and antibody production within the thymus. It was suggested that the failure of the thymus to react to circulating antigen was due to a barrier between blood and thymus parenchyma preventing the entry of antigen into the thymus (Marshall and White, 1961). Such experiments, however, do not exclude the entry into the thymus of extrinsic (non-thymic) lymphoid cells which may have accumulated there as a result of the injury inflicted by the intrathymic injection.

2. Antibody has been detected in thymus tissue transplanted to the anterior chamber of the eye of the mouse (Stoner and Hale, 1955).

3. Dissociated mouse-thymus cells can mount an immune attack (graft-versus-host reaction) against a foreign host but less effectively than mouse-lymph-node cells. Rat-thymus cells cannot produce such a reaction (Silvers, 1962). In the mouse, it is difficult to separate from the thymus the tiny thoracic lymph nodes which are embedded in or close to its capsule. Graft-versus-host disease in such experiments could have been caused by contaminating lymph-node cells—and this may explain why so many more cells were required to produce an effect with thymus preparations than with lymph-node suspensions.

4. In rabbits and guinea-pigs, thymus cells taken from presensitized donors or from normal donors and sensitized *in vitro* with Shigella antigens have produced little or no antibody in appropriate recipients, in complete contrast to non-thymic lymphocytes (Harris *et al.*, 1955; Dixon *et al.*, 1957).

The question is therefore still open whether there are cells in the thymus with immunological capabilities. If the thymus is at all responsible for maintaining immunological reactivity as a whole it is possible that thymectomy would reduce such reactivity.

The Thymectomized Animal

Thymectomy has been performed in many species and strains of *young adult* rodents, and has been without effect on growth rate, breeding behaviour, susceptibility to common infections, or longevity. It has, however, been associated with a fall in circulating lymphocytes, in both blood and thoracic duct, to about 30 to 40% below normal levels, and a fall in the weight of the spleen and lymph nodes by about 25% (Bierring, 1960; Metcalf, 1960). There is, however, no significant diminution in the immunological capabilities of the animal thymectomized in adult life (Fichtelius *et al.*, 1961), suggesting that "the thymus gland does not participate in the control of the immune response" (MacLean *et al.*, 1957). Since, however, the thymus is the major lymphoid organ in the newborn, other lymphoid structures being then poorly developed, it is possible that thymectomy of the newborn, in contrast to thymectomy of the adult, may be associated with more severe defects.

Mice thymectomized at birth develop normally for the first few weeks. After weaning, however, many die from a wasting disease characterized clinically by loss of weight, hunched posture, ruffled fur, diarrhoea, and death within a week (Fig. A). The earlier in life thymectomy is performed the higher the incidence of this syndrome. Some strains develop the syndrome as early as 4 weeks, others as late as 4 months. Thymectomy after one week of age is no longer associated with the development of this syndrome (Miller, 1963a). A similar syndrome has been described in rats (Janković *et al.*, 1962) and hamsters (Sherman

FIG. A.—Wasting disease: mouse on right shows signs of wasting disease 8 weeks after thymectomy at birth. Mouse on left is a litter mate also thymectomized at birth but grafted at 7 days of age with thymus tissue in the subcutaneous region of the axilla.

and Dameshek, 1963) thymectomized at birth. The wasting disease following neonatal thymectomy has obvious resemblances to the syndromes seen in graft-versus-host conditions, such as runt disease in infant mice, secondary disease in radiation chimeras, F_1 hybrid disease, and parabiotic intoxication.

The neonatally thymectomized mouse characteristically shows a persistently low peripheral blood-lymphocyte level (50 to 90% below normal) (Miller, 1961, 1963a) and a marked depletion of lymphocytes in lymphoid tissues (Figs. 2-6, Special Plate). The follicular structures of the lymph nodes has disappeared and germinal centres are absent (Miller, 1961, 1962a, 1962b). Spaces normally occupied by lymphoid follicles show reticulo-endothelial cells, histiocytes, macrophages with particles of cells and haemosiderin, and only a few scattered lymphocytes, most of which are immature. There is a marked deficiency of small lymphocytes and a moderate-to-marked reticulo-endothelial cell hyperplasia. In many lymph nodes there are only a few plasma cells or plasma-cell precursors. Follicles are usually present in the spleen although they are reduced in size and numbers. Peyer's patches are considerably reduced in size and often missing. Small lymphoid aggregates may be found in the submucosa of the ileum, the majority of cells consisting of reticulo-endothelial cells or immature lymphoid cells. Plasma cells may be found in the lamina propria, but are usually reduced in numbers. In neonatally thymectomized rats the characteristic feature is a deficiency of small lymphocytes: plasma cells and germinal centres are normal (Waksman *et al.*, 1962).

Small discrete necrotic patches are seen in the liver of about 50% of mice dying of wasting disease. They are acute necrotic areas, with no specific location in the lobules, and contain clusters of neutrophils and small numbers of macrophages. There is a general increase in the size and number of Kupffer cells. In the bone-marrow there is no apparent abnormality or any deficiency of lymphocyte-like cells.

The capacity of neonatally thymectomized mice to produce serum antibodies has been tested (Miller, 1963a). These studies have shown that 50 to 80% of such mice fail to produce detectable levels of antibodies to the H antigen of *Salmonella typhi*, to killed influenza virus, and to sheep erythrocytes. Similar deficiencies have been reported in rats (Janković *et al.*, 1962) and rabbits (Archer *et al.*, 1962) thymectomized at birth.

Mice subjected to a sham-operation at birth or to splenectomy at birth can later reject foreign skin grafts just as normal mice (Miller, 1962a). However, more than 70% of mice thymectomized at birth permanently fail to reject skin grafts, not only from donors closely related to the hosts (Miller, 1961) but also from distantly related donors such as rats (Miller, 1962a, 1962b). Such mice fail to reject the skin when grafted well before the clinical onset of wasting disease (Fig. B) and usually keep the graft until death. Thymectomy after 1 week of age is no longer associated with significant prolongation of survival of foreign skin grafts (Miller, 1962a). Temporary suppression of homograft immunity has also been reported in rats thymectomized at birth (Arnason *et al.*, 1962).

Thymectomy of the newborn mouse is therefore associated with (1) a fatal wasting disease occurring after weaning; (2) a marked diminution in the population of small lymphocytes in blood and tissues; (3) a diminished capacity to produce serum antibodies to some antigens; (4) a permanent failure of the capacity to reject foreign tissue grafts.

How can these results be explained? One might postulate that, in the newborn, cells with immunological potential are few or non-existent in non-thymus lymphoid organs. As the animal develops, these lymphoid organs receive from the thymus either a cellular or a non-cellular contribution which endows them with their full complement of immunological competent cells. Thymectomy in adult life would no longer be associated with severe defects, as

FIG. B.—Eight-weeks-old mouse thymectomized at birth and grafted at four weeks with skin from a mouse of a foreign strain. Note luxuriant tuft of hair on graft.

the other lymphoid organs can now carry on independently of the thymus. If, however, their immunological potential were to be destroyed or depleted, would the thymus again be required to restore this potential?

Experiments were therefore set up with mice thymectomized in adult life and exposed to total body irradiation. The results showed: (1) A marked diminution in lymphocytes, particularly small lymphocytes, in both blood and tissues in thymectomized irradiated mice, but no such

FIG. C.—Six-months-old mouse thymectomized and irradiated at 6 weeks of age and grafted with foreign skin at 10 weeks. Note greying of hair after irradiation and luxuriant tuft of hair on graft.

diminution in non-thymectomized irradiated controls two months after irradiation; (2) a failure to produce either a primary or a secondary response to sheep erythrocytes and to reject foreign skin grafts (Fig. C) 4 to 10 weeks after irradiation, but no such deficiencies in controls (Miller, 1962c; Miller *et al.*, 1963).

Clearly, therefore, the recovery of lymphopoietic and immune functions in an irradiated *adult* mouse is still dependent on a thymus-controlled mechanism. The adult thymus may thus still be essential to re-establish immunological mechanisms when the immune apparatus has been injured.

Protection Against the Effects of Thymectomy

A number of methods have been tested for their ability to protect against the effects of neonatal thymectomy.

1. A suspension of lymph-node cells or spleen from normal adult mice was found to be effective in preventing wasting disease and restoring lymphocyte counts and immune mechanisms when injected into mice of the same strain within a week after thymectomy (Miller, 1963c). Similarly, presensitized lymphoid cells were capable of breaking down an established skin graft on a neonatally thymectomized mouse (Miller, 1962a).

2. The effects of feeding or injecting thymus extracts to thymectomized animals have given equivocal results. Metcalf (1958) claimed that saline extracts of thymuses produced a lymphocytosis on injection into baby mice with intact thymuses. He suggested that the thymus elaborated a specific factor (L.S.F.) which would stimulate maturation or division of primitive lymphocytes into mature small lymphocytes. These claims have not been confirmed adequately (Duplan *et al.*, 1962). Saline extracts of young thymuses given repeatedly to mice thymectomized at birth failed to reduce the incidence of wasting disease, to enable the mice to reject foreign skin grafts, and to restore lymphocyte counts to normal (Miller, 1963c).

3. Mice thymectomized at birth and grafted within a week with an intact thymus from a newborn donor developed normally, had a normal life-span, normal lymphocyte populations in blood and tissues, and could reject foreign skin grafts as normal mice (Miller, 1961, 1962a, 1962b). When the donor of the thymus was of the same strain as the host, all foreign grafts could be rejected. When the donor was of a different strain the host usually showed some tolerance to grafts of thymus donor-type skin, but not to third-party skin grafts.

The protection afforded by thymus implants may be explained by assuming that the implant produces lymphocytes which populate the host and restore immune functions. If this is so, one would expect to find in the host lymphoid tissues cells with the cytogenetic characteristics of the thymus donor. Experiments were set up employing hosts whose cells were identifiable at metaphase by marker chromosomes. It was found that the majority of the cells multiplying in host lymphoid tissues and in the thymus implant itself were of host type and therefore could not have been contributed by the thymus donor (Miller, 1963a, 1963b, 1963c). There must therefore remain in the neonatally thymectomized mouse a population of cells which can acquire immunological competence, but only in the presence of thymus tissue. The thymus would thus exert its influence (a) either by producing a humoral factor inducing the maturation of such cells wherever they may be, or (b) by allowing the differentiation of these cells within its own environment: the cells would invade the implant (the lymphoid cells of which are known to degenerate after grafting), differentiate in it, and then colonize the lymphoid tissues where they could function immunologically.

Possible Role of the Thymus

Any current theory purporting to explain the role of the thymus in immunity must necessarily suffer from the lack of fundamental knowledge concerning many aspects of the physiology of lymphocytes in general and of the thymus in particular. For instance, we still do not know with any certainty whether many thymus lymphocytes actually leave the thymus, whether they can react immunologically either in or out of the thymus environment, whether they are the direct ancestors of all immunologically competent cells, and whether all small lymphocytes have immunological functions. Granted certain assumptions, one can offer two alternative hypotheses which are not mutually exclusive: a cellular hypothesis and a humoral hypothesis.

We have previously suggested that the thymus provides the most highly specialized environment for what might be termed the immunological education of cells (Miller, 1962c, 1963b, 1963c, 1963d). According to elective theories of immunity and tolerance, contact of antigen with a cell at a certain stage of its differentiation might eliminate in some way its potentiality of reacting to the antigen. Contact with the mature cell, on the other hand, would selectively induce clonal expansion of that cell. The thymus, having the greatest lymphopoietic activity of all the lymphoid tissues and being apparently shielded from the entry of extrinsic antigens, would thus seem to provide the ideal site for the differentiation and non-clonal expansion of cells with immunological potential. During ontogenesis, such cells might be directly derived from epithelial cells of the thymus rudiment (Auerbach, 1961). Thereafter, lymphoid precursor cells found elsewhere—for example, in bone-marrow—would have to migrate to the thymus for their education before they could function as immunologically competent cells in other tissues (Miller *et al.*, 1963). Such a migration might take place, for instance, after stress or irradiation. Hence this theory demands that all immunologically reactive cells must at some stage have passed through the thymus. To account for the negligible effects of thymectomy in the adult one must postulate either that such reactive cells are long-lived or that they can reproduce in a thymectomized animal.

A humoral factor elaborated specifically by the thymus, perhaps by the epithelial cells or by the specialized P.A.S. reticular cells, may conceivably act in one or more of the following ways:

1. It may allow the differentiation of lymphoid precursor cells to lymphocytes both in the thymus and elsewhere.
2. It may stimulate lymphopoiesis (Metcalf, 1958).
3. It may stimulate the recirculation of lymphocytes. Thus, after thymectomy of the adult, the lymphocyte content of the thoracic duct drops within a week to about 40% below normal levels. In the normal animal the percentage of newly formed cells in the thoracic duct is of the order of 1 to 2% per day. The drop after thymectomy cannot, therefore, be accounted for solely on the basis of a loss of newly formed cells.
4. It may endow non-thymic lymphocytes with the property of immunological competence. Recent experiments employing neonatally thymectomized mice bearing thymus tissue enclosed in millipore diffusion chambers strongly support this last possibility (Osoba and Miller, 1963).

Whatever hypothesis one may adopt, it is clear that the thymus has some role in establishing immunological reactivity. Does it play any part in the induction of immunological tolerance? Burnet (1962), for instance, suggested that it must be endowed with another function—"the elimination or inhibition of self-reacting clones." The fact that neonatally thymectomized mice bearing foreign thymus grafts are specifically tolerant of thymus-donor-type skin, but not of third-party skin (Miller, 1962b), suggests that some such function may be thymus-dependent. On the other hand, there is no reason to think that tolerance cannot develop outside the thymus. If we accept that antigen does not get into the thymus of an adult animal, the fact that such animals can be made tolerant suggests that lymphocytes outside the thymus environment can respond by becoming tolerant. Hence it is not quite clear whether the thymus plays any direct part in the induction of specific tolerance. On the other hand, it seems more likely that it plays a part in the breakdown of tolerance. The accelerated breakdown of immunological tolerance following total body irradiation presumably occurs as a result of the differentiation of new immunologically competent cells during the recovery phase (Makela and Nossal, 1962).

Since the thymus is essential for the recovery of immunological function after total body irradiation (Miller, 1962c), it seems likely that it is also responsible for the breakdown of tolerance under these circumstances. Further evidence for this comes from unpublished experiments (Miller, 1963e) which showed that tolerance breaks spontaneously in an intact animal but persists indefinitely after thymectomy. Is it possible, too, that the thymus is responsible for the breakdown of autotolerance, and hence for the development of autoimmune diseases? This will be referred to below.

Relationship to Thymic Syndromes in Man

The crucial role of the thymus in the development of the immune apparatus raises important questions regarding a possible relationship between the thymus and immunological abnormalities in man. Table III lists abnormalities of the thymus and associated clinical syndromes.

1. *Aplasia*.—Essential lymphocytopenia is a rare condition. It has a familial incidence and affects infants of both sexes. They are apparently normal at birth but fail to thrive. The syndrome is usually recognized at about the age of 3 months, when the presenting clinical features are wasting, diarrhoea, morbilliform rashes, bacterial and fungal infections, agammaglobulinaemia, and marked lymphopenia. The condition is fatal despite antibiotic therapy, blood transfusions, and gamma-globulin infusions. At necropsy there is marked atrophy of the lymphatic tissue and moderate to marked reticulo-endothelial hyperplasia. Lymph follicles and plasma cells are absent. The thymus is markedly atrophic and has few or no small lymphocytes and no Hassall's corpuscles. The primary cause of this condition may therefore be genetically determined agenesis of the thymus—a condition strikingly similar to that described above in mice thymectomized at birth.

2. *Hyperplasia with Germinal Centre Formation*.—The presence of an immunological type of lymphoid proliferation in the thymus—namely, germinal centre and plasma cell formation—is associated with some disturbances which are most probably autoimmune in nature. There is a strain of mice, NZB/BL, which has a genetically determined autoimmune haemolytic anaemia (Burnet and Holmes, 1962) and which characteristically shows in the thymus hyperplasia of the medulla, germinal centre formation, and accumulation of plasma cells and mast cells. Similar lesions occur in the thymus of patients with myasthenia gravis in whom both antinuclear factors and anti-muscle serum factors have been demonstrated (Strauss *et al.*, 1960; Strauss, 1963; White and Marshall, 1962). Hyperplastic thymus lesions have also been described in rheumatoid arthritis, thyrotoxicosis, and Addison's disease (Table III), in all of which circulating antibodies have been reported (Doniach and Roitt, 1962).

3. *Neoplasia*.—Thymus tumours are associated with some clinical disorders which characteristically show either primary

or secondary immunological disturbances. Some of these have conditions which are believed to be autoimmune in nature—for example, acute haemolytic anaemia, leucopenia, thrombocytopenia, thrombotic thrombocytopenic purpura. Furthermore, antinuclear factors have been described in others, such as myasthenia gravis (White and Marshall, 1962), myositis and myocarditis (Funkhouser, 1961), and aplastic anaemia (Holborow *et al.*, 1963). Reports of a clinical association between myasthenia gravis and systemic lupus erythematosus suggest the possibility of a common aetiology (Funkhouser, 1961; White and Marshall, 1962), but so far no mention has been made of the appearance of the thymus in patients with systemic lupus.

Patients with Hodgkin's disease have shown deficiencies in gamma-globulin, antibody formation, and homograft rejection (Geller, 1953; Good *et al.*, 1957; Miller *et al.*, 1961). The deficiencies in such patients are probably secondary to extreme lymphoid aplasia, which occurs either as a result of the widespread replacement of lymphoid tissue by granulomatous proliferations or owing to its destruction by cortisone or irradiation therapy.

Conclusion

The association between clinical immunological disorders and thymus lesions and the results obtained from experiments in mice certainly support the concept that the thymus has a key role in controlling immunological reactivity. Clearly, further work is required to evaluate its role in the initiation of autoimmune disease.

Figs. 2, 5, and 6 (Special Plate) are reproduced by permission of Professor Maisin, rédacteur-en-chef of *Acta Unio Internationalis Contra Cancrum*.

REFERENCES

- Andreasen, E., and Christensen, S. (1949). *Anat. Record*, **103**, 401.
 Archer, O. K., Pierce, J. C., Papermaster, B. W., and Good, R. A. (1962). *Nature (Lond.)*, **195**, 191.
 Arnason, B. G., Janković, B. D., Waksman, B. H., and Wennersten, C. (1962). *J. exp. Med.*, **116**, 177.
 Auerbach, R. (1961). *Develop. Biol.*, **3**, 336.
 Beard, J. (1902). *Zool. Jahrb. Jena, Abt. Anat.*, **17**, 403.
 Bierring, F. (1960). In *Haemopoiesis: Cell Production and its Regulation*, pp. 185–203. Ciba Symp. Churchill, London.
 Burnet, F. M. (1962). *Brit. med. J.*, **2**, 807.
 — and Holmes, M. C. (1962). *Nature (Lond.)*, **194**, 146.
 — and MacKay, I. R. (1962). *Lancet*, **2**, 1030.
 Castleman, B. (1955). In *Tumors of the Thymus Gland: Atlas of Tumor Pathology, Section V, Fascicle 19*. Armed Forces Inst. of Pathology, Washington, D.C.
 — (1960). In *Thymectomy for Myasthenia Gravis*, edited by H. R. Viets and R. S. Schwab. Thomas, Springfield.
 Davis, L. A., and McCreadie, S. R. (1962). *Amer. J. Roentgenol.*, **88**, 924.
 Dixon, F. J., Weigle, W. O., and Roberts, J. C. (1957). *J. Immunol.*, **78**, 56.
 Doniach, D., and Roitt, I. M. (1962). *Ann. Rev. Med.*, **13**, 213.
 Duplan, J. F., Foschi, G. V., and Manson, L. A. (1962). *Proc. Soc. exp. Biol. (N.Y.)*, **110**, 426.

TABLE III.—*Thymic Syndromes in Man*

Thymus Lesion	Associated Clinical Disorder	References
Aplasia	Hypogammaglobulinaemia Essential lymphocytopenia	Reviewed in Giltin <i>et al.</i> , 1959 Hitzig <i>et al.</i> , 1958; Hitzig and Willi, 1961
Hyperplasia with germinal centres	Myasthenia gravis Rheumatoid arthritis Thyrotoxicosis Addison's disease Acromegaly Hypogammaglobulinaemia Aplastic anaemia	Reviewed in Ringertz, 1951, and Castleman, 1955 Burnet and MacKay, 1962 McEachern and Parnell, 1948; Castleman, 1960 Good, 1954 Wintrobe, 1951
Neoplasia :	Hypogammaglobulinaemia (rarely hypergammaglobulinaemia)	Good and Varco, 1955; Martin <i>et al.</i> , 1956; Ramos, 1956; Lambie <i>et al.</i> , 1957; Soutter <i>et al.</i> , 1957; Gafni <i>et al.</i> , 1960.
Epithelial and lympho-epithelial	Haemopoietic insufficiency : Pure red-cell anaemia, pancytopenia, autoimmune haemolytic anaemia, thrombocytopenia, thrombotic thrombocytopenic purpura, leucopenia, pernicious anaemia Myasthenia gravis Myocarditis and myositis	Reviewed in Havard and Bodley Scott, 1960, and in Fisher, 1963 Reviewed in Ringertz, 1951, and Castleman, 1955 Mendelow and Jenkins, 1954; Waller <i>et al.</i> , 1957; Langston <i>et al.</i> , 1959; Funkhouser, 1961 Reviewed in Fisher, 1963 Thomson, 1955 Webster, 1961; Davis and McCreadie, 1962
Small-cell carcinoma ? Granulomatous Lymphosarcoma	Cushing's syndrome Hodgkin's disease Acute leukaemia	

- Fichtelius, K. E., Laurell, G., and Philipsson, L. (1961). *Acta path. microbiol. scand.*, **51**, 81.
- Fisher, E. R. (1963). *Proceedings of the Thymus Conference*. In press.
- Funkhouser, J. W. (1961). *New Engl. J. Med.*, **264**, 34.
- Gafni, J., Michaeli, D., and Heller, H. (1960). *Ibid.*, **263**, 536.
- Geller, W. (1953). *J. Lab. clin. Med.*, **42**, 232.
- Giltin, D., Janeway, C. A., Apt, L., and Craig, J. M. (1959). In *Cellular and Humoral Aspects of Hypersensitive States*, p. 375. Hoeber, New York.
- Good, R. A. (1954). *Bull. Univ. Minn. Hosp.*, **26**, 1.
- Rötstein, J., and Mazzitello, W. F. (1957). *J. Lab. clin. Med.*, **49**, 343.
- and Varco, R. L. (1955). *J. Lancet (Minneapolis)*, **75**, 245.
- Gowans, J. L., McGregor, D. D., Cowen, D. M., and Ford, C. E. (1962). *Nature (Lond.)*, **196**, 651.
- Hammar, J. A. (1908). *Arch. mikr. Anat.*, **73**, 1.
- Harris, T. N., Harris, S., and Farber, M. B. (1955). *J. Immunol.*, **75**, 112.
- Havard, C. W. H., and Bodley Scott, R. (1960). *Brit. J. Haematol.*, **6**, 178.
- Hitzig, W. H., Biró, Z., Bosch, H., and Huser, H. J. (1958). *Helv. paediat. Acta*, **13**, 551.
- and Willi, H. (1961). *Schweiz. med. Wschr.*, **91**, 1625.
- Holborow, E. J., Asherson, G. L., Johnson, G. D., Barnes, R. D. S., and Carmichael, D. S. (1963). *Brit. med. J.*, **1**, 656.
- Janković, B. D., Waksman, B. H., and Arnason, B. G. (1962). *J. exp. Med.*, **116**, 159.
- Lambie, A. T., Burrows, B. A., and Sommers, S. C. (1957). *Amer. J. clin. Path.*, **27**, 444.
- Langston, J. D., Wagman, G. F., and Dickenman, R. C. (1959). *Arch. Path.*, **68**, 367.
- McEachern, D., and Parnell, J. L. (1948). *J. clin. Endocr.*, **8**, 842.
- MacLean, L. D., Zak, S. J., Varco, R. L., and Good, R. A. (1957). *Transplant. Bull.*, **4**, 21.
- Makela, O., and Nossal, G. J. V. (1962). *J. Immunol.*, **88**, 613.
- Marshall, A. H. E., and White, R. G. (1961). *Brit. J. exp. Path.*, **42**, 379.
- Martin, C. M., Gordon, R. S., jun., and McCullough, N. B. (1956). *New Engl. J. Med.*, **254**, 449.
- Maurer, F. W. (1886). *Morphol. Jahrb.*, **11**, 129.
- Maximow, A. A. (1909). *Arch. mikr. Anat.*, **74**, 525.
- Mendelow, H., and Jenkins, G. (1954). *J. Mt Sinai Hosp.*, **21**, 218.
- Metcalf, D. (1958). *Ann. N.Y. Acad. Sci.*, **73**, 113.
- (1960). *Brit. J. Haematol.*, **6**, 324.
- and Ishidate, M. (1962). *Austr. J. exp. Biol.*, **40**, 57.
- Miller, D. G., Lizardo, J. G., and Snyderman, R. K. (1961). *J. nat. Cancer Inst.*, **26**, 569.
- Miller, J. F. A. P. (1961). *Lancet*, **2**, 748.
- (1962a). *Proc. roy. Soc. B.*, **156**, 415.
- (1962b). *Ann. N.Y. Acad. Sci.*, **99**, 340.
- (1962c). *Nature (Lond.)*, **195**, 1318.
- (1963a). In *Colloque sur la tolérance acquise et la tolérance naturelle à l'égard de substances antigéniques définies*, pp. 47-73. C.N.R.S., Paris.
- (1963b). *Lancet*, **1**, 43.
- (1963c). *Proceedings of the Thymus Conference*. In press.
- (1963d). *Brit. med. Bull.*, **19**. In press.
- (1963e). Unpublished observations.
- Doak, S. M. A., and Cross, A. M. (1963). *Proc. Soc. exp. Biol. (N.Y.)*, **112**, 785.
- Osoba, D., and Miller, J. F. A. P. (1963). *Nature (Lond.)*. In press.
- Ramos, A. J. (1956). *J. Amer. med. Ass.*, **160**, 1317.
- Ringertz, N. (1951). *Acta path. microbiol. scand.*, **29**, 9.
- Sainte-Marie, G., and Leblond, C. P. (1958). *Proc. Soc. exp. Biol. (N.Y.)*, **98**, 909.
- Sherman, J. D., and Dameshek, W. (1963). *Nature (Lond.)*, **197**, 469.
- Silvers, W. K. (1962). *J. cell. comp. Physiol.*, **60**, Suppl. 1, 164.
- Soutter, L., Sommers, S., Relman, A. S., and Emerson, C. P. (1957). *Ann. Surg.*, **146**, 424.
- Stöhr, P. (1906). *Anat. Hefte (Wiesb.)*, **31**, 409.
- Stoner, R. D., and Hale, W. M. (1955). *J. Immunol.*, **75**, 203.
- Strauss, A. J. L. (1963). *Brit. med. J.*, **1**, 56.
- Seegal, B. C., Hsu, K. C., Burkholder, P. M., Nastuk, W. L., and Osserman, K. E. (1960). *Proc. Soc. exp. Biol. (N.Y.)*, **105**, 184.
- Thomson, A. D. (1955). *Brit. J. Cancer*, **9**, 37.
- Waksman, B. H., Arnason, B. G., and Janković, B. D. (1962). *J. exp. Med.*, **116**, 187.
- Waller, J. V., Shapiro, M., and Paltauf, R. (1957). *Amer. Heart J.*, **53**, 479.
- Webster, R. (1961). *Med. J. Aust.*, **48**, 582.
- White, R. G., and Marshall, A. H. E. (1962). *Lancet*, **2**, 120.
- Wintrobe, M. M. (1951). *Clinical Haematology*, 3rd ed., p. 533. Kimpton, London.

QUATERNARY NITROGEN BASES IN HUMAN THYMUS GLANDS

BY

T. N. CALVEY, M.B., Ch.B., B.Sc., Ph.D.

A. WILSON, M.D., Ph.D., F.R.C.P.Glasg.

Department of Pharmacology and General Therapeutics, University of Liverpool

Although it is widely recognized that removal of the thymus gland is often of considerable benefit to patients with myasthenia gravis, the manner by which this effect is produced is a matter of conjecture. The problem has been explored in several ways, and these have been reviewed by Wilson and Wilson (1955). Extracts of thymus glands removed from patients with myasthenia gravis were reported to depress the response of the isolated rat diaphragm to stimulation of the phrenic nerve (Wilson, Obrist, and Wilson, 1953), and subsequently Nowell, Wilson, and Wilson (1959) showed that similar effects were produced on the cat tibialis/sciatic-nerve preparation by extracts of foetal whale thymus glands. These effects were not produced by similar extracts prepared from human lymph nodes and human voluntary muscle. The results of further work on the purification of extracts of foetal whale thymus glands by treatment with ion-exchange resin and paper chromatography were reported by Nowell and Wilson (1961), who identified in the extracts two quaternary nitrogen compounds with characteristic R_f values. The presence of two similar compounds was also detected in extracts of thymus glands removed from two patients with myasthenia gravis. These observations prompted a further investigation of the quaternary nitrogen bases in human thymus glands.

Methods

Collection of Thymus Glands

Thymus glands from patients with myasthenia gravis were obtained immediately after thymectomy. Adventitious

tissue was removed, and each gland was usually weighed before and after the excision of a specimen for histological examination. Each gland was placed in an airtight jar containing acetone A.R. and stored in the refrigerator. One thymus gland was also obtained from a patient with myasthenia gravis at post-mortem examination and stored in a similar way.

Normal thymus glands were obtained from neonatal infants and adults at necropsy and similarly stored in acetone. The term "normal" is not easy to define in this context, for many factors may influence the size and the histological structure of the thymus gland. For the present purposes it was thought desirable to exclude thymus glands from patients who had been given corticosteroids or A.C.T.H., or who had died from infectious or cachectic diseases. In one case a thymus gland was obtained from a female patient during operation for coarctation of the aorta.

Method of Extraction

Each thymus gland was extracted with acetone by the method previously described by Nowell *et al.* (1959). The extracts were further processed by passage through a Seitz centrifugal filter with an ultrafiltration membrane and centrifuged at 1,500 r.p.m. for 15 minutes. The filtrate was then passed through a 5.4 by 0.5 cm. "amberlite I.R.C. 50" (100-200 mesh) ion-exchange column and buffered to pH 6.86 with mixed phosphate buffer. After washing with distilled water the column was eluted with 0.2 M

J. F. A. P. MILLER: ROLE OF THE THYMUS IN IMMUNITY

FIG. 1.—Thymus medulla: 26-weeks-old human foetus. Note Hassall's corpuscles: groups of hypertrophic reticular cells or flattened cells concentrically arranged around a central hyaline or necrotic core. ($\times 395$.)

FIG. 2.—Lymph node of 2-months-old sham-thymectomized (control) mouse. Note close packing of small lymphocytes. ($\times 76$.)

FIG. 3.—Lymph node of 6-weeks-old mouse thymectomized at birth. Note poor cellularity and conspicuous deficiency of small lymphocytes. ($\times 70$.)

FIG. 4.—High-power view of cells in lymph node of 6-weeks-old mouse thymectomized at birth. Note predominance of reticulum cells and several mitotic cells. ($\times 362$.)

FIG. 5.—Spleen of 2-months-old mouse sham-thymectomized at birth. Note follicular structure and germinal centres. ($\times 76$.)

FIG. 6.—Spleen of 2-months-old mouse thymectomized at birth. Note almost complete absence of follicular structure. ($\times 76$.)