While admitting the importance of the curative side of the work, one wonders if we are perhaps neglecting the preventive side, which might yield more eventual success. This would necessitate more work in regard to carriers. Hsia, Paine, and Driscoll¹ showed how heterozygotes of phenylketonuric patients could be detected by their low tolerance for phenylalanine, etc. Perhaps a well-worth-while line of "treatment" on the preventive side would be to have a register of all phenylketonuric families and to practise some scientific marriage guidance in order to prevent, if possible, the matings of heterozygotes. I have been involved, successfully, in a little of this by "enlightened" families. All general practitioners might have knowledge of the phenylketonuric families in their practice and might recognize their responsibilities to them in this respect.

With regard to the genetics of the condition, Penrose² postulated that the condition was due to the action of a single autosomal recessive gene. One wonders what caused the gene to become defective so as to upset the enzyme which it controlled and so bring about the metabolic disturbance which resulted in the grave condition of phenylketonuria. I have noticed that in 50% of the families (albeit only eight families are involved) of my 10 phenylketonuric patients gross alcoholism appears in the parents or grandparents. Considering that alcohol is a liver poison and that the enzymatic action in the metabolism of phenylalanine takes place in the liver, one wonders what is the connexion between alcoholism and phenylketonuria, more especially since one considers gross alcoholism to be a form of metabolic disease rather than a psychiatric illness.—I am, etc.,

J. H. THOMPSON.

Northgate and District Hospital,
Morpeth, Northumberland.

REFERENCES

- ¹ Hsia, D. Y.-Y., Paine, R. S., and Driscoll, K. W., *J. ment. Defic. Res.*, 1957, 1, 53.
- ² Penrose, L. S., *Lancet*, 1935, 2, 192.

Framycetin and the Staphylococcus

SIR,—It has been shown¹ that the topical application of framycetin results in a significant reduction of the number of carriers of *Staphylococcus aureus* in a defined test population. Porter *et al.*² in a recent study of a hospital population have presented data which they have interpreted as showing framycetin to be of dubious value in the prophylaxis and control of nasal carriage of *Staph. aureus* strains. Some obvious criticisms of the conclusions of the latter group of workers have already been made (Rubbo *et al.*, July 27, p. 252). Our findings in Sheffield, based on investigations carried out on the nursing staff, have shown that framycetin is indeed of value. An outline of the studies which have been made, and upon which this opinion is based, is given below.

1. Short-term experiment

140 members of the nursing staff of the Children's Hospital, Sheffield,

were allocated at random to one of two groups. The carriage rate of *Staph. aureus* was determined by daily vestibular swabbing on each of five consecutive days. Swabs were cultured in nutrient broth for 24 hours. Broth cultures were subcultured on blood agar, and *Staph. aureus* colonies identified by their characteristic pigmentation. Identity was confirmed by the coagulase test. All isolates were phage typed. Staphylococcal carriers were defined as persons from whom the same phage type of *Staph. aureus* was isolated on at least two of the five days. The carriage rate in the 2 groups was 44% (20/45) and 33% (32/95) respectively. All 45 members of the first group were given nebulizers of framycetin, and instructed to use these once daily before coming on duty. The second group was not treated. The number of carriers in the two groups was determined after one week's treatment. The carriage rate was 9% (4/45) in the test group and 36% (34/95) in the control group.

2. Long-term experiment

Using the same protocol the efficacy of topical framycetin in the long-term control of nasal carriage of *Staph. aureus* was investigated. Of 200 members of the medical, nursing, and domestic staffs volunteering to participate, 98 submitted the requisite number of swabs over a 12-week treatment period, and these only have been analysed. The initial carriage rate prior to treatment was 33%. After one week's treatment this fell to 8%. Over the following six weeks the carriage rate rose to 17%. It remained more or less constant at this level during the remainder of the 12-week trial. Only one strain of *Staph. aureus* was found to be resistant to framycetin.

Analysis of the results in terms of occupation showed that the majority of the nursing and medical staff remained free of *Staph. aureus* during the trial period. The bulk of the 17% remaining persistently positive was made up of members of the domestic staff.

To confirm the impression that the success or failure of topical framycetin in the control of nasal carriage of *Staph. aureus* was largely a measure of the enthusiasm and discipline of the users, the following small-scale investigation was carried out. An intensive study was made of four chronic nasal carriers of *Staph. aureus*. Swabs were taken five times weekly for four weeks from the anterior nares, axillae, and perineum. All showed constant carriage in the anterior nares. Three yielded frequent isolations from the perineum and one frequent isolations from the axillae.

The anterior nares were treated by using the framycetin nebulizer once daily and the perineum and axillae by the application of framycetin cream (1.5%) (again once daily). Daily swabbing for four weeks failed to yield a growth of *Staph. aureus* after treatment had been initiated for two days. After four weeks treatment was discontinued. The previously colonized body sites recolonized in 1–2

weeks. In three cases recolonization was with the same phage type as had been isolated prior to treatment. The anterior nares remained free of *Staph. aureus* for 6–10 weeks after treatment had been discontinued. In all four subjects recolonization was with the same phage type as that isolated from the previously colonized body site. In the light of this one must seriously reconsider the statement of Porter *et al.* that the re-establishment of carriage in the anterior nares is due to persistence of small numbers of organisms in this site. One must also conclude that the value of topical framycetin, like that of any other therapeutic or prophylactic measure, is only as great as the enthusiasm and care with which it is used.—I am, etc.,

WILLIAM I. H. SHEDDEN.

Department of Bacteriology,
The University, Sheffield.

C. W. POTTER.

A. M. BRUCE.

Children's Hospital,
Sheffield.

REFERENCES

- ¹ Stratford, B., Rubbo, S. D., Christie, R., and Dixon, S., *Lancet*, 1960, 2, 1225.
- ² Porter, I. A., McNeill, I. F., Miller, I. T., and Green, C. A., *Brit. med. J.*, 1963, 1, 1515.

Future of General Practice

SIR,—From letters and articles in both the medical and the lay press it can be gathered that the medical profession is thoroughly demoralized, resulting in internal and external friction. Therefore, Sir, I would be grateful to air a few of my scattered thoughts in your columns, wondering how many of your members share my anxieties.

Salary is the most popular topic to-day. Though financially insecure I should be as glad of a lowering of expenses as of an increase in salary. I have no desire for my income to be made public, but if this be essential then I prefer that the figure be not distorted. I deplore the petty bickering over pay differentials between consultants and practitioners. It's about time each praised the other as they so often do on domiciliary visits.

Indeed, teamwork is essential—not only between specialist and general practitioner, but with the Ministry, social workers, the churches, local government, and teachers. The meetings and discussions held at frequent intervals between the doctors, clergy, and social workers in this area have been of great help to the participants. The combination of all our knowledge and resources has saved us duplicating work and made second opinions easily available on many problems, not necessarily medical. We realize and appreciate each other's functions and points of view, and are learning to work in harmony.

Of further causes of internal friction in the profession, a major one will be human experimentation. The art of medicine nowadays is the control of science, and so research must be a slave rather than a master. Many of my colleagues now find it necessary to interfere in the often over-enthusiastic treat-

ment of malignancy. In my view patients benefit more from a shorter period of relative comfort than prolonged misery through drastic treatment, even though the latter may lead more rapidly to eventual cures for other people.

Recently I witnessed on television a rather sordid dispute between medical people on human experimentation in hospital. I doubt that this will help when I have to persuade a nervous patient that hospital admission is essential. If asked by a patient, "Will they experiment on me?" what do I answer? So much of medical practice is experimental that we must be trusted not to carry this too far. We cannot believe in God and act like one at the same time.

Statistics, too, seem to be in the ascendancy, mainly, I assume, because of the convenience of interpreting them in accordance with the trend of the times. For instance, article upon article exhorts me to encourage smokers to desist, but are there any statistics to show what will happen when the heavy smokers, who are orally dependent, abstain? Will their demise be any further off, suffering from obesity and cardiac debility secondary to increased intake of confectionery? What percentage will degenerate under the influence of alcohol or drugs, and how many innocents will they slaughter on the roads? Indeed, will their increased sexual prowess be a problem of import? These are only a few of many questions that might be considered. Personally, I shall know some of the answers soon myself, having just given up smoking, mainly for financial reasons.

Finally, stress. The percentage of patients who require a doctor's help through being unable to stand the strain of modern living increases yearly. Many of the patients are of sufficient calibre to give one the impression that they have been subjected to more stress than the majority of persons could adequately have coped with, and do not come under the heading of inadequate personalities. Thus there is a great need for research by medical and other authorities into stress and the avoidance of excessive strain. This would indeed remove so many problems from, and increase greatly the standard of, medicine in general practice.—I am, etc.,

GRAHAM KNIGHT-WEBB.

Minchinhampton,
Gloucestershire.

Pulmonary Oedema

SIR,—The recent articles concerned with pulmonary oedema (July 13, pp. 80 and 83) give a further stimulus to those of us interested in the complications of intracranial and spinal injuries. The review by Dr. P. Richards (p. 83) is very valuable, but a study of pulmonary oedema of neurogenic origin is probably more fruitful if the factors of increased intracranial pressure and operative trauma are excluded. Blood products in the cerebral ventricles are a potent source of pulmonary oedema.¹

The vast amount of experimental physiology on the central autonomic system points strongly to the presence of numerous receptor centres, perhaps of almost pin-point size, and located chiefly in the hypothalamus. While it is agreed that conclusions drawn from animal experiments must be guarded in their application to the human, nevertheless such experiments provide a basis for ideas which can be applied to the study of clinical states. (With due respect to Dr. Richards, it is still controversial that bilateral division of the vagus of itself can produce pulmonary oedema except in the guinea-pig.)

Macroscopic study is not able to detect trauma to which nerve cells and their neurones may be subjected. It seems beyond doubt that gross damage to nerve cells and fibres may even be missed by the microscope.²⁻⁴ Moreover, a surprising amount of recovery may follow traumatic lesions in the central nervous system. Therefore it can be inferred that physiological lesions occur which are reversible.

There is abundant evidence in favour of the general concept of central receptor centres concerned with autonomic function. It is essential to appreciate that perverted activity of the autonomic system may result from damage to the afferent and efferent tracts associated with these centres, as well as trauma to the cells of the centres themselves. Hence it is unlikely that a central solitary and anatomically precise location will be found to explain every particular complication occurring in the periphery.

From a study of head and high spinal injuries over many years it was concluded⁵ that so-called hypostatic pneumonia was in essence a pulmonary oedema of neurogenic origin. It was further concluded that the damage to the central areas of autonomic function resulted in a perverted release of adrenergic activity in the periphery. Destruction of the peripheral sympathetic system does not precipitate pulmonary oedema.

In gross damage to the brain stem there is frequently observed a triad of complications consisting of pulmonary oedema, acute hypertension, and hyperthermia. Arnott⁶ suggested that the pulmonary oedema of head injuries was due to the acute rise in systemic vasoconstriction, and indeed the so-called adrenaline oedema may add support to that view. Nevertheless, as pointed out in a letter to the *B.M.J.*,⁷ acute hypertension is not invariable in pulmonary oedema, an observation confirmed by Dr. Richards. Certainly it is absent in the pulmonary oedema associated with high spinal injuries. There seems little doubt that neurogenic pulmonary oedema results from some peculiar disturbance of haemodynamics and hence the apt term, neurohaemodynamic, coined by Sarnoff.⁸ If this disturbance of haemodynamics is determined by some localized release of adrenergic function in the periphery, then inhibition of such adrenergic function should cure neurogenic pulmonary

oedema. Success has been claimed by Fontaine and Courtine⁹ by the injection of local anaesthetic into the upper dorsal sympathetic ganglia. An injection of ergotamine tartrate produced a dramatic effect in a severe pulmonary oedema associated with a high spinal transection.¹⁰

Experimentally the development of pulmonary oedema has been judged on the alteration of the heart-lung weight ratio. It is desirable to distinguish oedema of the lung tissues from the condition of oedema fluid in the air passages. In some work, unpublished as yet, I have had the temerity to coin the word pneumoedema for this latter state for the purpose of clarity in argument. There seems little doubt that these oedematous states can readily predispose to pneumonic changes. Such terminal pneumonia, described under several terms by pathologists, has no doubt tended to mask the true pathological changes. Moon¹¹ has reviewed these terminal states. Dr. Richards comments on the variable site of these changes, though they are most commonly basal. Certainly posture plays a leading part. The sheep on its back will quickly succumb with acute pulmonary oedema, a fact well known to country folk. The bucolic wisdom of going immediately to the sheep's assistance is well founded. Even so the distribution of blood flow and ventilation in the lungs is a study in itself, and the recent edition of the *British Medical Bulletin* on "Respiratory Physiology" is of great value to those interested in the problems of pulmonary oedema.—I am, etc.,

Wembley Hospital, J. S. CALLCUTT.
Wembley, Middlesex.

REFERENCES

- 1 Noordenbos, W., *Folia psychiat. neerl.*, 1954, 57th year No. 4/5, 405.
- 2 Greenfield, J. G., *Proc. roy. Soc. Med.*, 1938, 32, 43.
- 3 Lhermitte, J., *Folia psychiat. neerl.*, 1950 (Brouwer Mem. Vol.).
- 4 Strich Sabina, J., *Lancet*, 1961, 2, 443.
- 5 Callcutt, J. S., *Med. Wld (Lond.)*, 1943, 58, 279.
- 6 Arnott, W. M., *Brit. med. J.*, 1955, 2, 342.
- 7 Callcutt, J. S., *ibid.*, 1955, 2, 735.
- 8 Sarnoff, S. J., *J. Physiol.*, 1952, 117, 38P.
- 9 Fontaine, R., and Courtine, G., *Presse méd.*, 1940, 48, 711.
- 10 Callcutt, J. S., *Med. Wld (Lond.)*, 1945, 62, 467.
- 11 Moon, V. H., *Arch. Path.*, 1938, 26, 132.

Discharge from Mental Hospitals

SIR,—On four occasions so far this year I have been called by the police to attend patients who have recently been in a mental hospital. Three of these patients, a man and two women, were picked up wandering about in a confused state and required immediate readmission to a mental hospital. They had previously been in hospitals outside the Oxford area.

The fourth patient, an elderly man, was found dead on the floor of his squalid dwelling. He had been dead for about two months. Necropsy did not reveal the cause of death. He had been discharged from a mental hospital a few days before last being seen alive.