IAN DONALD: USE OF ULTRASONICS IN ABDOMINAL SWELLINGS

W. F. HARPER AND J. MacVICAR: HYDATIDIFORM
MOLE DIAGNOSED BY SONAR

LEGENDS: IAN DONALD

(In the transverse sections, R. and L., the view is that of the gynaecologist—that is, from below upwards—so that right in the picture is left on the patient, as in an x-ray. CR=cranial. CA=caudal.)

FIG. 9.—Hepatomegaly associated with drug-induced jaundice. (Longitudinal section.) (a) Liver. (b) Bowel echoes. (c) Mark at umbilicus.

FIG. 10.—Liver distorted and enlarged with metastases. (Longitudinal section.) (a) Liver.

FIG. 11A.—Enlarged spleen due to myelofibrosis, 1960.

FIG. 11B.—Same case in 1963, showing increase in splenomegaly. (Both in transverse section.) (a) Spleen.

FIG. 12.—Fibromyoma and pregnancy. Tumour is to left of picture and foetal echoes to right. (Longitudinal section.) (a) Fibromyoma. (b) Foetal echoes. (c) Bowel echoes.

FIG. 13.—Hydatidiform mole distending uterus. Longitudinal section. (a) Mole. (b) Blood clot.

FIG. 14.—Polycystic kidney. (Transverse section.) (a) Polycystic kidney. (b) Bowel echoes.

FIG. 15.—Large fairly homogeneous renal tumour in transverse section. (Spindle-cell sarcoma.) (a) Tumour mass.

FIG. 16.—Goitre straddling trachea. (Transverse section.) (a) Trachea. (b) Goitre.

FIG. 17.—Normal anteverted uterus demonstrated behind a full bladder. (Longitudinal section.) (a) Full bladder. (b) Anteverted uterus. (c) Endometrial cavity. (d) Bowel echoes.

LEGENDS: HARPER AND MacVICAR

FIG. 1.—Transverse section taken 6 cm. below umbilicus. Note typical speckled appearance of the echoes from the vesicles of hydatidiform mole.

FIG. 2.—Transverse section taken 12 cm. below umbilicus. There are definite echoes in a cystic cavity indicative of foetal parts.

FIG. 3.—Longitudinal scan from umbilicus, right, to symphysis pubis, left. The uterus is outlined and the mole can be seen at the fundus with foetal echoes in the lower pole.

plained of buzzing in the ears, and pins-and-needles and weakness in the legs. He was dysarthric and ataxic, with considerable loss of power in the legs, absent tendon reflexes, and impaired superficial sensation.

Investigations.—Blood count, blood urea, electrolytes, liver function tests, and chest x-ray picture: normal. W.R. negative. C.S.F. protein 80 mg./100 ml. Urinary 17-ketosteroids less than 2 mg./24 hours. Pyruvate tolerance and liver biopsy normal.

Progress.—After each admission he quickly responded to intravenous parentrovite, modified insulin, and chlorpromazine. His neurological symptoms and signs cleared after two weeks. After discharge or when on leave from the hospital he soon reverted to his old habits.

Discussion

In the patients described the average daily consumption of chlorodyne was at least 6 oz. (170 ml.). They were therefore taking 6 gr. (0.4 g.) of morphine and 360 min. (21 ml.) of chloroform daily. There existed an overpowering desire to continue taking the preparation, increasing the dosage from day to day. In the end they were totally dependent on it. Their progressive social and physical deterioration was no deterrent. All had been in an unconscious state several times, and one had been admitted to a general hospital in coma on no fewer than eight occasions. All were psychotic at one time or other, necessitating admission to a mental hospital. Visual hallucinations were present in two.

The similarity of presentation with severe physical debility, marked weight loss, sallow grey complexion, tremor, and dysarthria was striking. All had symptoms and signs of peripheral neuropathy, one of which, so far, seems irreversible.

Two patients had a low blood-pressure. This is a well-known effect of morphine and chloroform, but, together with the curious pigmentation of the skin and the low urinary 17-ketosteroids, suggested the presence of adrenal insufficiency. Two patients consented to liver biopsy. The extent of liver damage on biopsy in the first was remarkable. The hepatotoxic effects of chloroform and alcohol, however, are well known, and it is arguable that prolonged malnutrition may also have contributed. Unfortunately it is not possible to say with certainty whether the deleterious effects on the liver were due to morphine, alcohol, or chloroform.

All three patients showed behaviour disturbance. The degree of personality maladjustment prior to addiction was

difficult to determine, but they were emotionally immature, shy, sensitive people with a very low stress tolerance. Two were subject to marked mood changes from an early age, and the second patient was treated for depression by her family doctor at the age of 32. They were all unmarried and complained bitterly of the loneliness of their lives.

Over the past two years three other patients who were taking excessive amounts of chlorodyne have been admitted to this hospital, but the picture was complicated by addiction to barbiturates, amphetamines, or alcohol in addition.

A number of references have been made from time to time in the *Pharmaceutical Journal* to coroners' inquests on deaths due to chlorodyne-poisoning. In all cases there was a history of chronic addiction for many years. Coroners have commented on the ready availability of chlorodyne and the lack of legislation regarding its sale.

Seager and Foster (1958) described the cases of two patients who were addicted to the preparation, one of whom was acutely psychotic with visual and auditory hallucinations.

Personal inquiries from chemists in the area have established that there is no difficulty in obtaining at reasonable cost large quantities of this preparation, for which there is considerable demand. Its ready availability is to be deplored, and there is a strong case for amending existing legislation, so that chlorodyne becomes available on prescription only by inclusion in Schedule 4 of the Poisons Act.

Summary

The cases of three patients who were addicted to chlorodyne are described. There was evidence of serious mental and physical deterioration, including peripheral neuropathy.

The ready availability and the large quantities of morphine and chloroform consumed make chlorodyne a potentially dangerous drug. It is suggested that it would be better scheduled as a poison available only on prescription.

My thanks are due to Dr. J. J. O'Reilly, medical superintendent, for permission to publish, and to Dr. A. Paton, consultant physician, Dudley Road Hospital, for helpful advice and criticism throughout.

REFERENCES

- Seager, C. P., and Foster, A. R. (1958). *Brit. med. J.*, **2**, 950.
World Health Organization (1950). *Techn. Rep. Ser.* No. 21, p. 6.

Medical Memoranda

Hydatidiform Mole and Pregnancy Diagnosed By Sonar

[WITH SPECIAL PLATE]

The onset of pre-eclamptic toxæmia prior to the twentieth week of pregnancy is very suggestive of the presence of a hydatidiform mole. The definite diagnosis of hydatidiform mole can be very difficult, since biological tests for chorionadotrophin are often inaccurate; but recently the use of sonar has enabled the diagnosis to be made with certainty and immediately on examination (Donald and Brown, 1961; MacVicar, 1962; MacVicar and Donald, 1963).

CASE REPORT

A primigravida aged 26 who had had amenorrhoea for 16 weeks was admitted to hospital because of recent swelling of the ankles, feet, and hands. On admission she was found to have generalized oedema. Her blood-pressure was 190/100 mm. Hg, and the urine contained albumin (2.5 g./litre).

On examination the uterus, which felt cystic, appeared to be only slightly larger than the dates would suggest. On vaginal examination the cervix was long and not "taken up." No internal ballotement of foetal parts could be elicited. There was neither vaginal bleeding nor discharge. A "prepuerin" (Burroughs Wellcome Ltd.) pregnancy test was weakly positive only and a straight radiograph of the abdomen showed no foetal parts.

During the ensuing week the patient was treated with sedation and complete bed rest, but her blood-pressure remained raised, the highest reading being 220/125 mm. Hg. The urine, though microscopically clear, constantly contained albumin, the maximum amount being 4 g./litre. The oedema persisted. She

had no headache or visual upset. Seven days after admission the Aschheim-Zondek test was reported as positive in a dilution of 1 in 100, and it was decided to examine the patient by sonar for the possible presence of hydatidiform mole.

FIG. A.—Uterine contents showing the dead foetus still in its sac and the hydatidiform change involving the placenta.

Sonar Examination.—Several scans of the patient's abdomen were taken both in longitudinal and in transverse axes. The upper part of the uterus, which was clearly visualized by echosounding, seemed full of the typical speckled appearance characteristic of hydatidiform mole (Special Plate, Fig. 1), but in the lower portion of the uterus there was a well-marked cystic space, containing definite echoes (Special Plate, Fig. 2), which we recognize as being consistent with foetal parts (Special Plate, Fig. 3). The diagnosis was therefore made of the possibility of a twin pregnancy with hydatidiform mole having replaced one foetus, while the other was normal.

As the patient's condition did not improve, the decision to terminate the pregnancy was further reinforced by the sonar confirmation of the existence of hydatidiform mole.

Operation.—Under anaesthesia, pelvic examination revealed ballotable foetal parts, and it was decided to terminate by abdominal hysterotomy. The uterine contents were removed complete and intact. The specimen consisted of a missed abortion in a sac containing a dead foetus below, while the placenta was completely replaced by hydatidiform degeneration above (Fig. A). The ovaries were normal.

The patient made an uneventful recovery and was allowed home 14 days after operation. Her blood-pressure on dismissal was 140/90 mm. Hg, the urine contained only a trace of albumin, and the oedema had disappeared. An Aschheim-Zondek test one week after operation was negative in all dilutions.

Six weeks after operation the patient was well and had no complaints. Her blood-pressure was 120/80 mm. Hg; there was no oedema. The urine was normal. Pelvic examination revealed no abnormality and the Aschheim-Zondek test was again negative in all dilutions.

Histological Report.—Sections of the placenta showed moderate oedema and stromal degeneration of the villi, many of which were still normal, so that it was presumably still in

the early stages of hydatidiform degeneration. There was no evidence of syncytial proliferation or malignancy.

DISCUSSION

This patient developed the clinical picture of fulminating pre-eclamptic toxæmia in the second trimester of pregnancy, and it was known that her blood-pressure had been normal previously and that she had no history of renal disease. Clinical examination of the abdomen suggested a normal pregnancy rather than hydatidiform mole, since the uterus was cystic and not significantly larger than suggested by the dates. The pregnancy tests were contradictory, the prepuerin test being a weak positive and the quantitative Aschheim-Zondek test being positive at dilution of 1 in 100. Radiography showed no evidence of foetal parts, probably because the foetus had barely reached 16 weeks' gestation. We have found that sonar can detect foetal parts much earlier than this—in fact, between eight and nine weeks (MacVicar and Donald, 1963).

It is interesting that when this case was examined by sonar, not only were foetal echoes seen but also the typical picture of hydatidiform mole.

We should like to thank Dr. D. McKay Hart for his advice and help, and Professor Ian Donald for his confident diagnosis of an unusual sonar picture.

WILLIAM F. HARPER, M.B., Ch.B.,
B.Sc., M.R.C.O.G.,
Consultant Obstetrician and Gynaecologist.

Stobhill Hospital,
Glasgow.

JOHN MACVICAR, M.D., M.R.C.O.G.,
Lecturer in Midwifery.

University of Glasgow.

REFERENCES

- Donald, I., and Brown, T. G. (1961). *Brit. J. Radiol.*, **34**, 539.
MacVicar, J. (1962). *Proc. roy. Soc. Med.*, **55**, 638.
— and Donald, I. (1963). *J. Obstet. Gynaec. Brit. Cwlth*, **70**, 387.

Spontaneous Uterine Rupture in Association with Abortion

It is the purpose of this report to draw attention to spontaneous rupture of the uterus as a serious complication of abortion. Rupture of the uterus in late pregnancy or labour is well recognized, but its earlier occurrence is seldom reported.

CASE 1

A Cypriot woman aged 23 was admitted to another hospital on account of vaginal bleeding at about the twentieth week in her second pregnancy. The date of her last period was unknown. She had a normal confinement four years previously but no miscarriage or serious illness. The results of the usual antenatal investigations were normal.

No bleeding occurred after admission, but four days later her mental state necessitated moving her to a mental hospital.

After 10 days she complained of backache, followed by vaginal bleeding, which continued for eight days. Placenta praevia was suspected. Uterine contractions then started and she was transferred to the gynaecological unit at the Whittington Hospital. Her general condition was good, the blood-pressure being 130/70 mm. Hg. On pelvic examination some blood was found in the vagina, the cervix admitted one finger, and the head was presenting; no placenta was felt.

Four hours later a dead foetus, followed by its placenta, was passed. Bleeding amounting to about 500 ml. due to