origin and symptomless unless they become accidentally infected or very large. Symptoms arising in a previously quiescent diverticulum as a result of infection due to trauma, either from catheterization or the passage of the foetal head, would explain the high incidence of parous women in the series so far recorded.

Urethral diverticula, while apparently not so common in Great Britain as in some parts of the United States of America and the West Indies, are by no means very rare. I have personally encountered two in London in the past four years, and Mr. Geoffrey Dixon has recently drawn my attention to a third example which, like Dr. Phillips's case, contained calculi.

It is likely that if looked for carefully by British gynaecologists and urologists urethral diverticula would be found more often. This has certainly been the case elsewhere.—I am, etc.,

J. H. M. PINKERTON.

Department of Midwifery and
Gynaecology,
The Queen's University of Belfast.

REFERENCES

- ¹ Downer, I. G., and Virgilio, F. D., *J. Urol.*, 1945, **54**, 53.
- ² Pinkerton, J. H. M., *J. Obstet. Gynaec. Brit. Emp.*, 1956, **63**, 76.
- ³ McMahon, S., *J. Urol.*, 1946, **55**, 69.

Antibiotic Cover for Dental Extractions

SIR,—In the clinicopathological conference on valvular disease of the heart (August 31, p. 545) Dr. J. F. Goodwin stated that he had not seen subacute bacterial endocarditis from *Streptococcus viridans* develop after dental extractions in which antibiotic cover had been used. For this reason we should like to record the following case.

A primigravida aged 27 was seen in the second trimester of pregnancy and a diagnosis of severe rheumatic mitral stenosis and incompetence and severe aortic incompetence with cardiomegaly was made. She gave a past history of probable subacute bacterial endocarditis in other hospitals on three occasions. She had a non-toxic goitre and dental sepsis, and in view of the latter was referred for dental surgery. Extraction was advised and was carried out under a penicillin umbrella. The patient received 6,000,000 units of crystalline penicillin; the first dose of 1,000,000 units was given approximately one hour before the extraction, two further doses of 1,000,000 units were given on the day of extraction, two on the day following, and a final 1,000,000 units on the morning of the second post-operative day. In addition she was given penicillin V, 250 mg. three times a day for four days, starting on the day of extraction. Two weeks later she was admitted to hospital complaining of pain in the loin and the abdomen, and frequency. Her temperature was 97° F. (36.1° C.) and there were no stigmata of subacute bacterial endocarditis. For some days previously she had had a swinging fever up to 102° F. (38.9° C.) and had been treated by her practitioner with "terramycin" (oxytetracycline) and "furadantin" (nitrofurantoin) for four days for presumed pyelitis of pregnancy. Furadantin 100 mg. four times a day was given for a

further three days. Five days after all antibiotic treatment was discontinued her temperature rose to 102° F. (38.9° C.). Blood culture showed *Streptococcus viridans* sensitive to penicillin, "achromycin" (tetracycline), "chloromycetin" (chloramphenicol), terramycin, and streptomycin. She was treated with crystalline penicillin (2,000,000 units eight-hourly) for six weeks. On cessation of treatment she remained afebrile and blood culture was sterile. Spontaneous vaginal delivery of a living male child was made under penicillin cover at term. The puerperium was uncomplicated.

It seems likely that the streptococcal septicaemia in this case was related to the dental extraction, and at the time of the positive blood culture subacute bacterial endocarditis was actual or imminent. The patient had appeared ill within a few days of discontinuation of antibiotic prophylaxis and had been treated with terramycin for presumed pyelitis of pregnancy. Examination of the urine subsequent to her admission to hospital, though repeated, showed no evidence of urinary tract sepsis, although red blood cells were seen on one occasion. It seems therefore that streptococcal septicaemia persisted in spite of attempted prophylaxis with penicillin. The organism recovered from the blood-stream was sensitive to penicillin, and the patient recovered after prolonged treatment with penicillin. We are at a loss to explain the apparent failure of prophylaxis in this case.—We are, etc.,

C. G. BARNES.

ROSALINDE HURLEY.

Queen Charlotte's Maternity Hospital,
London W.6.

Nicotine and Colonic Muscle

SIR,—In the interesting preliminary communication published in your journal on the above-mentioned subject (September 14, p. 666) it was observed by Dr. D. J. Fishlock and Mr. A. G. Parks that nicotine did not cause stimulation of the human colonic muscle, *in vitro*, in any dose range. In their experiments the studies were made on the strips of colon obtained from patients under anaesthesia, and it is likely that these subjects were atropinized. On rabbit colon nicotine has been shown to cause stimulation which is abolished by atropine.¹

It is probable that on human colon also the initial pre-anaesthetic atropine may have been given which would block the muscarinic effect of nicotine. It would be interesting to know the type of pre-medication given to these patients. If atropine was in fact given, were the reported responses to acetylcholine quantitatively in a correct range, or were these due to excess acetylcholine overcoming the block?—We are, etc.,

K. C. DAVE.

K. S. SACHDEV.

Department of Pharmacology,
M. P. Shah Medical College,
Jamnagar, India.

REFERENCE

- ¹ Gillespie, J. S., and Mackenna, B. R., *J. Physiol. (Lond.)*, 1960, **152**, 191.

Pigmentation and Vitamin-B₁₂ Deficiency

SIR.—Our attention has just been drawn to an excellent article published in 1944 by Dr. Bramwell Cook entitled "A Vitamin B Deficiency Syndrome Allied to Sprue"¹ in which he gives a beautiful description of the type of pigmentation described in our article of June 29 last (p. 1713). He says, "The skin overlying the knuckles of the fingers looks darker than normal, and his complexion is sallow and muddy. There may be darkness of the skin around the neck and the lips. The patient is conscious and concerned about his blackness. . . ." And again, "The common sites of pigmentation are: on the knuckles, the tongue (black spots may appear on the dorsum of the tongue), around the lips and in the nasolabial folds, on the forehead, under the eyes, on the dorsum of the ankles, back of the wrist, extensor surface of the arms, front of the neck, front of the legs below the knees."

Dr. Cook also observed that this pigmentation was associated with a macrocytic anaemia, and that both it and the anaemia responded to crude liver extract.

As far as we are aware this is the first documented description of this type of pigmentation which we now know to be associated with vitamin-B₁₂ deficiency. —We are, etc.,

S. J. BAKER.

MERCY IGNATIUS.

S. JOHNSON.

S. K. VAISH.

Wellcome Research Unit,
Christian Medical College,
Vellore, South India.

REFERENCE

- ¹ Cook, B. A., *Indian med. Gaz.*, 1944, **79**, 429.

Toxic Reactions to Drugs

SIR.—The Subcommittee on Adverse Reactions of the Committee on Safety of Drugs intends to form a central register of toxic reactions to drugs both old and new and will shortly be issuing forms for the notifications of such reactions. One of the main tasks of the committee will be to assess the incidence of reactions in relation to the frequency of prescription of the drugs concerned. We therefore hope to make the central register as complete and as accurate as possible. We shall be grateful if any groups of workers or societies in the United Kingdom who are already keeping a register of adverse reactions or are contemplating doing so will get in touch with the Subcommittee on Adverse Reactions so that information can be exchanged and duplication avoided. Communications should be addressed to the Medical Secretary, Committee on Safety of Drugs, 1 Alexander Fleming House, Elephant and Castle, London S.E.1.—I am, etc.,

L. J. WITTS,

Chairman,

Subcommittee on Adverse Reactions,

The Radcliffe Infirmary,
Oxford.