USE OF ULTRASONICS IN DIAGNOSIS OF ABDOMINAL SWELLINGS

BY

IAN DONALD, M.B.E., M.D., B.A., F.R.C.S.Glas., F.R.C.O.G.

Regius Professor of Midwifery, University of Glasgow

[WITH SPECIAL PLATE]

Ultrasonic echo-sounding (sonar) has something to offer in the field of diagnosis, particularly in the case of large tumour masses within the abdomen (Donald *et al.*, 1958; Donald and Brown, 1961; MacVicar, 1962).

Our experience, predominantly of course gynaecological and obstetrical, extends over many hundreds of cases, and we are gradually building up a library of characteristic diagnostic appearances. Some of these are presented here.

Fig. A shows the apparatus at present in use at the Western Infirmary in Glasgow, which has been developed and built for us by Smiths Industrial Division Ltd. It will be noted that the patient does not have to be moved from her bed, and the only inconvenience is the application of a film of olive oil to the abdominal surface. With the use of a polaroid camera a picture is obtainable in longitudinal or transverse section within two or three minutes (Fig. 1, Special Plate—normal).

Fig. 2 (Special Plate) shows in diagrammatic form the scanning technique which we have developed in order to

depict the structures deep within the abdomen, detecting the echoes from them and projecting them in the form of light on the face of a cathode-ray tube which is photographed. The system is a multiple-angle sector-scanning technique already well known in radar.

Fig. 3 is an ultrasonogram in transverse section of a large ovarian cyst in which the vertebral column can be seen projecting into the posterior wall.

By contrast a fibroid (Fig. 4), because of its denser structure, is less transparent or, more correctly, less "transonic"; and here the far wall of the fibroid is not depicted at a given frequency of $2\frac{1}{2}$ megacycles, but Fig. 5 shows the same fibroid examined at a coarser frequency of $1\frac{1}{2}$ megacycles in which the posterior wall is now revealed. This difference is pathognomonic of fibroids.

Abdominal swelling due to ascites is shown in transverse section in Fig. 6. The free fluid is visible in each flank, whereas the bowel floats up into the centre, thus reproducing the well-recognized clinical signs on percussion.

This ultrasonogram is of a man suffering from cirrhosis of the liver and without evidence of malignant disease.

Fig. 7 reveals the very different picture of malignant ascites due to ovarian carcinomatosis in a woman of 43. The very bizarre picture seen here we now recognize as characteristic.

Fig. 8A is an ultrasonogram in longitudinal section from a case of cardiac ascites. The patient, a woman of 72, was in right-sided congestive cardiac failure. It will be seen that the large liver (on the right of Fig. 8A) extends almost down to the umbilicus (shown with a vertical mark on the abdominal surface), the bowel echoes all float up towards the centre, and the free fluid wells up out of the pelvis (left of picture).

The same case is shown in Fig. 8B after three and a half weeks' treatment with diuretics and proper care in a medical ward. It will be

FIG. A.—Present apparatus.

LEGENDS FOR SPECIAL PLATE

(In the transverse sections, R. and L., the view is that of the gynaecologist—that is, from below upwards—so that right in the picture is left on the patient, as in an x-ray. CR=cranial. CA=caudal.)

FIG. 1: Normal.—(A) Longitudinal and (B) transverse sections (same case). (a) Probe rocking movement over skin. (b) Abdominal wall. (c) Bowel echoes.

FIG. 2.—Diagram of multiple-angle sector-scanning technique employed.

FIG. 3.—Large ovarian cyst in transverse section. (a) Cyst cavity. (b) Vertebral column.

FIG. 4.—Fibromyoma—not transonic to $2\frac{1}{2}$ megacycles ultrasound. (Longitudinal section.) (a) Fibromyoma. (b) Posterior wall not visible.

FIG. 5.—Same fibromyoma as in Fig. 4. Posterior wall now revealed using frequency of $1\frac{1}{2}$ megacycles. (Longitudinal section.) (a) Fibromyoma. (b) Posterior surface of fibromyoma now visible.

FIG. 6.—Ascites—non-malignant. (Transverse section.) (a) Free fluid in flanks. (b) Bowel echoes floating up into centre of abdomen.

FIG. 7.—Ascites due to ovarian carcinomatosis. (Transverse section.)

FIG. 8A.—Ascites due to congestive cardiac failure with hepatic enlargement (to right of picture). (a) Liver. (b) Bowel echoes. (c) Mark at umbilicus. (d) Free fluid.

FIG. 8B.—Same case after treatment. Ascites now gone (Both cases in longitudinal section.) (a) Liver. (b) Bowel echoes. (c) Mark at umbilicus. No free fluid.

IAN DONALD: USE OF ULTRASONICS IN ABDOMINAL SWELLINGS

FIG. 1

FIG. 8

IAN DONALD: USE OF ULTRASONICS IN ABDOMINAL SWELLINGS

IAN DONALD: USE OF ULTRASONICS IN ABDOMINAL SWELLINGS

W. F. HARPER AND J. MacVICAR: HYDATIDIFORM
MOLE DIAGNOSED BY SONAR

LEGENDS: IAN DONALD

(In the transverse sections, R. and L., the view is that of the gynaecologist—that is, from below upwards—so that right in the picture is left on the patient, as in an x-ray. CR=cranial. CA=caudal.)

FIG. 9.—Hepatomegaly associated with drug-induced jaundice. (Longitudinal section.) (a) Liver. (b) Bowel echoes. (c) Mark at umbilicus.

FIG. 10.—Liver distorted and enlarged with metastases. (Longitudinal section.) (a) Liver.

FIG. 11A.—Enlarged spleen due to myelofibrosis, 1960.

FIG. 11B.—Same case in 1963, showing increase in splenomegaly. (Both in transverse section.) (a) Spleen.

FIG. 12.—Fibromyoma and pregnancy. Tumour is to left of picture and foetal echoes to right. (Longitudinal section.) (a) Fibromyoma. (b) Foetal echoes. (c) Bowel echoes.

FIG. 13.—Hydatidiform mole distending uterus. Longitudinal section. (a) Mole. (b) Blood clot.

FIG. 14.—Polycystic kidney. (Transverse section.) (a) Polycystic kidney. (b) Bowel echoes.

FIG. 15.—Large fairly homogeneous renal tumour in transverse section. (Spindle-cell sarcoma.) (a) Tumour mass.

FIG. 16.—Goitre straddling trachea. (Transverse section.) (a) Trachea. (b) Goitre.

FIG. 17.—Normal anteverted uterus demonstrated behind a full bladder. (Longitudinal section.) (a) Full bladder. (b) Anteverted uterus. (c) Endometrial cavity. (d) Bowel echoes.

LEGENDS: HARPER AND MacVICAR

FIG. 1.—Transverse section taken 6 cm. below umbilicus. Note typical speckled appearance of the echoes from the vesicles of hydatidiform mole.

FIG. 2.—Transverse section taken 12 cm. below umbilicus. There are definite echoes in a cystic cavity indicative of foetal parts.

FIG. 3.—Longitudinal scan from umbilicus, right, to symphysis pubis, left. The uterus is outlined and the mole can be seen at the fundus with foetal echoes in the lower pole.

seen that the liver has shrunk considerably and the ascitic fluid has gone.

Fig. 9 is from a case of hepatomegaly associated with jaundice which turned out to be drug-induced. In longitudinal section the liver is seen to be much enlarged but clearly transonic and not grossly distorted (right of picture).

Fig. 10, on the other hand, shows a large liver which is much distorted by reflecting material within its substance. (The fine dots are artifacts due to interference.) This was in a case of primary carcinoma of the colon with multiple

FIG. B.—Latest Glasgow echo-sounder, being supplied to the Queen Mother's Hospital.

metastases in the liver. Our experience, however, is not yet great enough to enable us to be dogmatic about the ultrasonic appearances of hepatic metastases. Such cases are not very common in gynaecological practice, and we certainly need more practice and refinement in technique before we can make a confident diagnosis in this respect.

Spleens are much easier. Figs. 11A and B illustrate a case of myelofibrosis in transverse section, looking towards the patient's head. Fig. 11A was taken in 1960, and shows a much-enlarged spleen coming out from underneath the left costal margin. Fig. 11B shows the same case three years later, in May, 1963. The spleen is now very much larger.

We are often confronted with gross abdominal distension in pregnancy, and Fig. 12 shows a case of a fibroid associated with pregnancy, in longitudinal section. The tumour is to the left and the uterus containing the foetal echoes is to the right.

A uterus enormously distended with hydatidiform mole is shown in Fig. 13. We have now diagnosed 20 of these cases by ultrasonic echo-sounding; the appearance of a uterus full of dots is characteristic (MacVicar and Donald, 1963).

We have occasionally had an opportunity to study renal tumours, as, for example, in Fig. 14, which shows an echogram from a man aged 38 with a large renal swelling. The trabeculated appearance and the intervening spaces are from a case of polycystic kidney.

The case of a woman aged 24 who presented with a large renal tumour in the second trimester of pregnancy is demonstrated in Fig. 15. Intravenous pyelography showed

no excretion and only the suggestion of a soft mass. Clinically it was believed to be a large hydronephrosis, but on the strength of this ultrasonogram we reported a non-cystic homogeneous tumour, possibly malignant. At operation it was found to be a large spindle-cell sarcoma.

Within recent months we have been attempting to scan goitres, as shown in Fig. 16, which represents a transverse section of the neck, in which the trachea can be seen in the centre with the goitre straddling it.

We have learned to use the full bladder as a transmitting medium for viewing structures deep in the pelvis, such as a uterus which is not even enlarged, and Fig. 17 is from a patient who was persuaded not to pass urine for some hours. Lying immediately behind the bladder can be seen a normal-sized anteverted uterus, with, apparently, the endometrial cavity depicted down its centre. She had seven weeks' amenorrhoea, but there is no question here of an intrauterine pregnancy.

The latest version of the Glasgow echo-sounder developed for us by Smiths Industrial Division Ltd. is shown in Fig. B and has been ordered by the Scottish Home and Health Department for the new Queen Mother's Hospital, Glasgow, now nearing completion.

Summary

The possible uses of ultrasonic echo-sounding (sonar) in the differential diagnosis of abdominal swellings are illustrated by a number of examples.

These two-dimensional ultrasonograms are obtained by photographing the cathode-ray tube face on which the echoes are displayed. They differ, often characteristically, according to the configuration of the mass, its density, the presence or absence of fluid, and its relationship to gas-containing structures such as bowel.

The help which we have received over the years from the Research Department of Smiths Industrial Division Ltd. is gratefully acknowledged, and we are also indebted to the Scottish Hospital Endowments Research Trust for financial assistance.

REFERENCES

- Donald, I., and Brown, T. G. (1961). *Brit. J. Radiol.*, **34**, 539.
 — MacVicar, J., and Brown, T. G. (1958). *Lancet*, **1**, 1188.
 MacVicar, J. (1962). *Proc. roy. Soc. Med.*, **55**, 638.
 — and Donald, I. (1963). *J. Obstet. Gynaec. Brit. Cwlth.*, **70**, 387.

At the end of February this year the Royal Infirmary at Edinburgh had to close 106 beds, and a committee of inquiry was set up by the South-Eastern Regional Hospital Board to look into the matter. The committee's main findings are as follows. The pattern of nurse staffing in the Infirmary should be reviewed so as to make more use of enrolled nurses and auxiliary staff. Changes should be made to allow more economical use of the nursing staff—for example, some duties may not really require nurses, or at any rate trained nurses, and some services might be provided centrally that would relieve the nurses of particular tasks. When it is proposed to expand services for patients closer attention should be paid to the financial and other resources available (for example, skilled staff or central hospital facilities). There should be changes in the arrangements for admission and in the medical ward organization with the object of evening out (in co-operation with other Edinburgh hospitals) the incidence of emergency admissions and the strain they impose. There is also a more general recommendation that the effect on staffing of the nurse-training requirements needs to be looked at at national level. With a view to implementing the recommendations on nursing the Regional Board are setting up a work-study team to carry out a full review of the use of nurses in the Infirmary.