Review:

DNA-damage response network at the crossroads of cell-cycle checkpoints, cellular senescence and apoptosis*

SCHMITT Estelle¹, PAQUET Claudie¹, BEAUCHEMIN Myriam¹, BERTRAND Richard^{†‡1,2}

(¹Notre Dame Hospital and Montreal Cancer Institute, Research Centre of University of Montreal Hospital Centre (CRCHUM), Montreal (Que) H2L 4M1, Canada)

(²Medicine Department, University of Montreal, Montreal (Que) H3C 3J7, Canada)

[†]E-mail: richard.bertrand@umontreal.ca

Received Apr. 9, 2007; revision accepted Apr. 23, 2007

Abstract: Tissue homeostasis requires a carefully-orchestrated balance between cell proliferation, cellular senescence and cell death. Cells proliferate through a cell cycle that is tightly regulated by cyclin-dependent kinase activities. Cellular senescence is a safeguard program limiting the proliferative competence of cells in living organisms. Apoptosis eliminates unwanted cells by the coordinated activity of gene products that regulate and effect cell death. The intimate link between the cell cycle, cellular senescence, apoptosis regulation, cancer development and tumor responses to cancer treatment has become eminently apparent. Extensive research on tumor suppressor genes, oncogenes, the cell cycle and apoptosis regulatory genes has revealed how the DNA damage-sensing and -signaling pathways, referred to as the DNA-damage response network, are tied to cell proliferation, cell-cycle arrest, cellular senescence and apoptosis. DNA-damage responses are complex, involving “sensor” proteins that sense the damage, and transmit signals to “transducer” proteins, which, in turn, convey the signals to numerous “effector” proteins implicated in specific cellular pathways, including DNA repair mechanisms, cell-cycle checkpoints, cellular senescence and apoptosis. The Bcl-2 family of proteins stands among the most crucial regulators of apoptosis and performs vital functions in deciding whether a cell will live or die after cancer chemotherapy and irradiation. In addition, several studies have now revealed that members of the Bcl-2 family also interface with the cell cycle, DNA repair/recombination and cellular senescence, effects that are generally distinct from their function in apoptosis. In this review, we report progress in understanding the molecular networks that regulate cell-cycle checkpoints, cellular senescence and apoptosis after DNA damage, and discuss the influence of some Bcl-2 family members on cell-cycle checkpoint regulation.

Key words: DNA-damage response network, Cell cycle, Cellular senescence, Apoptosis, Bcl-2 family

doi:10.1631/jzus.2007.B0377

Document code: A

CLC number: R73

INTRODUCTION

The DNA-damage response network is highly complex and involves a multitude of proteins that sense the damage, transduce signals into cells and execute cellular responses. DNA double-strand break “sensor” proteins constitute NBS1/Mre11/Rad50 multiprotein complexes recruited at DNA damage sites, where they create rapidly-expanding nuclear

foci referred to as DNA damage heterochromatin foci, with phosphorylated histone γ -H2AX and other proteins, including BRCA1, 53BP1 and MDC1. The “transducer” protein of DNA double-strand breaks is the phosphoinositide 3-kinase-related kinase ATM, which, in turn, activates multiple “effector” proteins, including p53, Mdm2, and CHK2. DNA-PK, another phosphoinositide 3-kinase-related kinase recruited at DNA damage heterochromatin foci, is involved in the non-homologous end-joining repair pathway, whereas Rad51 is implicated in DNA recombination repair pathways. Although less well-characterized, DNA replication fork stalling and DNA single-strand break

[‡] Corresponding author

* Project supported by the Canadian Institutes of Health Research and the Cancer Research Society, and fellowships by the Health Research Funds of Quebec, Canada

alarms include ATR, ATRIP, RPA, Rad9/Rad1/Hus1, Rad17/RSR and claspin. ATR, a phosphoinositide 3-kinase-related kinase, in turn, activates multiple "effector" proteins, including p53 and CHK1 (reviewed in Bakkenist and Kastan, 2004; Shiloh, 2006). Activation of these "effector" proteins influences cell-cycle progression and arrest, cellular senescence and apoptosis activation.

CELL-CYCLE CHECKPOINT REGULATION

Timely progression through the cell division cycle ensures the correct transmission of genetic information from a cell to its daughters. Once stimulated in proper environment with suitable growth factors, quiescent cells leave a resting phase of the cell cycle, called gap 0 (G0), and enter the gap 1 (G1) phase, prior to the DNA replication or synthesis (S) phase, followed by a second gap (G2) phase, and cell division or mitosis (M). Cell-cycle checkpoints maintain the order and fidelity of cell-cycle events in response to replicative stress and DNA strand breaks (Nigg, 2001). The molecular mechanisms associated with these cell-cycle checkpoints entail the transient inactivation of a series of specific cyclin-dependent kinases (cdks) and their respective regulatory cyclin subunits. During normal phase progression and at key cell-cycle transition phases, specific cdk-cyclin complexes are activated by dephosphorylation, mediated by the dual specificity cdc25 phosphatase family, including cdc25A, -B and -C. The cdk-cyclin complexes influencing G1 progression and G1/S cell-cycle checkpoint are primarily cdk4-cyclinD, cdk6-cyclinD and cdk2-cyclinE (Koff *et al.*, 1992; Sherr and Roberts, 1995). In response to DNA damage, ATM/ATR protein kinases activate the checkpoint serine/threonine kinases, CHK1 and CHK2. To prevent entry into S phase, CHK1 and CHK2 target the cell-cycle regulatory phosphatase cdc25A. Phosphorylation of cdc25A on several residues, including Ser123, leads to its ubiquitin-mediated proteolysis, and to sustained phosphorylation and inhibition of cdk2-cyclinE complexes, that normally promote G1/S transition (Falck *et al.*, 2001). Several small proteins, the cdk-cyclin inhibitors (cdkIs), which bind to and inhibit cdk-cyclin activities, also play critical role in the G1/S cell-cycle checkpoint. The most potent cdkIs

are p21, p27 and p57 for cdk2-cyclinE, and p16 for cdk4-cyclinD or cdk6-cyclinD (Sherr and Roberts, 1995). Activation of these cdkIs causes G1/S cell-cycle arrest after DNA damage (Kohn *et al.*, 1994; Hunter and Pines, 1994; Kastan *et al.*, 1995; Kohn, 1996; Kaufmann and Paules, 1996; Jacks, 1996; Tanaka *et al.*, 1996). Indeed, ATM/ATR and CHK1/CHK2 kinases also target p53. In turn, p53 phosphorylation and stabilization by ATM/ATR and CHK1/CHK2 lead to accumulation of the cdkI p21, which is involved in more sustained G1/S cell-cycle arrest (Shiloh, 2001).

Progression into the S phase and transition from G2 into M are regulated by the serine-threonine cdk2 and cdk1 (cdc2), respectively. Activation of cdk2 and cdk1 (cdc2) requires the association of their positive subunits, cyclinA and cyclinB, respectively, and the phosphorylation of Thr161 by cdk-activating kinase. These cdk-cyclin complexes are also negatively regulated by phosphorylation at Thr14 and Tyr15, which are catalyzed by inhibitory protein kinases, including Wee1, Myt1 and Mik1 (Nigg, 2001; Draetta and Beach, 1988; Draetta *et al.*, 1989; Riabowol *et al.*, 1989; Ducommun *et al.*, 1991; King *et al.*, 1994; Labib *et al.*, 1995; McGowan and Russell, 1995; Mueller *et al.*, 1995; Liu *et al.*, 1997; O'Connell *et al.*, 1997; Booher *et al.*, 1997). During S phase progression and at G2/M phase transition, dephosphorylation of Thr14 and Tyr15 by the cdc25 protein phosphatase cdc25A, -B, and -C, triggers cdk2-cyclinA (intra-S phase) and cdk1 (cdc2)-cyclinB (at G2/M) activation. During the DNA-damage response, activation of ATM/ATR and the downstream checkpoint kinases CHK1/CHK2 leads to the phosphorylation of all 3 cdc25 phosphatases, which creates binding sites for 14-3-3 proteins and sequestration of the phosphatases away from cdk2-cyclinA and cdk1 (cdc2)-cyclinB (Borgne and Meijer, 1996; Peng *et al.*, 1997; Zeng *et al.*, 1998; Sanchez *et al.*, 1997; Matsuoka *et al.*, 1998; Brown *et al.*, 1999; Liu *et al.*, 2000; Furnari *et al.*, 1999; 1997; Lammer *et al.*, 1998; Hofmann *et al.*, 1998; Aguda, 1999; Schmitt *et al.*, 2006). These molecular events provoke S-phase slowing and G2/M cell-cycle arrest. The importance of p53 in the intra-S phase and G2/M checkpoints is uncertain because most cells lacking p53 retain the ability to slow S-phase progression and to arrest at G2/M after DNA damage (O'Connor *et al.*, 1997; O'Connor, 1997;

Gupta *et al.*, 1997; Passalaris *et al.*, 1999; Cliby *et al.*, 2002; Flatten *et al.*, 2005). However, studies have indicated that p53 attenuates the activity of cyclinB1 promoter or promotes the transcriptional activation of the sequestering protein 14-3-3 (Hermeking *et al.*, 1997; Innocente *et al.*, 1999). Others reported that the p53-regulated GADD45 protein associates with and inhibits cdk1 (cdc2)-cyclinB kinase activity in response to DNA damage (Wang *et al.*, 1999a; Zhan *et al.*, 1999). Additional regulators of the G2/M checkpoint are the mitotic serine/threonine Polo-like (PLK) and Aurora-like kinases. PLK1 promotes mitotic entry by phosphorylating and activating cdc25C, and in response to DNA damage, PLK1 is inhibited by an ATM-dependent mechanism. Unlike PLK1, PLK3 is involved in the G2/M arrest by phosphorylating and inhibiting cdc25C (Smits *et al.*, 2000; van Vugt *et al.*, 2001; Xie *et al.*, 2001). AuroraA kinase is normally required for recruitment and activation of cdk1 (cdc2)/cyclinB and commitment of cells into M (Hirota *et al.*, 2003). In response to DNA damage, inhibition of AuroraA kinase is associated to activation of G2/M checkpoint, whereas its overexpression results in checkpoint bypass (Cazales *et al.*, 2005; Marumoto *et al.*, 2002).

Although these pathways leading to cell-cycle delay have been identified during the past few years, the dynamic spatio-temporal coordination and regulation of the DNA-damage response network remain poorly understood. Today, a new challenge in cell-cycle checkpoint biology is to understand how cell-cycle checkpoint pathways are coordinated and organized in space and time, to trigger cellular response to DNA damage (Lukas *et al.*, 2004).

CELLULAR SENESENCE

Recent investigations in the field of cellular senescence have revealed the importance of DNA-damage responses and cell-cycle checkpoints to initiate cell-cycle arrest associated with cellular senescence. Replicative (or the Hayflick limit) and premature senescence share a common signal, now recognized as DNA strand breaks. In replicative senescence, telomere erosion and shortening yield DNA double-strand breaks at chromosome ends that initiate DNA damage responses (Herbig *et al.*, 2004; d'Adda

di Fagagna *et al.*, 2003). In premature senescence, oncogenic activation leads to elevated intracellular levels of reactive species, augmented numbers of active replicons, alterations in DNA replication fork progression, and the appearance of DNA single- and double-strand breaks that initiate DNA-damage responses (Takahashi *et al.*, 2006; Di Micco *et al.*, 2006; Bartkova *et al.*, 2006; Mallette *et al.*, 2007; Hemann and Narita, 2007). Similarly, DNA-damaging chemotherapeutic and radiotherapeutic agents provoke DNA replication fork stalling and/or DNA single- and double-strand breaks with activation of the DNA-damage response network (Holliday and Tarrant, 1972; Kung *et al.*, 1990; Di Leonardo *et al.*, 1994; Waldman *et al.*, 1996; Robles and Adami, 1998; Chang *et al.*, 1999a; 1999b; Suzuki *et al.*, 2001a; Campisi, 2001; Johnstone *et al.*, 2002; Schmitt *et al.*, 2002; Han *et al.*, 2002; Lee and Schmitt, 2003). Indeed, in models of cellular senescence, ATM/ATR mediate the activation of cell-cycle checkpoints associated with cellular senescence, mainly via p53, CHK1 and CHK2, with the participation of p21, p16 and Rb (reviewed in Schmitt, 2007). It has been suggested previously that after genotoxic drug treatment, cell-cycle arrest and senescence development are dependent on the p53/p21 pathway in HCT116 colon cancer cells (Han *et al.*, 2002). Similar observations concerning the importance of the p53/p21 pathway have been reported with other chemotherapeutic agents or irradiation (Di Leonardo *et al.*, 1994; Waldman *et al.*, 1996; Bunz *et al.*, 1998; Linke *et al.*, 1997). Likewise, studies with bleomycin, actinomycin D and cyclophosphamide have revealed that cells respond to these agents by engaging a p53/p16-dependent long-term senescence program (Robles and Adami, 1998; Schmitt *et al.*, 2002). However, others have reported senescence phenotypes in p53-mutated cell lines derived from various types of human solid tumors, including cervical, larynx and colon carcinoma, glioma, osteosarcoma (Chang *et al.*, 1999a) and lymphoma cell lines (Schmitt *et al.*, 2007). Typically, senescent cells are observed by flow cytometry after DNA staining as broader G1 or G2/M peaks, and with the appearance of a senescence-associated β -galactosidase activity (pH 6.0).

The DNA-damage responses induced in the early phase of cellular senescence initiation explain how cells will activate cell-cycle checkpoints and

undergo cell-cycle arrest at specific phases of the cell cycle. However, after DNA damage, cell-cycle arrests do not always lead to cellular senescence. Indeed, cell-cycle arrests often afford cells the opportunity to repair DNA damage before progressing to the next phase of the cycle. In turn, inappropriate progression of damaged cells through the cell cycle is often associated with mitotic catastrophe, apoptosis and cell death. Thus, the stabilization of cell-cycle arrests to irreversible senescence involves other yet unknown critical events. Recently, epigenetic changes, including histone H3 lysine K9 trimethylation (H3K9) associated with gene silencing, have been suggested as critical mechanisms associated with the stabilization of cell-cycle arrest to cellular senescence, as senescent cells show focal histone H3K9 trimethylation. These foci are now referred to as senescence-associated heterochromatin foci (SAHF) (Schmitt, 2007; Strunnikova et al., 2005; Guney and Sedivy, 2006; Zhang et al., 2007). Early evidence indicated that H3K9 trimethylation controls DNA methylation in several model organisms (Tamaru and Selker, 2001; Tamaru et al., 2003; Jackson et al., 2002). Methylation of CpG islands in promoters and locus control regions is strongly associated with H3K9 trimethylation and gene silencing. Indeed, others have indicated that some gene promoters are hypomethylated, whereas others undergo hypermethylation during cellular senescence, suggesting that differential DNA methylation patterns at specific gene promoters and locus control regions may stabilize the senescence phenotype (Tollefsbol and Andrews, 1993; Machwe et al., 2000; Kang et al., 2001; Neumeister et al., 2002; Lopatina et al., 2002; Sun and Arceci, 2005).

BCL-2 FAMILY MEMBERS AND APOPTOSIS

The Bcl-2 gene was first identified at the chromosomal breakpoint of t(14:18)-bearing B cell lymphomas (Tsujimoto et al., 1984; Cleary et al., 1986), and was unexpectedly found to act as a new class of oncogenes that function to prevent apoptosis instead of directly promoting cellular proliferation (Hockenbery et al., 1990; Korsmeyer, 1992; McDonnell et al., 1993; Vaux et al., 1988). Bcl-2 is the founder of a large family of apoptosis regulators that either promote cell survival or facilitate cell death.

The pro-survival members are multidomain proteins that contain BH1, BH2 and BH3 (A1/Bfl-1), BH2 and BH4 (Bcl-xES), and those most similar to Bcl-2 have all 4 BH domains (Bcl-2, Bcl-xL, Mcl-1, Bcl-w, Brag-1, Boo/Diva, Bcl-B/Bcl-2L-10) (Cleary et al., 1986; Boise et al., 1993; Choi et al., 1995; Das et al., 1996; Gibson et al., 1996; Inohara et al., 1998a; Ke et al., 2001; Kozopas et al., 1993; Lin et al., 1993; Schmitt et al., 2004; Song et al., 1999; Zhang et al., 2001). According to their structure and biochemical function, the pro-apoptotic members can be further divided into 2 subgroups: multidomain proteins composed of BH domains 1 to 3 (Bax, Bak, Bok/Mtd, Bcl-rambo), BH2 and BH3 (Bcl-gL) or BH4 and BH3 (Bcl-xS) (Boise et al., 1993; Chittenden et al., 1995a; Farrow et al., 1995; Guo et al., 2001; Hsu et al., 1997a; Inohara et al., 1998b; Kataoka et al., 2001; Kiefer et al., 1995; Oltvai et al., 1993), and BH3-only proteins built of 1 or 2 BH3 domains (Bad, Bid, Bik/Nbk/Blk, Bim/Bod, Hrk/Dp5, Bnip1, Bnip3, Nix, Puma/Bbc3, Noxa, Bcl-gS, Mcl-1s, Map1, Bmf, Spike and ApoL6) (Guo et al., 2001; Bae et al., 2000; Bingle et al., 2000; Boyd et al., 1994; 1995; Chen et al., 1999; Han et al., 1996; 2001; Hegde et al., 1998; Hsu et al., 1998; Imaizumi et al., 1999; Inohara et al., 1997; Matsushima et al., 1998; Nakano and Vausden, 2001; O'Connor et al., 1998; Oda et al., 2000; Puthalakath et al., 2001; Tan et al., 2001; Wang et al., 1996; Yang et al., 1995; Yasuda et al., 1999; Yu et al., 2001; Liu et al., 2005). In addition to the BH domains, some proteins contain a flexible loop, a pore-forming domain and a transmembrane domain at the carboxy terminal extremity (Adams and Cory, 2001; Chao and Korsmeyer, 1998; Reed, 1997; 1998; Vander Heiden and Thompson, 1999; Zamzami et al., 1998; Cory and Adams, 2002).

The various Bcl-2 homologues function, at least in part, through protein-protein interactions, forming a complex dynamic network of homo- and heterooligomers, depending on the cellular environment and their subcellular localization. Mutagenesis experiments have suggested that the dimerization potential of Bcl-2-like proteins is greatly influenced by their specific combination of BH domains (Chittenden et al., 1995b; Kelekar and Thompson, 1998; Yin et al., 1994). Thus, elucidation of the 3-dimensional structure of the anti-apoptotic Bcl-xL has provided a first model in which the α helices located in the BH1, BH2

and BH3 domains create an elongated hydrophobic pocket where a BH3 amphipathic α helix of another Bcl-2-like protein can bind, analogously to a peptide ligand binding to its receptor (Muchmore *et al.*, 1996; Diaz *et al.*, 1997; Sattler *et al.*, 1997). Heterodimerization between pro- and anti-apoptotic molecules may neutralize one another's activity, suggesting that the relative concentration of one faction versus the other greatly influences the susceptibility of cells to a death signal (Oltvai *et al.*, 1993; Sedlak *et al.*, 1995). Because it provides binding ability, the BH3 motif has become a potent mediator of cell death and is often uniquely required for cell-killing activity (Chittenden *et al.*, 1995b; Polster *et al.*, 2001). Two distinct models have been proposed to explain the killing activity of the BH3-only proteins (reviewed in Adams and Cory, 2007). The direct activation model proposes that after a death stimuli, BH3-only proteins promote apoptosis by binding to and inhibiting pro-survival molecules, like Bcl-2, Bcl-xL, Bcl-w, Mcl-1 and A1, or by binding to other death agonists, like the multidomain Bax and Bak proteins, which become activated and exert their pro-apoptotic activities at the mitochondrial level. BH3-only proteins that neutralize pro-survival proteins are referred to as BH3 "enabler or sensitizing" proteins, while those that bind pro-apoptotic Bax or Bak are called BH3 "activator or activating" proteins (Chittenden, 2002; Letai *et al.*, 2002; Kuwana *et al.*, 2005; Kim *et al.*, 2006). The indirect activation model suggests that all BH3-only proteins engage only the pro-survival proteins and act by preventing them from inhibiting Bax or Bak activation (Chen *et al.*, 2005; Willis *et al.*, 2005; 2007).

The p53-mediated pathways have been explored extensively in the context of tumor suppression and their response to DNA-damaging agents (Kohn *et al.*, 1994; Hunter and Pines, 1994; Kastan *et al.*, 1995; Kohn, 1996; Kaufmann and Paules, 1996; Jacks, 1996; Tanaka *et al.*, 1996; O'Connor *et al.*, 1997; O'Connor, 1997; Gupta *et al.*, 1997; Passalaris *et al.*, 1999; Hermeking *et al.*, 1997; Innocente *et al.*, 1999; Wang *et al.*, 1999a; Zhan *et al.*, 1999; Johnstone *et al.*, 2002; Levine, 1997; Lowe *et al.*, 1993; 1994; Gottlieb and Oren, 1998). At the level of transcription, multiple p53 target genes, including those of the Bcl-2 family, play crucial roles in promoting apoptosis induced by genotoxic agents. Among them, the Bax gene was one of the first p53-dependent apoptotic targets to be

identified (Miyashita and Reed, 1995; Miyashita *et al.*, 1994); then, other members of the Bcl-2 family were reported to be p53-downstream targets, such as the BH3-only members Bid (Sax *et al.*, 2002), Puma (Nakano and Voutsden, 2001; Yu *et al.*, 2001; Villunger *et al.*, 2003) and Noxa (Oda *et al.*, 2000; Shibue *et al.*, 2003). More recently, others have revealed that p53 itself targets mitochondria, where it associates with and activates the multidomain pro-apoptotic Bax and Bak at the protein level (Chipuk *et al.*, 2004; Leu *et al.*, 2004; Perfettini *et al.*, 2004; Mihara *et al.*, 2003; Dumont *et al.*, 2003; Chipuk *et al.*, 2005).

Much less is known about the effector genes involved in p53-independent responses after DNA damage. However, most human tumors lose the expression of a functional p53, and they likely represent difficult targets in cancer therapies. Paradoxically, several observations have disclosed that cells lacking the ability to transduce a cell-cycle checkpoint, as a consequence of targeted disruption of p53 or mutated p53, could display hypersensitivity to DNA-damaging treatments, including DNA topoisomerase I and II inhibitors, and undergo apoptosis (Gupta *et al.*, 1997; Waldman *et al.*, 1996; Kaufmann, 1989; 1998; Slichenmyer *et al.*, 1993; Thomas *et al.*, 1996; Bertrand *et al.*, 1991; 1993; Solary *et al.*, 1992; Dubrez *et al.*, 1995; 1996; Shimizu and Pommier, 1997; Kaufmann and Earnshaw, 2000; Droin *et al.*, 1998).

p73 and p63, other p53 family members that can be activated when cells are damaged, also contribute to programmed cell death induction, in the presence or absence of p53 after DNA damage (Jost *et al.*, 1997; Yang *et al.*, 1998; Yuan *et al.*, 1999; Gong *et al.*, 1999; Agami *et al.*, 1999; Costanzo *et al.*, 2002; Flores *et al.*, 2002; Bergamaschi *et al.*, 2003; Freebern *et al.*, 2003; Nozell *et al.*, 2003). While the p73 and p63 functional domains necessary for mediating apoptosis have been characterized, much less is known about the p73 or p63 target genes of Bcl-2 family members (Jost *et al.*, 1997; Yang *et al.*, 1998; Yuan *et al.*, 1999; Gong *et al.*, 1999; Agami *et al.*, 1999; Costanzo *et al.*, 2002; Flores *et al.*, 2002; Bergamaschi *et al.*, 2003; Freebern *et al.*, 2003; Nozell *et al.*, 2003; Fontemaggi *et al.*, 2002). Recently, Melino *et al.* (2004) reported that p73 isoform overexpression can directly induce Puma transactivation.

E2F1 transcription factor has been shown to

participate in the induction of apoptosis in the presence or absence of p53 (Hunt *et al.*, 1997; Phillips *et al.*, 1997; Holmberg *et al.*, 1998). Overexpression of E2F1 itself can directly activate p73 transcription (Stiewe and Pützer, 2000) and regulate the expression of several genes involved in apoptosis, including the multi-domain pro-apoptotic Bak, the BH3-only proteins Bid, Bad, Bim, Noxa, Puma, Hrk, the adaptor protein Apaf-1 and caspase-3 and -7 (Muller *et al.*, 2001; Stanelle *et al.*, 2002; Pediconi *et al.*, 2003; Hershko and Ginsberg, 2004). Some of these targeted genes are also regulated by E2F1 following cisplatin or doxorubicin treatment of p53-deficient mouse embryo fibroblasts (Hershko and Ginsberg, 2004), highlighting its importance for drug responses.

NF κ B activation after DNA topoisomerase inhibitor treatment was shown to be p53-independent (Piret *et al.*, 1999), but reported to be essential in p53-mediated cell death induced by DNA damage, highlighting its important role (Ryan *et al.*, 2000). Activation of NF κ B and c-Jun/AP-1 contributes either to protect cells from death or activates cell death, depending on the cellular environment (Foo and Nolan, 1999; Mercurio and Manning, 1999; Grumont *et al.*, 1998), and a growing number of genes related to apoptosis have been demonstrated to be induced by NF κ B and c-Jun/AP-1 in response to multiple stress-inducing agents, including DNA damage (Chu *et al.*, 1997; Stehlik *et al.*, 1998; Wang *et al.*, 1998; 1999b; Tamatani *et al.*, 1999; Zong *et al.*, 1999; Stroka *et al.*, 1999).

Bcl-2 family members generally localize to separate subcellular compartments in the absence of a death signal. Anti-apoptotic members, like Bcl-2 and Bcl-xL, usually reside on the cytoplasmic face of the outer mitochondrial membrane (OMM), endoplasmic reticulum and nuclear envelope (Hockenbery *et al.*, 1990; Krajewski *et al.*, 1993; Zamzami *et al.*, 1996). In contrast, death-inducing members, such as Bax, Bad and Bid, are rather free in unstimulated cells, except for the multidomain Bak which seems to constitutively reside in the OMM (Gross *et al.*, 1998; Korsmeyer *et al.*, 2000; Hsu *et al.*, 1997b). In turn, the BH3-only Bim and Bmf are linked to cytoskeleton components via binding to the large microtubular dynein motor complex and myosin V actin motor complex, respectively (Puthalakath *et al.*, 1999; 2001). After a death signal, multidomain pro-apoptotic pro-

teins, like Bax, undergo a conformational change which gives them the capacity to target and insert into intracellular membranes, especially in the mitochondria. The transmembrane domain appears to be important in conferring membrane-docking capacity to both pro- and anti-apoptotic members as well as in providing protein stability, even though reports indicate that deletion of this domain does not abrogate the function of most Bcl-2-like proteins (Borner *et al.*, 1994; Nguyen *et al.*, 1994).

Regulation of mitochondrial membrane permeabilization is the major means by which Bcl-2-like proteins exert their apoptosis regulatory effect. As a site of convergence for multiple death-inducing stimuli, the mitochondria are a pivotal decision centre that controls life and death by releasing apoptogenic factors in the cytosol. These death-inducing molecules are located within the mitochondrial intermembrane space and include cytochrome c (Kluck *et al.*, 1997; Yang *et al.*, 1997), DIABLO/Smac, a factor that promotes caspase activation by its effect on IAPs (inhibitor of apoptosis proteins) (Du *et al.*, 2000; Verhagen *et al.*, 2000), the nuclease activator AIF (apoptosis-inducing factor) (Susin *et al.*, 1996; 1999a), Endo G (an apoptotic DNase) (Li *et al.*, 2001; Parrish *et al.*, 2001), HtrA2/Omi (an inhibitor of IAPs that also contains pro-apoptotic serine protease activity) (Suzuki *et al.*, 2001b), and some pro-caspases (Krajewski *et al.*, 1999; Mancini *et al.*, 1998; Qin *et al.*, 2001; Susin *et al.*, 1999b). Permeabilization of the OMM is usually followed by caspase-9 activation and irreversible apoptosis. Bcl-2 family members may control the mitochondrial permeabilization phase (Desagher and Martinou, 2000) via 3 prevailing models: first, by forming autonomous channels in the OMM; second, by interfering with mitochondrial permeability transition pores (mPTP); and, finally, by promoting dynamic alterations in the structure of the lipid bilayer (Gross *et al.*, 1998; Kluck *et al.*, 1997; Yang *et al.*, 1997; Antonsson *et al.*, 1997; 2000; Minn *et al.*, 1997; Chou *et al.*, 1999; McDonnell *et al.*, 1999; Schendel *et al.*, 1998; Schlesinger *et al.*, 1997; Bossy-Wetzel *et al.*, 1998; Vander Heiden *et al.*, 1997; Saito *et al.*, 2000; Pavlov *et al.*, 2001; Basañez *et al.*, 1999; Kudla *et al.*, 2000; Epand *et al.*, 2002; Kuwana *et al.*, 2002). Anti-apoptotic proteins seem to exert their function in part by inhibiting the formation of death-inducing pores by pro-apoptotic proteins, and a

study has reported that the Bcl-2 channel could potentially inhibit apoptosis by evoking H^+ efflux from the mitochondria, preserving mitochondrial transmembrane potential ($\Delta\Psi_m$) (Shimizu *et al.*, 1998). Anti-apoptotic proteins may also protect cells from death by inhibiting the production of reactive oxygen species (Hockenbery *et al.*, 1993; Kane *et al.*, 1993), by preventing intracellular acidification (Gottlieb *et al.*, 1996), by increasing mitochondrial Ca^{2+} capacity (Susin *et al.*, 1996; Shimizu *et al.*, 1998), and by providing overall stabilization of the membrane into which they are inserted (Vander Heiden *et al.*, 1997; Zamzami *et al.*, 1995). These pro-survival mechanisms are not well defined and may or may not involve the pore-forming capacity of anti-apoptotic proteins.

In the first model, Bcl-2-like proteins modulate OMM permeabilization without altering mitochondrial function, which means that $\Delta\Psi_m$ is maintained as well as ATP production. This is crucial because in the absence of dATP or ATP, the apoptosome cannot form, caspase-9 is not activated, and the cells tend to die by necrosis rather than by apoptosis (Chautan *et al.*, 1999; Nicotera *et al.*, 1999). The second model of OMM permeabilization involves mPTP, or megachannels, which form at contact sites between the outer and inner mitochondrial membranes (Zamzami *et al.*, 1998; Zoratti and Szabo, 1995). Its core components are the voltage-dependent anion channel 1 (VDAC1), the adenine nucleotide translocator (ANT), the peripheral benzodiazepine receptor, the mitochondrial matrix cyclophilin as well as hexokinase and creatine kinase, which all create a large conductance pore permeable to solute with a molecular mass of ≤ 1.5 kDa (Bernardi *et al.*, 1999; Crompton, 1999). mPTP opening results in mitochondrial depolarization and the subsequent dissipation of $\Delta\Psi_m$, uncoupling of oxidative phosphorylation, and swelling of the mitochondrial matrix (Bernardi *et al.*, 1999). OMM rupture may eventually occur as a consequence of matrix expansion, releasing the intermembrane space contents (cytochrome c, DIABLO/Smac, Omni/HtrA2, AIF) into the cytosol in a nonspecific manner (Harris and Thompson, 2000; Zamzami and Kroemer, 2001). Interactions between some members of the Bcl-2 family and the components of mPTP have been proposed by several studies. Pro-apoptotic molecules like Bax could bind to ANT and promote mPTP

opening, an effect prevented by the anti-apoptotic Bcl-2 (Marzo *et al.*, 1998). The multidomains Bax and Bak could also interact with the VDAC1, enhancing its channel activity to allow passage of cytochrome c, while the BH4 domain of anti-apoptotic Bcl-2 and Bcl-xL rather stimulates closure of the channel (Shimizu *et al.*, 1999; 2000; Shimizu and Tsujimoto, 2000). Recently, Sugiyama *et al.* (2002) have shown that the BH3-only Bim, but not Bid, may also interact with the VDAC1 and promote its opening, which results in loss of $\Delta\Psi_m$ and cytochrome c release. By interacting with mPTP, pro-apoptotic proteins thus appear to be capable of enhancing the basal permeability of mPTP, allowing molecules larger than 1.5 kDa to diffuse through.

BCL-2 FAMILY MEMBERS AND CELL-CYCLE CHECKPOINTS

In addition to their central role in controlling cell death, numerous studies have linked some Bcl-2-like family members with cell cycle progression (summarized in Fig.1). First, Bcl-2 itself was shown to slow entry from the quiescent G0 into the G1 phase of the cell cycle prior to DNA replication in multiple cell lineages and transgenic mice (Fig.1). In contrast, Bcl-2^{-/-} knockout cells enter the S phase more quickly (Borner, 1996; Brady *et al.*, 1996; Vairo *et al.*, 1996; Mazel *et al.*, 1996; Linette *et al.*, 1996; O'Reilly *et al.*, 1996; 1997a; 1997b; Huang *et al.*, 1997; Knowlton *et al.*, 1998; Ogilvy *et al.*, 1999; Gillardon *et al.*, 1999; Domen *et al.*, 2000; Vail *et al.*, 2002). This effect of Bcl-2 on cell proliferation is genetically distinct from its function on apoptosis (Borner, 1996; Brady *et al.*, 1996; Vairo *et al.*, 1996; Huang *et al.*, 1997; Knowlton *et al.*, 1998). Mutation of a tyrosine residue (Tyr28) within the BH4 domain has no impact on the ability of Bcl-2 to inhibit apoptosis and does not prevent its heterodimerization with death-inducing family members, including Bax, Bak, Bad, and Bik. However, when starved quiescent cells expressing the mutated protein are stimulated by serum, they re-enter the cell cycle faster than those expressing wild-type Bcl-2 protein (Huang *et al.*, 1997).

Mcl-1, another Bcl-2 homologue known to function as an anti-apoptotic protein (Kozopas *et al.*, 1993), inhibits cell-cycle progression through the S phase of

the cell cycle (Fig.1). The cell-cycle regulatory function of Mcl-1 is partially mediated through its interaction with proliferating cell nuclear antigen (PCNA), a cell cycle regulator that plays a critical role in DNA replication (Fujise *et al.*, 2000). The binding motif for PCNA resides between the 221st to 228th amino acids of Mcl-1. Mutations introduced within this region do not alter the anti-apoptotic function of Mcl-1, but impede its binding to PCNA and significantly reduce its effect on cell cycle progression (Fujise *et al.*, 2000). Thus, this effect of Mcl-1 on cell proliferation is genetically distinct from its function on apoptosis. More recently, others have revealed that a proteolytic fragment of Mcl-1 regulates cell growth via interaction with cdk1 (cdc2) (Jamil *et al.*, 2005).

Fig.1 Schematic representation of the effects of Bcl-2, Mcl-1, Bid and Bcl-xL on cell cycle

Microtubule inhibitors widely used in cancer chemotherapy, including paclitaxel, taxotere, vinblastine and vincristine, induce mitotic arrest linked with Bcl-2, Bcl-xL and Mcl-1 phosphorylation (Haldar *et al.*, 1995; 1996; Blagosklonny *et al.*, 1997; Poruchynsky *et al.*, 1998; Domina *et al.*, 2000). However, studies have revealed that Bcl-2 and Bcl-xL phosphorylation is tightly coupled with normal mitotic events (Domina *et al.*, 2000; Ling *et al.*, 1998; Scatena *et al.*, 1998; Fan *et al.*, 2000a). Bcl-2 phosphorylation multisites (Ser70, Ser87, Thr69) are located within the unstructured loop domain of the protein, a region not essential for its anti-apoptotic function (Chang *et al.*, 1997; Haldar *et al.*, 1998; Fang *et al.*, 1998; Srivastava *et al.*, 1999; Yamamoto *et al.*, 1999). Multiple kinases have been proposed to phosphorylate Bcl-2 (and presumably Bcl-xL), including an Ask1/Jun-N-terminal protein kinase 1 pathway, normally activated at G2/M (Fan *et al.*, 2000a; 2000b; Yamamoto *et al.*, 1999; Blagosklonny *et al.*, 1996; Ruvolo *et al.*, 1998; Srivastava *et al.*, 1998; Maundrell *et al.*, 1997; Kharbanda *et al.*, 2000).

In addition, Kharbanda *et al.* (2000) have reported that Bcl-xL is phosphorylated on Thr47, within the unstructured loop domain, and on Thr115, adjacent to the $\alpha 3$ helix of Bcl-xL, in response to ionizing radiation.

Recent works have also indicated a role for Bid in the intra-S phase checkpoint after replicative stress in response to DNA-damaging agents (Fig.1). This role of Bid is mediated through its phosphorylation at Ser78 and Ser61/64, by the DNA-damage kinase ATM (Zinkel *et al.*, 2005; Kamer *et al.*, 2005). More recently, we showed that in addition to its mitochondrial effect that delays apoptosis, Bcl-xL co-localizes and binds to cdk1 (cdc2) during the G2/M cell-cycle checkpoint, and its overexpression stabilizes a G2/M-arrest senescence program in surviving cells after DNA damage (Fig.1). Bcl-xL potentially inhibits cdk1 (cdc2) kinase activities, which is reversible by a synthetic peptide between the 41st to 60th amino acids surrounding the Thr47 and Ser62 phosphorylation sites, and Asn52 deamidation site, within the flexible loop domain of Bcl-xL. A mutant deleted of this region does not alter the anti-apoptotic function of Bcl-xL, but impedes its effect on cdk1 (cdc2) activities and on the G2/M-arrest senescence program after DNA damage. The nuclear interaction of Bcl-xL and cdk1 (cdc2) suggests that Bcl-xL is coupled to the stabilization of a cell-cycle checkpoint induced by DNA damage, and this effect is genetically distinct from its function on apoptosis (Schmitt *et al.*, 2007).

CONCLUSION

Recent progress in understanding the molecular mechanisms of DNA-damage response illustrates how the key components interplay into different processes to protect genomic integrity of cells. Already, strong efforts are being made by several laboratories to develop new compounds targeting cell-cycle progression, cell-cycle checkpoint and apoptosis regulatory proteins. Several natural products and synthetic chemicals occupied the BH3-binding site of anti-apoptotic Bcl-2 family members and some of these compounds promote apoptosis of tumor cells (summarized in Table 1) (reviewed in Reed, 2006). Knowing more about the molecular interdependence of the DNA-damage response network will certainly be helpful in designing rational therapeutic protocols

with these new compounds in combination with traditional chemotherapy. Insights into the dynamic nature of these processes and the spatio-temporal orchestration of their regulation will suggest novel potential therapeutic strategies to prevent and treat cancer in the near future.

Table 1 Natural compounds and synthetic chemicals targeting antiapoptotic Bcl-2 family members*

Compounds	Organization	Status
Genasense (antisense)	Genta, Inc.	Phase III
Gossypol	Ascenta (NCI-USA)	Phase I/II
Polyphenol E	Mayo Clinic (Mitsui Norin)	Phase I
GX-15-070	Gemin X (Canada)	Phase I
HA14-1 analogs	Raylight Chemokine Pharm. Inc.	Preclinical
ABT-737	Abbott Laboratories	Preclinical
Apogossypol	Burnham Institute/NCI-USA	Preclinical
BH3I-1/-2	Havard University	Preclinical
Antimycin A3	University of Washington	Preclinical
Compound 6	University of Michigan	Preclinical
Terphenyl derivative	Yale University	Preclinical

* From (Reed, 2006)

ACKNOWLEDGEMENT

The editorial work of Mr. Ovid Da Silva of the Research Support Office, CHUM, is much appreciated.

References

- Adams, J.M., Cory, S., 2001. Life-or-death decisions by the Bcl-2 protein family. *Trends Biochem. Sci.*, **26**(1):61-66. [doi:10.1016/S0968-0004(00)01740-0]
- Adams, J.M., Cory, S., 2007. The Bcl-2 apoptotic switch in cancer development and therapy. *Oncogene*, **26**(9):1324-1337. [doi:10.1038/sj.onc.1210220]
- Agami, R., Blandino, G., Oren, M., Shaul, Y., 1999. Interaction of c-Abl and p73^a and their collaboration to induce apoptosis. *Nature*, **399**(6738):809-813. [doi:10.1038/21697]
- Aguda, B.D., 1999. A quantitative analysis of the kinetics of the G(2) DNA damage checkpoint system. *Proc. Natl. Acad. Sci. (USA)*, **96**(20):11352-11357. [doi:10.1073/pnas.96.20.11352]
- Antonsson, B., Conti, F., Ciavatta, A., Montessuit, S., Lewis, S., Martinou, I., Bernasconi, L., Bernard, A., Mermod, J.J., Mazzei, G., et al., 1997. Inhibition of Bax channel-forming activity by Bcl-2. *Science*, **277**(5324):370-372. [doi:10.1126/science.277.5324.370]
- Antonsson, B., Montessuit, S., Lauper, S., Eskes, R., Martinou, J.C., 2000. Bax oligomerization is required for channel-forming activity in liposomes and to trigger cytochrome c release from mitochondria. *Biochem. J.*, **345**(2):271-278. [doi:10.1042/0264-6021:3450271]
- Bae, J., Leo, C.P., Hsu, S.Y., Hsueh, A.J., 2000. Mcl-1S, a splicing variant of the antiapoptotic Bcl-2 family member Mcl-1, encodes a proapoptotic protein possessing only the BH3 domain. *J. Biol. Chem.*, **275**(33):25255-25261. [doi:10.1074/jbc.M909826199]
- Bakkenist, C.J., Kastan, M.B., 2004. Initiating cellular stress responses. *Cell*, **118**(1):9-17. [doi:10.1016/j.cell.2004.06.023]
- Bartkova, J., Rezaei, N., Liontos, M., Karakaidos, P., Kletsas, D., Issaeva, N., Vassiliou, L.V., Kolettas, E., Niforou, K., Zoumpourlis, V.C., et al., 2006. Oncogene-induced senescence is part of the tumorigenesis barrier imposed by DNA damage checkpoints. *Nature*, **444**(7119):633-637. [doi:10.1038/nature05268]
- Basañez, G., Nechushtan, A., Drozhinin, O., Chanturiya, A., Choe, E., Tutt, S., Wood, K.A., Hsu, Y., Zimmerberg, J., Youle, R.J., 1999. Bax, but not Bcl-x_L, decreases the lifetime of planar phospholipid bilayer membranes at subnanomolar concentrations. *Proc. Natl. Acad. Sci. (USA)*, **96**(10):5492-5497. [doi:10.1073/pnas.96.10.5492]
- Bergamaschi, D., Gasco, M., Hiller, L., Sullivan, A., Syed, N., Trigiante, G., Yulug, I., Merlano, M., Numico, G., Comino, A., et al., 2003. p53 polymorphism influences response in cancer chemotherapy via modulation of p73-dependent apoptosis. *Cancer Cell*, **3**(4):387-402. [doi:10.1016/S1535-6108(03)00079-5]
- Bernardi, P., Scorrano, L., Colonna, R., Petronilli, V., Di Lisa, F., 1999. Mitochondria and cell death. Mechanistic aspects and methodological issues. *Eur. J. Biochem.*, **264**(3):687-701. [doi:10.1046/j.1432-1327.1999.00725.x]
- Bertrand, R., Sarang, M., Jenkin, J., Kerrigan, D., Pommier, Y., 1991. Differential induction of secondary DNA fragmentation by topoisomerase II inhibitors in human tumor cell lines with amplified c-myc expression. *Cancer Res.*, **51**:6280-6285.
- Bertrand, R., Solary, E., Jenkins, J., Pommier, Y., 1993. Apoptosis and its modulation in human promyelocytic HL-60 cells treated with DNA topoisomerase I and II inhibitors. *Exp. Cell Res.*, **207**(2):388-397. [doi:10.1006/excr.1993.1206]
- Bingle, C.D., Craig, R.W., Swales, B.M., Singleton, V., Zhou, P., Whyte, M.K., 2000. Exon skipping in Mcl-1 results in a bcl-2 homology domain 3 only gene product that promotes cell death. *J. Biol. Chem.*, **275**(29):22136-22146. [doi:10.1074/jbc.M909572199]
- Blagosklonny, M.V., Schulte, T., Nguyen, P., Trepel, J., Neckers, L.M., 1996. Taxol-induced apoptosis and phosphorylation of Bcl-2 protein involves c-Raf-1 and represents a novel c-Raf-1 signal transduction pathway. *Cancer Res.*, **56**:1851-1854.
- Blagosklonny, M.V., Giannakakou, P., el Deiry, W.S., Kingston, D.G., Higgs, P.I., Neckers, L., Fojo, T., 1997. Raf-1/Bcl-2

- phosphorylation: a step from microtubule damage to cell death. *Cancer Res.*, **57**:130-135.
- Boise, L.H., Gonzalez-Garcia, M., Postema, C.E., Ding, L., Lindsten, T., Turka, L.A., Mao, X., Nunez, G., Thompson, C.B., 1993. Bcl-x, a bcl-2-related gene that functions as a dominant regulator of apoptotic cell death. *Cell*, **74**(4): 597-608. [doi:10.1016/0092-8674(93)90508-N]
- Booher, R.N., Holman, P.S., Fattaey, A., 1997. Human Myt1 is a cell cycle-regulated kinase that inhibits Cdc2 but not Cdk2 activity. *J. Biol. Chem.*, **272**(35):22300-22306. [doi:10.1074/jbc.272.35.22300]
- Borgne, A., Meijer, L., 1996. Sequential dephosphorylation of p34^{cdc2} on Thr-14 and Tyr-15 at the prophase/metaphase transition. *J. Biol. Chem.*, **271**(44):27847-27854. [doi:10.1074/jbc.271.44.27847]
- Borner, C., 1996. Diminished cell proliferation associated with the death-protective activity of Bcl-2. *J. Biol. Chem.*, **271**:12695-12698.
- Borner, C., Martinou, I., Mattmann, C., Irmeler, M., Schaefer, E., Martinou, J.C., Tschopp, J., 1994. The protein bcl-2 alpha does not require membrane attachment, but two conserved domains to suppress apoptosis. *J. Cell. Biol.*, **126**(4):1059-1068. [doi:10.1083/jcb.126.4.1059]
- Bossy-Wetzel, E., Newmeyer, D.D., Green, D.R., 1998. Mitochondrial cytochrome C release in apoptosis occurs upstream of DEVD-specific caspase activation and independently of mitochondrial transmembrane depolarization. *EMBO J.*, **17**(1):37-49. [doi:10.1093/emboj/17.1.37]
- Boyd, J., Malstrom, S., Subramanian, T., Venkatesh, L.K., Schaeper, U., Elangovan, B., D'Sa-Eipper, C., Chinnadurai, G., 1994. E1B 19 kDa and Bcl-2 proteins interact with a common set of cellular proteins. *Cell*, **79**(2):341-351. [doi:10.1016/0092-8674(94)90202-X]
- Boyd, J.M., Gallo, G.J., Elangovan, B., Houghton, A.B., Malstrom, S., Avery, B.J., Ebb, R.G., Subramanian, T., Chittenden, T., Lutz, R.J., et al., 1995. Bik, a novel death-inducing protein shares a distinct sequence motif with Bcl-2 family proteins and interacts with viral and cellular survival-promoting proteins. *Oncogene*, **11**:1921-1928.
- Brady, H.J.M., Gil-Gómez, G., Kirberg, J., Berns, A.J.M., 1996. Bax-a perturbs T cell development and affects cell cycle entry of T cells. *EMBO J.*, **15**:6991-7001.
- Brown, A.L., Lee, C.H., Schwarz, J.K., Mitiku, N., Pwnica-Worms, H., Chung, J.H., 1999. A human Cds1-related kinase that functions downstream of ATM protein in the cellular response to DNA damage. *Proc. Natl. Acad. Sci. (USA)*, **96**(7):3745-3750. [doi:10.1073/pnas.96.7.3745]
- Bunz, F., Dutriaux, A., Lengauer, C., Waldman, T., Zhou, S., Brown, J.P., Sedivy, J.M., Kinzler, K.W., Vogelstein, B., 1998. Requirement for p53 and p21 to sustain G2 arrest after DNA damage. *Science*, **282**(5393):1497-1501. [doi:10.1126/science.282.5393.1497]
- Campisi, J., 2001. Cellular senescence as a tumor-suppressor mechanism. *Trends Cell Biol.*, **11**(11):S27-S31. [doi:10.1016/S0962-8924(01)02151-1]
- Cazales, M., Schmitt, E., Montembault, E., Dozier, C., Prigent, C., Ducommun, B., 2005. CDC25B phosphorylation by Aurora-A occurs at the G2/M transition and is inhibited by DNA damage. *Cell Cycle*, **4**:1233-1238.
- Chang, B.S., Minn, A.J., Muchmore, S.W., Fesik, S.W., Thompson, C.B., 1997. Identification of a novel regulatory domain in Bcl-X(L) and Bcl-2. *EMBO J.*, **16**(5):968-977. [doi:10.1093/emboj/16.5.968]
- Chang, B.D., Broude, E.V., Dokmanovic, M., Zhu, H., Ruth, A., Xuan, Y., Kandel, E.S., Lausch, E., Christov, K., Roninson, I.B., 1999a. A senescence-like phenotype distinguishes tumor cells that undergo terminal proliferation arrest after exposure to anticancer agents. *Cancer Res.*, **59**:3761-3767.
- Chang, B.D., Xuan, Y., Broude, E.V., Zhu, H., Schott, B., Fang, J., Roninson, I.B., 1999b. Role of p53 and p21waf1/cip1 in senescence-like terminal proliferation arrest induced in human tumor cells by chemotherapeutic drugs. *Oncogene*, **18**(34):4808-4818. [doi:10.1038/sj.onc.1203078]
- Chao, D.T., Korsmeyer, S.J., 1998. Bcl-2 family: regulators of cell death. *Annu. Rev. Immunol.*, **16**(1):395-419. [doi:10.1146/annurev.immunol.16.1.395]
- Chautan, M., Chazal, G., Cecconi, F., Gruss, P., Golstein, P., 1999. Interdigital cell death can occur through a necrotic and caspase-independent pathway. *Curr. Biol.*, **9**(17): 967-970. [doi:10.1016/S0960-9822(99)80425-4]
- Chen, G., Cizeau, J., Vande Velde, C., Park, J.H., Bozek, G., Bolton, J., Shi, L., Dubik, D., Greenberg, A., 1999. Nix and Nip3 form a subfamily of pro-apoptotic mitochondrial proteins. *J. Biol. Chem.*, **274**(1):7-10. [doi:10.1074/jbc.274.1.7]
- Chen, L., Willis, S.N., Wei, A., Smith, B.J., Fletcher, J.I., Hinds, M.G., Colman, P.M., Day, C.L., Adams, J.M., Huang, D.C., 2005. Differential targeting of prosurvival Bcl-2 proteins by their BH3-only ligands allows complementary apoptotic function. *Mol. Cell*, **17**(3):393-403. [doi:10.1016/j.molcel.2004.12.030]
- Chipuk, J.E., Kuwana, T., Bouchier-Hayes, L., Droin, N.M., Newmeyer, D.D., Schuler, M., Green, D.R., 2004. Direct activation of Bax by p53 mediates mitochondrial membrane permeabilization and apoptosis. *Science*, **303**(5660): 1010-1014. [doi:10.1126/science.1092734]
- Chipuk, J.E., Bouchier-Hayes, L., Kuwana, T., Newmeyer, D.D., Green, D.R., 2005. Puma couples the nuclear and cytoplasmic proapoptotic function of p53. *Science*, **309**(5741):1732-1735. [doi:10.1126/science.1114297]
- Chittenden, T., 2002. BH3 domains: intracellular death-ligands critical for initiating apoptosis. *Cancer Cell*, **2**(3):165-166. [doi:10.1016/S1535-6108(02)00128-9]
- Chittenden, T., Harrington, E.A., O'Connor, R., Flemington, C., Lutz, R.J., Evan, G.I., Guild, B.C., 1995a. Induction of apoptosis by the Bcl-2 homologue Bak. *Nature*, **374**(6524):733-736. [doi:10.1038/374733a0]
- Chittenden, T., Flemington, C., Houghton, A.B., Ebb, R.G., Gallo, G.J., Elangovan, B., Chinnadurai, G., Lutz, R.J., 1995b. A conserved domain in Bak, distinct from BH1

- and BH2, mediates cell death and protein binding functions. *EMBO J.*, **14**:5589-5596.
- Choi, S.S., Park, I.C., Yun, J.W., Sung, Y.C., Hong, S.I., Shin, H.S., 1995. A novel Bcl-2 related gene, Bfl-1, is overexpressed in stomach cancer and preferentially expressed in bone marrow. *Oncogene*, **11**:1693-1698.
- Chou, J.J., Li, H., Salvesen, G.S., Yuan, J., Wagner, G., 1999. Solution structure of BID, an intracellular amplifier of apoptotic signaling. *Cell*, **96**(5):615-624. [doi:10.1016/S0092-8674(00)80572-3]
- Chu, Z.L., McKinsey, T.A., Liu, L., Gentry, J.J., Malim, M.H., Ballard, D.W., 1997. Suppression of tumor necrosis factor-induced cell death by inhibitor of apoptosis C-Iap2 is under NF-kappa-B control. *Proc. Natl. Acad. Sci. (USA)*, **94**(19):10057-10062. [doi:10.1073/pnas.94.19.10057]
- Cleary, M.L., Smith, S.D., Sklar, J., 1986. Cloning and structural analysis of cDNAs for *bcl-2* and a hybrid *bcl-2*/immunoglobulin transcript resulting from the t(14;18) translocation. *Cell*, **47**(1):19-28. [doi:10.1016/0092-8674(86)90362-4]
- Cliby, W.A., Lewis, K.A., Lilly, K.K., Kaufmann, S.H., 2002. S phase and G2 arrests induced by topoisomerase I poisons are dependent on ATR kinase function. *J. Biol. Chem.*, **277**(2):1599-1606. [doi:10.1074/jbc.M106287200]
- Cory, S., Adams, J.M., 2002. The bcl2 family: regulators of the cellular life-or-death switch. *Nat. Rev. Cancer*, **2**(9):647-656. [doi:10.1038/nrc883]
- Costanzo, A., Merlo, P., Pediconi, N., Fulco, M., Sartorelli, V., Cole, P.A., Fontemaggi, G., Fanciulli, M., Schiltz, L., Blandino, G., Balsano, C., Levrero, M., 2002. DNA damage-dependent acetylation of p73 dictates the selective activation of apoptotic target genes. *Mol. Cell*, **9**(1):175-186. [doi:10.1016/S1097-2765(02)00431-8]
- Crompton, M., 1999. The mitochondrial permeability transition pore and its role in cell death. *Biochem. J.*, **341**(2):233-249. [doi:10.1042/0264-6021:3410233]
- d'Adda di Fagnana, F., Reaper, P.M., Clay-Farrace, L., Fiegler, H., Carr, P., von Zglinicki, T., Saretzki, G., Carter, N.P., Jackson, S.P., 2003. A DNA damage checkpoint response in telomere-initiated senescence. *Nature*, **426**(6963):194-198. [doi:10.1038/nature02118]
- Das, R., Reddy, E.P., Chatterjee, D., Andrews, D.W., 1996. Identification of a novel Bcl-2 related gene, Brag-1, in human glioma. *Oncogene*, **12**:947-951.
- Desagher, S., Martinou, J.C., 2000. Mitochondria as the central control point of apoptosis. *Trends Cell Biol.*, **10**(9):369-377. [doi:10.1016/S0962-8924(00)01803-1]
- Di Leonardo, A., Linke, S.P., Clarkin, K., Wahl, G.M., 1994. DNA damage triggers a prolonged p53-dependent G1 arrest and long-term induction of Cip1 in normal human fibroblasts. *Genes. Dev.*, **8**:2540-2551.
- Di Micco, R., Fumagalli, M., Cicalese, A., Piccinin, S., Gasparini, P., Luise, C., Schurra, C., Garre, M., Nuciforo, P.G., Bensimon, A., et al., 2006. Oncogene-induced senescence is a DNA damage response triggered by DNA hyper-replication. *Nature*, **444**(7119):638-642. [doi:10.1038/nature05327]
- Diaz, J.L., Oltersdorf, T., Horne, W., McConnell, M., Wilson, G., Weeks, S., Garcia, T., Fritz, L.C., 1997. A common binding site mediates heterodimerization and homodimerization of Bcl-2 family members. *J. Biol. Chem.*, **272**(17):11350-11355. [doi:10.1074/jbc.272.17.11350]
- Domen, J., Cheshier, S.H., Weissman, I.L., 2000. The role of apoptosis in the regulation of hematopoietic stem cells. Overexpression of bcl-2 increases both their number and repopulation potential. *J. Exp. Med.*, **191**(2):253-264. [doi:10.1084/jem.191.2.253]
- Domina, A.M., Smith, J.H., Craig, R.W., 2000. Myeloid cell leukemia 1 is phosphorylated through two distinct pathways, one associated with extracellular signal-regulated kinase activation and the other with G2/M accumulation or protein phosphatase 1/2A inhibition. *J. Biol. Chem.*, **275**(28):21688-21694. [doi:10.1074/jbc.M000915200]
- Draetta, G., Beach, D., 1988. Activation of cdc2 protein kinase during mitosis in human cells: cell cycle-dependent phosphorylation and subunit rearrangement. *Cell*, **54**(1):17-26. [doi:10.1016/0092-8674(88)90175-4]
- Draetta, G., Luca, F., Westendorf, J., Brizuela, L., Ruderman, J., Beach, D., 1989. Cdc2 protein kinase is complexed with both cyclin A and B: evidence for proteolytic inactivation of MPF. *Cell*, **56**(5):829-838. [doi:10.1016/0092-8674(89)90687-9]
- Droin, N., Dubrez, L., Eymin, B., Renvoize, C., Breard, J., Dimanche-Boitrel, M.T., Solary, E., 1998. Upregulation of Casp genes in human tumor cells undergoing etoposide-induced apoptosis. *Oncogene*, **16**(22):2885-2894. [doi:10.1038/sj.onc.1201821]
- Du, C., Fang, M., Li, Y., Li, L., Wang, X., 2000. Smac, a mitochondrial protein that promotes cytochrome c-dependent caspase activation by eliminating IAP inhibition. *Cell*, **102**(1):33-42. [doi:10.1016/S0092-8674(00)00008-8]
- Dubrez, L., Goldwasser, F., Genne, P., Pommier, Y., Solary, E., 1995. The role of cell cycle regulation and apoptosis triggering in determining the sensitivity of leukemic cells to topoisomerase I and II inhibitors. *Leukemia*, **9**:1013-1024.
- Dubrez, L., Savoy, I., Hamman, A., Solary, E., 1996. Pivotal role of a DEVD-sensitive step in etoposide-induced and Fas-mediated apoptotic pathways. *EMBO J.*, **15**:5504-5512.
- Ducommun, B., Brambilla, P., Felix, M.A., Franza, B.R.Jr, Karsenti, E., Draetta, G., 1991. Cdc2 phosphorylation is required for its interaction with cyclin. *EMBO J.*, **10**:3311-3319.
- Dumont, P., Leu, J.I., Della Pietra, A.C., George, D.L., Murphy, M., 2003. The codon 72 polymorphic variants of p53 have markedly different apoptotic potential. *Nat. Genet.*, **33**(3):357-365. [doi:10.1038/ng1093]
- Epand, R.F., Martinou, J.C., Fornallaz-Mulhauser, M., Hughes, D.W., Epand, R.M., 2002. The apoptotic protein tBid promotes leakage by altering membrane curvature. *J. Biol. Chem.*, **277**(36):32632-32639. [doi:10.1074/jbc.M202396

- 200]
- Falck, J., Mailand, N., Syljuasen, R.G., Bartek, J., Lukas, J., 2001. The ATM-Chk2-Cdc25A checkpoint pathway guards against radioresistant DNA synthesis. *Nature*, **410**(6830):842-847. [doi:10.1038/35071124]
- Fan, M., Du, C., Stone, A.A., Gilbert, K.M., Chambers, T.C., 2000a. Modulation of mitogen-activated protein kinases and phosphorylation of Bcl-2 by vinblastine represent persistent forms of normal fluctuations at G2-M. *Cancer Res.*, **60**:6403-6407.
- Fan, M., Goodwin, M., Vu, T., Brantley-Finley, C., Gaarde, W.A., Chambers, T.C., 2000b. Vinblastine-induced phosphorylation of Bcl-2 and Bcl-XL is mediated by JNK and occurs in parallel with inactivation of the Raf-1/MEK/ERK cascade. *J. Biol. Chem.*, **275**(39):29980-29985. [doi:10.1074/jbc.M003776200]
- Fang, G.F., Chang, B.S., Kim, C.N., Perkins, C., Thompson, C.B., Bhalla, K.N., 1998. Loop domain is necessary for taxol-induced mobility shift and phosphorylation of Bcl-2 as well as for inhibiting taxol-induced cytosolic accumulation of cytochrome c and apoptosis. *Cancer Res.*, **58**:3202-3208.
- Farrow, S.N., White, J.H., Martinou, I., Raven, T., Pun, K.T., Grinham, C.J., Martinou, J.C., Brown, R., 1995. Cloning of a bcl-2 homologue by interaction with adenovirus E1B 19K. *Nature*, **374**(6524):731-733. [doi:10.1038/374731a0]
- Flatten, K., Dai, N.T., Vroman, B.T., Loegering, D., Erlichman, C., Karnitz, L.M., Kaufmann, S.H., 2005. The role of checkpoint kinase 1 in sensitivity to topoisomerase I poisons. *J. Biol. Chem.*, **280**(14):14349-14355. [doi:10.1074/jbc.M411890200]
- Flores, E.R., Tsai, K.Y., Crowley, D., Sengupta, S., Yang, A., McKeon, F., Jacks, T., 2002. p63 and p73 are required for p53-dependent apoptosis in response to DNA damage. *Nature*, **416**(6880):560-564. [doi:10.1038/416560a]
- Fontemaggi, G., Kela, I., Amariglio, N., Rechavi, G., Krishnamurthy, J., Strano, S., Sacchi, A., Givol, D., Blandino, G., 2002. Identification of direct p73 target genes combining DNA microarray and chromatin immunoprecipitation analyses. *J. Biol. Chem.*, **277**(45):43359-43368. [doi:10.1074/jbc.M205573200]
- Foo, S.Y., Nolan, G.P., 1999. NF-kappaB to the rescue: RELs, apoptosis and cellular transformation. *Trends Genet.*, **15**(6):229-235. [doi:10.1016/S0168-9525(99)01719-9]
- Freebern, W.J., Smith, J.L., Chaudhry, S.S., Haggerty, C.M., Gardner, K., 2003. Novel cell-specific and dominant negative anti-apoptotic roles of p73 in transformed leukemia cells. *J. Biol. Chem.*, **278**(4):2249-2255. [doi:10.1074/jbc.M208517200]
- Fujise, K., Zhang, D., Liu, J., Yeh, E.T., 2000. Regulation of apoptosis and cell cycle progression by MCL1. Differential role of proliferating cell nuclear antigen. *J. Biol. Chem.*, **275**(50):39458-39465. [doi:10.1074/jbc.M006626200]
- Furnari, B., Rhind, N., Russell, P., 1997. Cdc25 mitotic inducer targeted by Chk1 DNA damage checkpoint kinase. *Science*, **277**(5331):1495-1497. [doi:10.1126/science.277.5331.1495]
- Furnari, B., Blasina, A., Boddy, M.N., McGowan, C.H., Russell, P., 1999. Cdc25 inhibited in vivo and in vitro by checkpoint kinases Cds1 and Chk1. *Mol. Biol. Cell*, **10**:833-845.
- Gibson, L., Holmgreen, S.P., Huang, D.C.S., Bernard, O., Copeland, N.G., Jenkins, N.A., Sutherland, G.R., Baker, E., Adams, J.M., Cory, S., 1996. Bcl-W, a novel member of the Bcl-2 family, promotes cell survival. *Oncogene*, **13**:665-675.
- Gillard, F., Moll, I., Meyer, M., Michaelidis, T.M., 1999. Alterations in cell death and cell cycle progression in the UV-irradiated epidermis of bcl-2-deficient mice. *Cell Death Differ.*, **6**(1):55-60. [doi:10.1038/sj.cdd.4400455]
- Gong, J.G., Costanzo, A., Yang, H.Q., Melino, G., Kaelin, W.G.Jr, Levvero, M., Wang, J.Y., 1999. The tyrosine kinase c-Abl regulates p73 in apoptotic response to cis-platin-induced DNA damage. *Nature*, **399**(6738):806-809. [doi:10.1038/21690]
- Gottlieb, T.M., Oren, M., 1998. p53 and apoptosis. *Semin. Cancer Biol.*, **8**(5):359-368. [doi:10.1006/scbi.1998.0098]
- Gottlieb, R.A., Nordberg, J., Skowronski, E., Babior, B.M., 1996. Apoptosis induced in Jurkat cells by several agents is preceded by intracellular acidification. *Proc. Natl. Acad. Sci. (USA)*, **93**(2):654-658. [doi:10.1073/pnas.93.2.654]
- Gross, A., Jockel, J., Wei, M.C., Korsmeyer, S.J., 1998. Enforced dimerization of Bax results in its translocation, mitochondrial dysfunction and apoptosis. *EMBO J.*, **17**(14):3878-3885. [doi:10.1093/emboj/17.14.3878]
- Grumont, R.J., Rourke, I.J., O'Reilly, L.A., Strasser, A., Miyake, K., Sha, W., Gerondakis, S., 1998. B lymphocytes differentially use the Rel and nuclear factor kappa-B1 (NF-kappa-B1) transcription factors to regulate cell cycle progression and apoptosis in quiescent and mitogen-activated cells. *J. Exp. Med.*, **187**(5):663-674. [doi:10.1084/jem.187.5.663]
- Guney, I., Sedivy, J.M., 2006. Cellular senescence, epigenetic switches and c-Myc. *Cell Cycle*, **5**:2319-2323.
- Guo, B., Godzik, A., Reed, J.C., 2001. Bcl-G, a novel pro-apoptotic member of the bcl-2 family. *J. Biol. Chem.*, **276**(4):2780-2785. [doi:10.1074/jbc.M005889200]
- Gupta, M., Fan, S.J., Zhan, Q.M., Kohn, K.W., O'Connor, P.M., Pommier, Y., 1997. Inactivation of p53 increases the cytotoxicity of camptothecin in human colon HCT116 and breast MCF-7 cancer cells. *Clin. Cancer Res.*, **3**:1653-1660.
- Haldar, S., Jena, N., Croce, C.M., 1995. Inactivation of Bcl-2 by phosphorylation. *Proc. Natl. Acad. Sci. (USA)*, **92**(10):4507-4511. [doi:10.1073/pnas.92.10.4507]
- Haldar, S., Chintapalli, J., Croce, C.M., 1996. Taxol induces Bcl-2 phosphorylation and death of prostate cancer cells. *Cancer Res.*, **56**:1253-1255.
- Haldar, S., Basu, A., Croce, C.M., 1998. Serine-70 is one of the critical sites for drug-induced Bcl2 phosphorylation in cancer cells. *Cancer Res.*, **58**:1609-1615.
- Han, J., Sabbatini, P., White, E., 1996. Induction of apoptosis by human Nbk/Bik, a BH3-containing protein that interacts with E1b 19k. *Mol. Cell Biol.*, **16**:5857-5864.

- Han, J., Flemington, C., Houghton, A.B., Gu, Z., Zambetti, G.P., Lutz, R.J., Zhu, L., Chittenden, T., 2001. Expression of Bbc3, a pro-apoptotic BH3-only gene, is regulated by diverse cell death and survival signals. *Proc. Natl. Acad. Sci. (USA)*, **98**(20):11318-11323. [doi:10.1073/pnas.201208798]
- Han, Z., Wei, W., Dunaway, S., Darnowski, J.W., Calabresi, P., Sedivy, J., Hendrickson, E.A., Balan, K.V., Pantazis, P., Wyche, J.H., 2002. Role of p21 in apoptosis and senescence of human colon cancer cells treated with camptothecin. *J. Biol. Chem.*, **277**(19):17154-17160. [doi:10.1074/jbc.M112401200]
- Harris, M.H., Thompson, C.B., 2000. The role of the Bcl-2 family in the regulation of outer mitochondrial membrane permeability. *Cell Death Differ.*, **7**(12):1182-1191. [doi:10.1038/sj.cdd.4400781]
- Hegde, R., Srinivasula, S.M., Ahmad, M., Fernandes-Alnemri, T., Alnemri, E.S., 1998. Blk, a BH3-containing mouse protein that interacts with Bcl-2 and Bcl-xL, is a potent death agonist. *J. Biol. Chem.*, **273**(14):7783-7786. [doi:10.1074/jbc.273.14.7783]
- Hemann, M.T., Narita, M., 2007. Oncogenes and senescence: breaking down in the fast line. *Genes Dev.*, **21**(1):1-5. [doi:10.1101/gad.1514207]
- Herbig, U., Jobling, W.A., Chen, B.P., Chen, D.J., Sedivy, J.M., 2004. Telomere shortening triggers senescence of human cells through a pathway involving ATM, p53, and p21(Cip1), but not p16(INK4a). *Mol. Cell*, **14**(4):501-513. [doi:10.1016/S1097-2765(04)00256-4]
- Hermeking, H., Lengauer, C., Polyak, K., He, T., Zhang, L., Thiagalingam, S., 1997. 14-3-3 sigma is a p53-regulated inhibitor of G2/M progression. *Mol. Cell*, **1**(1):3-11. [doi:10.1016/S1097-2765(00)80002-7]
- Hershko, T., Ginsberg, D., 2004. Up-regulation of Bcl-2 homology 3 (BH3)-only proteins by E2F1 mediates apoptosis. *J. Biol. Chem.*, **279**(10):8627-8634. [doi:10.1074/jbc.M312866200]
- Hirota, T., Kunitoku, N., Sasayama, T., Marumoto, T., Zhang, D., Nitta, M., Hatakeyama, K., Saya, H., 2003. Aurora-A and an interacting activator, the LIM protein Ajuba, are required for mitotic commitment in human cells. *Cell*, **114**(5):585-598. [doi:10.1016/S0092-8674(03)00642-1]
- Hockenbery, D., Nunez, G., Millman, C., Schreiber, R.D., Korsmeyer, S.J., 1990. Bcl-2 is an inner mitochondrial membrane protein that blocks programmed cell death. *Nature*, **348**(6299):334-336. [doi:10.1038/348334a0]
- Hockenbery, D.M., Oltvai, Z.N., Yin, X.M., Millman, C.L., Korsmeyer, S.J., 1993. Bcl-2 functions in an antioxidant pathway to prevent apoptosis. *Cell*, **75**(2):241-251. [doi:10.1016/0092-8674(93)80066-N]
- Hofmann, K., Bucher, P., Kajava, A.V., 1998. A model of Cdc25 phosphatase catalytic domain and Cdk-interaction surface based on the presence of a rhodanese homology domain. *J. Mol. Biol.*, **282**(1):195-208. [doi:10.1006/jmbi.1998.1998]
- Holliday, R., Tarrant, G.M., 1972. Altered enzymes in ageing human fibroblasts. *Nature*, **238**(5358):26-30. [doi:10.1038/238026a0]
- Holmberg, C., Helin, K., Sehested, M., Karlstrom, O., 1998. E2F-1-induced p53-independent apoptosis in transgenic mice. *Oncogene*, **17**(2):143-155. [doi:10.1038/sj.onc.1201915]
- Hsu, S.Y., Kaipia, A., McGee, E., Lomeli, M., Hsueh, A.J.W., 1997a. Bok is a pro-apoptotic Bcl-2 protein with restricted expression in reproductive tissues and heterodimerizes with selective anti-apoptotic Bcl-2 family members. *Proc. Natl. Acad. Sci. (USA)*, **94**(23):12401-12406. [doi:10.1073/pnas.94.23.12401]
- Hsu, Y.T., Wolter, K.G., Youle, R.J., 1997b. Cytosol-to-membrane redistribution of Bax and Bcl-X-L during apoptosis. *Proc. Natl. Acad. Sci. (USA)*, **94**(8):3668-3672. [doi:10.1073/pnas.94.8.3668]
- Hsu, S.Y., Lin, P., Hsueh, A.J.W., 1998. Bod (Bcl-2-related ovarian death gene) is an ovarian BH3 domain-containing proapoptotic Bcl-2 protein capable of dimerization with diverse antiapoptotic Bcl-2 members. *Mol. Endocrinol.*, **12**(9):1432-1440. [doi:10.1210/me.12.9.1432]
- Huang, D.C.S., O'Reilly, L.A., Strasser, A., Cory, S., 1997. The anti-apoptosis function of Bcl-2 can be genetically separated from its inhibitory effect on cell cycle entry. *EMBO J.*, **16**(15):4628-4638. [doi:10.1093/emboj/16.15.4628]
- Hunt, K.K., Deng, J., Liu, T.J., Wilson-Heiner, M., Swisher, S.G., Clayman, G., Hung, M.C., 1997. Adenovirus-mediated overexpression of the transcription factor E2F-1 induces apoptosis in human breast and ovarian carcinoma cell lines and does not require p53. *Cancer Res.*, **57**:4722-4726.
- Hunter, T., Pines, J., 1994. Cyclins and cancer. 2: Cyclin D and CDK inhibitors come of age. *Cell*, **79**(4):573-582. [doi:10.1016/0092-8674(94)90543-6]
- Imaizumi, K., Morihara, T., Mori, Y., Katayama, T., Tsuda, M., Furuyama, T., Wanaka, A., Takeda, M., Tohyama, M., 1999. The cell death-promoting gene DP5, which interacts with the BCL2 family, is induced during neuronal apoptosis following exposure to amyloid beta protein. *J. Biol. Chem.*, **274**(12):7975-7981. [doi:10.1074/jbc.274.12.7975]
- Innocente, S.A., Abrahamson, J.L., Cogswell, J.P., Lee, J.M., 1999. p53 regulates a G2 checkpoint through cyclin B1. *Proc. Natl. Acad. Sci. (USA)*, **96**(5):2147-2152. [doi:10.1073/pnas.96.5.2147]
- Inohara, N., Ding, L.Y., Chen, S., Nunez, G., 1997. Harakiri, a novel regulator of cell death, encodes a protein that activates apoptosis and interacts selectively with survival-promoting proteins Bcl-2 and Bcl-X-L. *EMBO J.*, **16**(7):1686-1694. [doi:10.1093/emboj/16.7.1686]
- Inohara, N., Gourley, T.S., Carrio, R., Muniz, M., Merino, J., Garcia, I., Koseki, T., Hu, Y., Chen, S., Nunez, G., 1998a. Diva, a Bcl-2 homologue that binds directly to Apaf-1 and induces BH3-independent cell death. *J. Biol. Chem.*, **273**(49):32479-32486. [doi:10.1074/jbc.273.49.32479]
- Inohara, N., Ekhterae, D., Garcia, I., Carrio, R., Merino, J., Merry, A., Chen, S., Nunez, G., 1998b. Mtd, a novel Bcl-2 family member activates apoptosis in the absence of het-

- erodimerization with Bcl-2 and Bcl-X-L. *J. Biol. Chem.*, **273**(15):8705-8710. [doi:10.1074/jbc.273.15.8705]
- Jacks, T., 1996. Tumor suppressor gene mutations in mice. *Ann. Rev. Genet.*, **30**(1):603-636. [doi:10.1146/annurev.genet.30.1.603]
- Jackson, J.P., Lindroth, A.M., Cao, X., Jacobsen, S.E., 2002. Control of CpNpG DNA methylation by the KRYPTONITE histone H3 methyltransferase. *Nature*, **416**(6880):556-560. [doi:10.1038/nature731]
- Jamil, S., Sobouti, R., Hojabrpour, P., Raj, M., Kast, J., Duronio, V., 2005. A proteolytic fragment of Mcl-1 exhibits nuclear localization and regulates cell growth via interaction with Cdk1. *Biochem. J.*, **387**(3):659-667. [doi:10.1042/BJ20041596]
- Johnstone, R.W., Ruefli, A.A., Lowe, S.W., 2002. Apoptosis. A link between cancer genetics and chemotherapy. *Cell*, **108**(2):153-164. [doi:10.1016/S0092-8674(02)00625-6]
- Jost, C.A., Marin, M.C., Kaelin, W.G., 1997. p73 is a human p53-related protein that can induce apoptosis. *Nature*, **389**(6647):191-194. [doi:10.1038/38298]
- Kamer, I., Sarig, R., Zaltsman, Y., Niv, H., Oberkovitz, G., Regev, L., Haimovich, G., Lerenthal, Y., Marcellus, R., Gross, A., 2005. Proapoptotic Bid is an ATM effector in the DNA-damage response. *Cell*, **122**(4):593-603. [doi:10.1016/j.cell.2005.06.014]
- Kane, D.J., Sarafian, T.A., Anton, R., Hahn, H., Gralla, E.B., Valentine, J.S., Ord, T., Bredesen, D.E., 1993. Bcl-2 inhibition of neural death: decreased generation of reactive oxygen species. *Science*, **262**(5137):1274-1277. [doi:10.1126/science.8235659]
- Kang, Y.K., Koo, D.B., Park, J.S., Choi, Y.H., Lee, K.K., Han, Y.M., 2001. Differential inheritance modes of DNA methylation between euchromatic and heterochromatic DNA sequences in ageing fetal bovine fibroblasts. *FEBS Lett.*, **498**(1):1-5. [doi:10.1016/S0014-5793(01)02472-3]
- Kastan, M.B., Canman, C.E., Leonard, C.J., 1995. p53, cell cycle control and apoptosis: implications for cancer. *Cancer Metast. Rev.*, **14**(1):3-15. [doi:10.1007/BF00690207]
- Kataoka, T., Holler, N., Micheau, O., Martinon, F., Tinel, A., Hofmann, K., Tschopp, J., 2001. Bcl-rambo, a novel Bcl-2 homologue that induces apoptosis via its unique C-terminal extension. *J. Biol. Chem.*, **276**(22):19548-19554. [doi:10.1074/jbc.M010520200]
- Kaufmann, S.H., 1989. Induction of endonucleolytic DNA cleavage in human acute myelogenous leukemia cells by etoposide, camptothecin, and other cytotoxic anticancer drugs: a cautionary note. *Cancer Res.*, **49**:5870-5878.
- Kaufmann, S.H., 1998. Cell death induced by topoisomerase-targeted drugs: more questions than answers. *Biochim. Biophys. Acta*, **1400**:195-211.
- Kaufmann, W.K., Paules, R.S., 1996. DNA damage and cell cycle checkpoints. *FASEB J.*, **10**:238-247.
- Kaufmann, S.H., Earnshaw, W.C., 2000. Induction of apoptosis by cancer chemotherapy. *Exp. Cell Res.*, **256**(1):42-49. [doi:10.1006/excr.2000.4838]
- Ke, N., Godzik, A., Reed, J.C., 2001. Bcl-B, a novel Bcl-2 family member that differentially binds and regulates Bax and Bak. *J. Biol. Chem.*, **276**(16):12481-12484. [doi:10.1074/jbc.C000871200]
- Kelekar, A., Thompson, C.B., 1998. Bcl-2-family proteins: the role of the BH3 domain in apoptosis. *Trends Cell Biol.*, **8**(8):324-330. [doi:10.1016/S0962-8924(98)01321-X]
- Kharbanda, S., Saxena, S., Yoshida, K., Pandey, P., Kaneki, M., Wang, Q., Cheng, K., Chen, Y.N., Campbell, A., Sudha, T., et al., 2000. Translocation of SAPK/JNK to mitochondria and interaction with Bcl-x(L) in response to DNA damage. *J. Biol. Chem.*, **275**(1):322-327. [doi:10.1074/jbc.275.1.322]
- Kiefer, M.C., Brauer, M.J., Powers, V.C., Wu, J.J., Umansky, S.R., Tomei, L.D., Barr, P.J., 1995. Modulation of apoptosis by the widely distributed Bcl-2 homologue Bak. *Nature*, **374**(6524):736-739. [doi:10.1038/374736a0]
- Kim, H., Rafiuddin-Shah, M., Tu, H.C., Jeffers, J.R., Zambetti, G.P., Hsieh, J.J.D., Cheng, E.H.Y., 2006. Hierarchical regulation of mitochondrion-dependent apoptosis by BCL-2 subfamilies. *Nat. Cell Biol.*, **8**(12):1348-1358. [doi:10.1038/ncb1499]
- King, R.W., Jackson, P.K., Kirschner, M.W., 1994. Mitosis in transition. *Cell*, **79**(4):563-571. [doi:10.1016/0092-8674(94)90542-8]
- Kluck, R.M., Bossy-Wetzel, E., Green, D.R., Newmeyer, D.D., 1997. The release of cytochrome C from mitochondria: a primary site for Bcl-2 regulation of apoptosis. *Science*, **275**(5303):1132-1136. [doi:10.1126/science.275.5303.1132]
- Knowlton, K., Mancini, M., Creason, S., Morales, C., Hockenbery, D., Anderson, B.O., 1998. Bcl-2 slows in vitro breast cancer growth despite its antiapoptotic effect. *J. Surg. Res.*, **76**(1):22-26. [doi:10.1006/jsre.1998.5277]
- Koff, A., Giordano, A., Desai, D., Yamashita, K., Harper, J.W., Elledge, S., Nishimoto, T., Morgan, D.O., Franza, B.R., Roberts, J.M., 1992. Formation and activation of a cyclinE-cdk2 complex during the G1 phase of the human cell cycle. *Science*, **257**(5077):1689-1694. [doi:10.1126/science.1388288]
- Kohn, K.W., 1996. Regulatory genes and drug sensitivity. *J. Natl. Cancer Inst.*, **88**(18):1255-1256. [doi:10.1093/jnci/88.18.1255]
- Kohn, K.W., Jackman, J., O'Connor, P.M., 1994. Cell cycle control and cancer chemotherapy. *J. Cell. Biochem.*, **54**(4):440-452. [doi:10.1002/jcb.240540411]
- Korsmeyer, S.J., 1992. Bcl-2 initiates a new category of oncogenes: regulators of cell death. *Blood*, **80**:879-886.
- Korsmeyer, S.J., Wei, M.C., Saito, M., Weiler, S., Oh, K.J., Schlesinger, P.H., 2000. Pro-apoptotic cascade activates Bid, which oligomerizes Bak or Bax into pores that result in the release of cytochrome c. *Cell Death Differ.*, **7**(12):1166-1173. [doi:10.1038/sj.cdd.4400783]
- Kozopas, K.M., Yang, T., Buchan, H.L., Zhou, P., Craig, R.W., 1993. Mcl1, a gene expressed in programmed myeloid cell differentiation, has sequence similarity to Bcl-2. *Proc. Natl. Acad. Sci. (USA)*, **90**(8):3516-3520. [doi:10.1073/pnas.90.8.3516]

- Krajewski, S., Tanaka, S., Takayama, S., Schibler, M.J., Fenton, W., Reed, J.C., 1993. Investigation of the subcellular distribution of the bcl-2 oncoprotein: residence in the nuclear envelope, endoplasmic reticulum, and outer mitochondrial membranes. *Cancer Res.*, **53**:4701-4714.
- Krajewski, S., Krajewska, M., Ellerby, L.M., Welsh, K., Xie, Z.H., Deveraux, Q.L., Salvesen, G.S., Bredesen, D.E., Rosenthal, R.E., Fiskum, G., Reed, J.C., 1999. Release of caspase-9 from mitochondria during neuronal apoptosis and cerebral ischemia. *Proc. Natl. Acad. Sci. (USA)*, **96**(10):5752-5757. [doi:10.1073/pnas.96.10.5752]
- Kudla, G., Montessuit, S., Eskes, R., Berrier, C., Martinou, J.C., Ghazi, A., Antonsson, B., 2000. The destabilization of lipid membranes induced by the C-terminal fragment of caspase 8-cleaved bid is inhibited by the N-terminal fragment. *J. Biol. Chem.*, **275**(30):22713-22718. [doi:10.1074/jbc.M003807200]
- Kung, A.L., Zetterberg, A., Sherwood, S.W., Schimke, R.T., 1990. Cytotoxic effects of cell cycle phase specific agents: result of cell cycle perturbation. *Cancer Res.*, **50**:7307-7317.
- Kuwana, T., Mackey, M.R., Perkins, G., Ellisman, M., Latterich, M., Schneider, R., Green, D., Newmeyer, D., 2002. Bid, bax, and lipids cooperate to form supramolecular openings in the outer mitochondrial membrane. *Cell*, **111**(3):331-342. [doi:10.1016/S0092-8674(02)01036-X]
- Kuwana, T., Bouchier-Hayes, L., Chipuk, J.E., Bonzon, C., Sullivan, B.A., Green, D.R., Newmeyer, D.D., 2005. BH3 domains of BH3-only proteins differentially regulate Bax-mediated mitochondrial membrane permeabilization both directly and indirectly. *Mol. Cell*, **17**(4):525-535. [doi:10.1016/j.molcel.2005.02.003]
- Labib, K., Craven, R.A., Crawford, K., Nurse, P., 1995. Dominant mutants identify new roles for p34cdc2 in mitosis. *EMBO J.*, **14**:2155-2165.
- Lammer, C., Wagerer, S., Saffrich, R., Mertens, D., Ansorge, W., Hoffmann, I., 1998. The cdc25B phosphatase is essential for the G2/M phase transition in human cells. *J. Cell Sci.*, **111**:2445-2453.
- Lee, S., Schmitt, C.A., 2003. Chemotherapy response and resistance. *Curr. Opin. Genet. Dev.*, **13**(1):90-96. [doi:10.1016/S0959-437X(02)00014-X]
- Letai, A., Bassik, M., Walensky, L., Sorcinelli, M., Weiler, S., Korsmeyer, S., 2002. Distinct BH3 domains either sensitize or activate mitochondrial apoptosis, serving as prototype cancer therapeutics. *Cancer Cell*, **2**(3):183-192. [doi:10.1016/S1535-6108(02)00127-7]
- Leu, J.I.J., Dumont, P., Hafey, M., Murphy, M.E., George, D.L., 2004. Mitochondrial p53 activates Bak and causes disruption of a Bak-Mcl1 complex. *Nat. Cell Biol.*, **6**(5):443-450. [doi:10.1038/ncb1123]
- Levine, A.J., 1997. p53, the cellular gatekeeper for growth and division. *Cell*, **88**(3):323-331. [doi:10.1016/S0092-8674(00)81871-1]
- Li, L.Y., Luo, X., Wang, X., 2001. Endonuclease G is an apoptotic DNase when released from mitochondria. *Nature*, **412**(6842):95-99. [doi:10.1038/35083620]
- Lin, E.Y., Orlofsky, A., Berger, M.S., Prystowsky, M.B., 1993. Characterization of A1, a novel hemopoietic-specific early-response gene with sequence similarity to bcl-2. *J. Immunol.*, **151**:1979-1988.
- Linette, G.P., Li, Y., Roth, K., Korsmeyer, S.J., 1996. Cross talk between cell death and cell cycle progression: BCL-2 regulates NFAT-mediated activation. *Proc. Natl. Acad. Sci. (USA)*, **93**(18):9545-9552. [doi:10.1073/pnas.93.18.9545]
- Ling, Y.H., Tornos, C., Perez-Soler, R., 1998. Phosphorylation of Bcl-2 is a marker of M phase events and not a determinant of apoptosis. *J. Biol. Chem.*, **273**(30):18984-18991. [doi:10.1074/jbc.273.30.18984]
- Linke, S.P., Clarkin, K.C., Wahl, G.M., 1997. p53 mediates permanent arrest over multiple cell cycles in response to gamma-irradiation. *Cancer Res.*, **57**:1171-1179.
- Liu, F., Stanton, J.J., Wu, Z., Piwnicka-Worms, H., 1997. The human Myt1 kinase preferentially phosphorylates Cdc2 on threonine 14 and localizes to the endoplasmic reticulum and Golgi complex. *Mol. Cell Biol.*, **17**:571-583.
- Liu, Q., Guntuku, S., Cui, X.S., Matsuoka, S., Cortez, D., Tamai, K., Luo, G., Carattini-Rivera, S., DeMayo, F., Bradley, A., Donehower, L.A., Elledge, S.J., 2000. Chk1 is an essential kinase that is regulated by Atr and required for the G(2)/M DNA damage checkpoint. *Genes Dev.*, **14**(12):1448-1459.
- Liu, Z., Lu, H., Jiang, Z.H., Pastuszyn, A., Hu, C.A., 2005. Apolipoprotein L6, a novel proapoptotic Bcl-2 homology 3-only protein, induces mitochondria-mediated apoptosis in cancer cells. *Mol. Cancer Res.*, **3**:21-31.
- Lopatina, N., Haskell, J.F., Andrews, L.G., Poole, J.C., Saldanha, S., Tollefsbol, T., 2002. Differential maintenance and de novo methylating activity by three DNA methyltransferases in aging and immortalized fibroblasts. *J. Cell. Biochem.*, **84**(2):324-334. [doi:10.1002/jcb.10015]
- Lowe, S.W., Ruley, H.E., Jacks, T., Housman, D.E., 1993. p53-dependent apoptosis modulates the cytotoxicity of anticancer agents. *Cell*, **74**(6):957-967. [doi:10.1016/0092-8674(93)90719-7]
- Lowe, S.W., Bodis, S., McClatchey, A., Remington, L., Ruley, H.E., Fisher, D.E., Housman, D.E., Jacks, T., 1994. p53 status and the efficacy of cancer therapy in vivo. *Science*, **266**(5186):807-810. [doi:10.1126/science.7973635]
- Lukas, J., Lukas, C., Bartek, J., 2004. Mammalian cell cycle checkpoints: signalling pathways and their organization in space and time. *DNA Repair (Amst.)*, **3**(8-9):997-1007. [doi:10.1016/j.dnarep.2004.03.006]
- Machwe, A., Orren, D.K., Bohr, V.A., 2000. Accelerated methylation of ribosomal RNA genes during the cellular senescence of Werner syndrome fibroblasts. *FASEB J.*, **14**(12):1715-1724. [doi:10.1096/fj.99-0926com]
- Mallette, F.A., Gaumont-Leclerc, M.F., Ferbeyre, G., 2007. The DNA damage signaling pathway is a critical mediator of oncogene-induced senescence. *Genes Dev.*, **21**(1):43-48. [doi:10.1101/gad.1487307]
- Mancini, M., Nicholson, D.W., Roy, S., Thornberry, N.A., Peterson, E.P., Casciola-Rosen, L.A., Rosen, A., 1998.

- The caspase-3 precursor has a cytosolic and mitochondrial distribution: implications for apoptotic signaling. *J. Cell Biol.*, **140**(6):1485-1495. [doi:10.1083/jcb.140.6.1485]
- Marumoto, T., Hirota, T., Morisaki, T., Kunitoku, N., Zhang, D., Ichikawa, Y., Sasayama, T., Kuninaka, S., Mimori, T., Tamaki, N., Kimura, M., Okano, Y., Saya, H., 2002. Roles of aurora-A kinase in mitotic entry and G2 checkpoint in mammalian cells. *Genes Cells*, **7**(11):1173-1182. [doi:10.1046/j.1365-2443.2002.00592.x]
- Marzo, I., Brenner, C., Zamzami, N., Jurgensmeier, J.M., Susin, S.A., Vieira, H.L., Prevost, M.C., Xie, Z., Matsuyama, S., Reed, J.C., Kroemer, G., 1998. Bax and adenine nucleotide translocator cooperate in the mitochondrial control of apoptosis. *Science*, **281**(5385):2027-2031. [doi:10.1126/science.281.5385.2027]
- Matsuoka, S., Huang, M., Elledge, S.J., 1998. Linkage of ATM to cell cycle regulation by the Chk2 protein kinase. *Science*, **282**(5395):1893-1897. [doi:10.1126/science.282.5395.1893]
- Matsushima, M., Fujiwara, T., Takahashi, E., Minaguchi, T., Eguchi, Y., Tsujimoto, Y., Suzumori, K., Nakamura, Y., 1998. Isolation, mapping, and functional analysis of a novel human cDNA (*BNIP3L*) encoding a protein homologous to human NIP3. *Genes. Chromos. Cancer*, **21**(3):230-235. [doi:10.1002/(SICI)1098-2264(199803)21:3<230::AID-GCC7>3.3.CO;2-G]
- Maundrell, K., Antonsson, B., Magnenat, E., Camps, M., Muda, M., Chabert, C., Gillieron, C., Boschert, U., Vial-Knecht, E., Martinou, J.C., Arkinstall, S., 1997. Bcl-2 undergoes phosphorylation by C-Jun N-terminal kinase stress-activated protein kinases in the presence of the constitutively active GTP-binding protein Rac1. *J. Biol. Chem.*, **272**(40):25238-25242. [doi:10.1074/jbc.272.40.25238]
- Mazel, S., Burtrum, D., Petrie, H.T., 1996. Regulation of cell division cycle progression by bcl-2 expression: a potential mechanism for inhibition of programmed cell death. *J. Exp. Med.*, **183**(5):2219-2226. [doi:10.1084/jem.183.5.2219]
- McDonnell, T.J., Marin, M.C., Hsu, B., Brisbay, S.M., McConnell, K., Tu, S.M., Campbell, M.L., Rodriguez-Villanueva, J., 1993. The Bcl-2 oncogene: apoptosis and neoplasia. *Radiation Res.*, **136**(3):307-312. [doi:10.2307/3578541]
- McDonnell, J.M., Fushman, D., Millman, C.L., Korsmeyer, S.J., Cowburn, D., 1999. Solution structure of the proapoptotic molecule BID: a structural basis for apoptotic agonists and antagonists. *Cell*, **96**(5):625-634. [doi:10.1016/S0092-8674(00)80573-5]
- McGowan, C.H., Russell, P., 1995. Cell cycle regulation of human Wee1. *EMBO J.*, **14**:2166-2175.
- Melino, G., Bernassola, F., Ranalli, M., Yee, K., Zong, W.X., Corazzari, M., Knight, R.A., Green, D.R., Thompson, C., Vousden, K.H., 2004. p73 induces apoptosis via PUMA transactivation and Bax mitochondrial translocation. *J. Biol. Chem.*, **279**(9):8076-8083. [doi:10.1074/jbc.M307469200]
- Mercurio, F., Manning, A.M., 1999. NF- κ B as a primary regulator of the stress response. *Oncogene*, **18**(45):6163-6171. [doi:10.1038/sj.onc.1203174]
- Mihara, M., Erster, S., Zaika, A., Petrenko, O., Chittenden, T., Pancoska, P., Moll, U.M., 2003. p53 has a direct apoptogenic role at the mitochondria. *Mol. Cell*, **11**(3):577-590. [doi:10.1016/S1097-2765(03)00050-9]
- Minn, A.J., Velez, P., Schendel, S.L., Liang, H., Muchmore, S.W., Fesik, S.W., Fill, M., Thompson, C.B., 1997. Bcl-xL forms an ion channel in synthetic lipid membranes. *Nature*, **385**(6614):353-357. [doi:10.1038/385353a0]
- Miyashita, T., Reed, J.C., 1995. Tumor suppressor p53 is a direct transcriptional activator of the human bax gene. *Cell*, **80**(2):293-299. [doi:10.1016/0092-8674(95)90412-3]
- Miyashita, T., Krajewski, S., Krajewska, M., Wang, H.G., Lin, H.K., Liebermann, D.A., Hoffman, B., Reed, J.C., 1994. Tumor suppressor p53 is a regulator of bcl-2 and bax gene expression in vitro and in vivo. *Oncogene*, **9**:1799-1805.
- Muchmore, S.W., Sattler, M., Liang, H., Meadows, R.P., Harlan, J.E., Yoon, H.S., Nettesheim, D., Chang, B.S., Thompson, C.B., Wong, S.L., Ng, S.C., Fesik, S.W., 1996. X-ray and NMR structure of human Bcl-xL, an inhibitor of programmed cell death. *Nature*, **381**(6580):335-341. [doi:10.1038/381335a0]
- Mueller, P.R., Coleman, T.R., Kumagai, A., Dunphy, W.G., 1995. Myt1: a membrane-associated inhibitory kinase that phosphorylates Cdc2 on both threonine-14 and tyrosine-15. *Science*, **270**(5233):86-90. [doi:10.1126/science.270.5233.86]
- Muller, H., Bracken, A.P., Vernell, R., Moroni, M.C., Christians, F., Grassilli, E., Prosperini, E., Vigo, E., Oliner, J.D., Helin, K., 2001. E2Fs regulate the expression of genes involved in differentiation, development, proliferation, and apoptosis. *Genes Dev.*, **15**(3):267-285. [doi:10.1101/gad.864201]
- Nakano, K., Vousden, K.H., 2001. Puma, a novel proapoptotic gene, is induced by p53. *Mol. Cell*, **7**(3):683-694. [doi:10.1016/S1097-2765(01)00214-3]
- Neumeister, P., Albanese, C., Balent, B., Grealley, J., Pestell, R.G., 2002. Senescence and epigenetic dysregulation in cancer. *Int. J. Biochem. Cell Biol.*, **34**(11):1475-1490. [doi:10.1016/S1357-2725(02)00079-1]
- Nguyen, M., Branton, P.E., Walton, P.A., Oltvai, Z.N., Korsmeyer, S.J., Shore, G.C., 1994. Role of membrane anchor domain of Bcl-2 in suppression of apoptosis caused by E1B-defective adenovirus. *J. Biol. Chem.*, **269**:16521-16524.
- Nicotera, P., Leist, M., Ferrando-May, E., 1999. Apoptosis and necrosis: different execution of the same death. *Biochem. Soc. Symp.*, **66**:69-73.
- Nigg, E.A., 2001. Mitotic kinases as regulators of cell division and its checkpoints. *Nat. Rev. Mol. Cell Biol.*, **2**(1):21-32. [doi:10.1038/35048096]
- Nozell, S., Wu, Y., McNaughton, K., Liu, G., Willis, A., Paik, J.C., Chen, X., 2003. Characterization of p73 functional domains necessary for transactivation and growth suppression. *Oncogene*, **22**(28):4333-4347. [doi:10.1038/

- sj.onc.1206470]
- O'Connell, M.J., Raleigh, J.M., Verkade, H.M., Nurse, P., 1997. Chk 1 is a wee 1 kinase in the G2 DNA damage checkpoint inhibiting cdc2 by Y15 phosphorylation. *EMBO J.*, **16**(3):545-554. [doi:10.1093/emboj/16.3.545]
- O'Connor, P.M., 1997. Mammalian G1 and G2 phase checkpoints. *Cancer Surv.*, **29**:151-182.
- O'Connor, P.M., Jackman, J., Bae, I., Myers, T.G., Fan, S., Mutoh, M., Scudiero, D.A., Monks, A., Sausville, E.A., Weinstein, J.N., et al., 1997. Characterization of the p53 tumor suppressor pathway in cell lines of the National Cancer Institute drug screen and correlations with the growth-inhibitory potency of 123 anticancer agents. *Cancer Res.*, **57**:4285-4300.
- O'Connor, L., Strasser, A., O'Reilly, L.A., Hausmann, G., Adams, J.M., Cory, S., Huang, D.C., 1998. Bim: a novel member of the Bcl-2 family that promotes apoptosis. *EMBO J.*, **17**(2):384-395. [doi:10.1093/emboj/17.2.384]
- Oda, E., Ohki, R., Murasawa, H., Nemoto, J., Shibue, T., Yamashita, Y., Tokino, T., Taniguchi, T., Tanaka, N., 2000. Noxa, a BH3-only member of the bcl-2 family and candidate mediator of p53-induced apoptosis. *Science*, **288**(5468):1053-1058. [doi:10.1126/science.288.5468.1053]
- Ogilvy, S., Metcalf, D., Print, C.G., Bath, M.L., Harris, A.W., Adams, J.M., 1999. Constitutive Bcl-2 expression throughout the hematopoietic compartment affects multiple lineages and enhances progenitor cell survival. *Proc. Natl. Acad. Sci. (USA)*, **96**(26):14943-14948. [doi:10.1073/pnas.96.26.14943]
- Oltvai, Z.N., Millman, C.L., Korsmeyer, S.J., 1993. Bcl-2 heterodimerizes in vivo with a conserved homolog, Bax, that accelerates programmed cell death. *Cell*, **74**(4):609-619. [doi:10.1016/0092-8674(93)90509-O]
- O'Reilly, L.A., Huang, D.C.S., Strasser, A., 1996. The death inhibitor Bcl-2 and its homologues influence control of cell cycle. *EMBO J.*, **15**:6979-6990.
- O'Reilly, L.A., Harris, A.W., Strasser, A., 1997a. Bcl-2 transgene expression promotes survival and reduces proliferation of CD3-CD4-CD8-T cell progenitors. *Int. Immunol.*, **9**(9):1291-1301. [doi:10.1093/intimm/9.9.1291]
- O'Reilly, L.A., Harris, A.W., Tarlinton, D.M., Corcoran, L.M., Strasser, A., 1997b. Expression of a Bcl-2 transgene reduces proliferation and slows turnover of developing B lymphocytes in vivo. *J. Immunol.*, **159**:2301-2311.
- Parrish, J., Li, L., Klotz, K., Ledwich, D., Wang, X., Xue, D., 2001. Mitochondrial endonuclease G is important for apoptosis in *C. elegans*. *Nature*, **412**(6842):90-94. [doi:10.1038/35083608]
- Passalaris, T.M., Benanti, J.A., Gewin, L., Kiyono, T., Gallo-way, D.A., 1999. The G(2) checkpoint is maintained by redundant pathways. *Mol. Cell Biol.*, **19**:5872-5881.
- Pavlov, E.V., Priault, M., Pietkiewicz, D., Cheng, E.H.Y., Antonsson, B., Manon, S., Korsmeyer, S.J., Mannella, C.A., Kinnally, K.W., 2001. A novel, high conductance channel of mitochondria linked to apoptosis in mammalian cells and Bax expression in yeast. *J. Cell Biol.*, **155**(5):725-732. [doi:10.1083/jcb.200107057]
- Pediconi, N., Ianari, A., Costanzo, A., Belloni, L., Gallo, R., Cimino, L., Porcellini, A., Screpanti, I., Balsano, C., Alesse, E., Gulino, A., Massimo Levrero, M., 2003. Differential regulation of E2F1 apoptotic target genes in response to DNA damage. *Nat. Cell Biol.*, **5**(6):552-558. [doi:10.1038/ncb998]
- Peng, C.Y., Graves, P.R., Thoma, R.S., Wu, Z., Shaw, A.S., Piwnica-Worms, H., 1997. Mitotic and G2 checkpoint control: regulation of 14-3-3 protein binding by phosphorylation of Cdc25C on serine-216. *Science*, **277**(5331):1501-1505. [doi:10.1126/science.277.5331.1501]
- Perfettini, J.L., Kroemer, R.T., Kroemer, G., 2004. Fatal liaisons of p53 with bax and Bak. *Nat. Cell Biol.*, **6**(5):386-387. [doi:10.1038/ncb0504-386]
- Phillips, A.C., Bates, S., Ryan, K.M., Helin, K., Vousden, K.H., 1997. Induction of DNA synthesis and apoptosis are separable functions of E2F-1. *Genes Dev.*, **11**:1853-1863.
- Piret, B., Schoonbroodt, S., Piette, J., 1999. The ATM protein is required for sustained activation of NF-kappa B following DNA damage. *Oncogene*, **18**(13):2261-2271. [doi:10.1038/sj.onc.1202541]
- Polster, B.M., Kinnally, K.W., Fiskum, G., 2001. BH3 death domain peptide induces cell type-selective mitochondrial outer membrane permeability. *J. Biol. Chem.*, **276**:37887-37894.
- Poruchynsky, M.S., Wang, E.E., Rudin, C.M., Blagosklonny, M.V., Fojo, T., 1998. Bcl-X(L) is phosphorylated in malignant cells following microtubule disruption. *Cancer Res.*, **58**:3331-3338.
- Puthalakath, H., Huang, D.C.S., O'Reilly, L.A., King, S.M., Strasser, A., 1999. The proapoptotic activity of the Bcl-2 family member Bim is regulated by interaction with the dynein motor complex. *Mol. Cell*, **3**(3):287-296. [doi:10.1016/S1097-2765(00)80456-6]
- Puthalakath, H., Villunger, A., O'Reilly, L.A., Beaumont, J.G., Coultas, L., Cheney, R.E., Huang, D.C., Strasser, A., 2001. Bmf: a proapoptotic BH3-only protein regulated by interaction with the myosin V actin motor complex, activated by anoikis. *Science*, **293**(5536):1829-1832. [doi:10.1126/science.1062257]
- Qin, Z.H., Wang, Y., Kikly, K.K., Sapp, E., Kegel, K.B., Aronin, N., DiFiglia, M., 2001. Pro-caspase-8 is predominantly localized in mitochondria and released into cytoplasm upon apoptotic stimulation. *J. Biol. Chem.*, **276**(11):8079-8086. [doi:10.1074/jbc.M007028200]
- Reed, J.C., 1997. Double identity for proteins of the Bcl-2 family. *Nature*, **387**(6635):773-776. [doi:10.1038/42867]
- Reed, J.C., 1998. Bcl-2 family proteins. *Oncogene*, **17**(25):3225-3236. [doi:10.1038/sj.onc.1202591]
- Reed, J.C., 2006. Drug insight: cancer therapy strategies based on restoration of endogenous cell death mechanisms. *Nat. Clin. Pract. Oncol.*, **3**(7):388-398. [doi:10.1038/ncponc.0538]
- Riabowol, K., Draetta, G., Brizuela, L., Vandre, D., Beach, D., 1989. The cdc2 kinase is a nuclear protein that is essential

- for mitosis in mammalian cells. *Cell*, **57**(3):393-401. [doi:10.1016/0092-8674(89)90914-8]
- Robles, S.J., Adami, G.R., 1998. Agents that cause DNA double strand breaks lead to p16INK4a enrichment and the premature senescence of normal fibroblasts. *Oncogene*, **16**(9):1113-1123. [doi:10.1038/sj.onc.1201862]
- Ruvolo, P.P., Deng, X.M., Carr, B.H., May, W.S., 1998. A functional role for mitochondrial protein kinase C- α in Bcl2 phosphorylation and suppression of apoptosis. *J. Biol. Chem.*, **273**(39):25436-25442. [doi:10.1074/jbc.273.39.25436]
- Ryan, K.M., Ernst, M.K., Rice, N.R., Vousden, K.H., 2000. Role of NF κ B in p53-mediated programmed cell death. *Nature*, **404**(6780):892-896. [doi:10.1038/35009130]
- Saito, M., Korsmeyer, S.J., Schlesinger, P.H., 2000. Bax-dependent transport of cytochrome c reconstituted in pure liposomes. *Nat. Cell. Biol.*, **2**(8):553-555. [doi:10.1038/35019596]
- Sanchez, Y., Wong, C., Thoma, R.S., Richman, R., Wu, Z., Piwnicka-Worms, H., Elledge, S.J., 1997. Conservation of the Chk1 checkpoint pathway in mammals: linkage of DNA damage to Cdk regulation through Cdc25. *Science*, **277**(5331):1497-1501. [doi:10.1126/science.277.5331.1497]
- Sattler, M., Liang, H., Nettesheim, D., Meadows, R.P., Harlan, J.E., Eberstadt, M., Yoon, H.S., Shuker, S.B., Chang, B.S., Minn, A.J., et al., 1997. Structure of Bcl-x(L)-Bak peptide complex: recognition between regulators of apoptosis. *Science*, **275**(5302):983-986. [doi:10.1126/science.275.5302.983]
- Sax, J.K., Fei, P., Murphy, M.E., Bernhard, E., Korsmeyer, S.J., El-Deiry, W.S., 2002. BID regulation by p53 contributes to chemosensitivity. *Nat. Cell Biol.*, **4**(11):842-849. [doi:10.1038/ncb866]
- Scatena, C.D., Stewart, Z.A., Mays, D., Tang, L.J., Keefer, C.J., Leach, S.D., Pietenpol, J.A., 1998. Mitotic phosphorylation of Bcl-2 during normal cell cycle progression and Taxol-induced growth arrest. *J. Biol. Chem.*, **273**(46):30777-30784. [doi:10.1074/jbc.273.46.30777]
- Schendel, S.L., Montal, M., Reed, J.C., 1998. Bcl-2 family proteins as ion-channels. *Cell Death Differ.*, **5**(5):372-380. [doi:10.1038/sj.cdd.4400365]
- Schlesinger, P.H., Gross, A., Yin, X.M., Yamamoto, K., Saito, M., Waksman, G., Korsmeyer, S.J., 1997. Comparison of the ion channel characteristics of proapoptotic Bax and antiapoptotic Bcl-2. *Proc. Natl. Acad. Sci. (USA)*, **94**(21):11357-11362. [doi:10.1073/pnas.94.21.11357]
- Schmitt, C.A., 2007. Cellular senescence and cancer treatment. *Biochim. Biophys. Acta*, **1775**:5-20.
- Schmitt, C.A., Fridman, J.S., Yang, M., Lee, S., Baranov, E., Hoffman, R.M., Lowe, S.W., 2002. A senescence program controlled by p53 and p16(INK4a) contributes to the outcome of cancer therapy. *Cell*, **109**(3):335-346. [doi:10.1016/S0092-8674(02)00734-1]
- Schmitt, E., Paquet, C., Beauchemin, M., Bertrand, R., 2004. Bcl-xES, a BH4- and BH2-containing antiapoptotic protein, delays Bax oligomer formation and binds Apaf-1, blocking procaspase 9 activation. *Oncogene*, **23**(22):3915-3931. [doi:10.1038/sj.onc.1207554]
- Schmitt, E., Boutros, R., Froment, C., Monsarrat, B., Ducommun, B., Dozier, C., 2006. CHK1 phosphorylates CDC25B during the cell cycle in the absence of DNA damage. *J. Cell Sci.*, **119**(20):4269-4275. [doi:10.1242/jcs.03200]
- Schmitt, E., Beauchemin, M., Bertrand, R., 2007. Nuclear co-localization and interaction between bcl-xL and cdk1(cdc2) during G2/M cell cycle checkpoint. *Oncogene*, in press. [doi:10.1038/sj.onc.1210396]
- Sedlak, T.W., Oltvai, Z.N., Yang, E., Wang, K., Boise, L.H., Thompson, C.B., Korsmeyer, S.J., 1995. Multiple Bcl-2 family members demonstrate selective dimerizations with Bax. *Proc. Natl. Acad. Sci. (USA)*, **92**(17):7834-7838. [doi:10.1073/pnas.92.17.7834]
- Sherr, C.J., Roberts, J.M., 1995. Inhibitors of mammalian G1 cyclin-dependent kinases. *Genes. Dev.*, **9**:1149-1163.
- Shibue, T., Takeda, K., Oda, E., Tanaka, H., Murasawa, H., Takaoka, A., Morishita, Y., Akira, S., Taniguchi, T., Tanaka, N., 2003. Integral role of Noxa in p53-mediated apoptotic response. *Genes Dev.*, **17**(18):2233-2238. [doi:10.1101/gad.1103603]
- Shiloh, Y., 2001. ATM and ATR: networking cellular responses to DNA damage. *Curr. Opin. Genet. Dev.*, **11**(1):71-77. [doi:10.1016/S0959-437X(00)00159-3]
- Shiloh, Y., 2006. The ATM-mediated DNA-damage response: taking shape. *Trends Biochem. Sci.*, **31**(7):402-410. [doi:10.1016/j.tibs.2006.05.004]
- Shimizu, T., Pommier, Y., 1997. Camptothecin-induced apoptosis in p53-null human leukemia HL60 cells and their isolated nuclei; Effects of the protease inhibitors z-VAD-fmk and dichloroisocoumarin suggest an involvement of both caspases and serine proteases. *Leukemia*, **11**(8):1238-1244. [doi:10.1038/sj.leu.2400734]
- Shimizu, S., Tsujimoto, Y., 2000. Proapoptotic BH3-only Bcl-2 family members induce cytochrome c release, but not mitochondrial membrane potential loss, and do not directly modulate voltage-dependent anion channel activity. *Proc. Natl. Acad. Sci. (USA)*, **97**(2):577-582. [doi:10.1073/pnas.97.2.577]
- Shimizu, S., Eguchi, Y., Kamiike, W., Funahashi, Y., Mignon, A., Lacronique, V., Matsuda, H., Tsujimoto, Y., 1998. Bcl-2 prevents apoptotic mitochondrial dysfunction by regulating proton flux. *Proc. Natl. Acad. Sci. (USA)*, **95**(4):1455-1459. [doi:10.1073/pnas.95.4.1455]
- Shimizu, S., Narita, M., Tsujimoto, Y., 1999. Bcl-2 family proteins regulate the release of apoptogenic cytochrome c by the mitochondrial channel VDAC. *Nature*, **399**(6735):483-487. [doi:10.1038/20959]
- Shimizu, S., Konishi, A., Kodama, T., Tsujimoto, Y., 2000. BH4 domain of antiapoptotic bcl-2 family members closes voltage-dependent anion channel and inhibits apoptotic mitochondrial changes and cell death. *Proc. Natl. Acad. Sci. (USA)*, **97**(7):3100-3105. [doi:10.1073/pnas.97.7.3100]
- Slichenmyer, W.J., Nelson, W.G., Slebos, R.J., Kastan, M.B.,

1993. Loss of a p53-associated G1 checkpoint does not decrease cell survival following DNA damage. *Cancer Res.*, **53**:4164-4168.
- Smits, V.A.J., Klompmaier, R., Arnaud, L., Rijksen, G., Nigg, E.A., Medema, R.H., 2000. Polo-like kinase-1 is a target of the DNA damage checkpoint. *Nat. Cell Biol.*, **2**(9): 672-676. [doi:10.1038/35023629]
- Solary, E., Bertrand, R., Jenkins, J., Pommier, Y., 1992. Radiolabeling of DNA can induce its fragmentation in HL-60 promyelocytic leukemic cells. *Exp. Cell Res.*, **203**(2):495-498. [doi:10.1016/0014-4827(92)90027-6]
- Song, Q.Z., Kuang, Y.P., Dixit, V.M., Vincenz, C., 1999. Boo, a novel negative regulator of cell death, interacts with Apaf-1. *EMBO J.*, **18**(1):167-178. [doi:10.1093/emboj/18.1.167]
- Srivastava, R.K., Srivastava, A.R., Korsmeyer, S.J., Nesterova, M., Chochung, Y.S., Longo, D.L., 1998. Involvement of microtubules in the regulation of Bcl2 phosphorylation and apoptosis through cyclic AMP-dependent protein kinase. *Mol. Cell Biol.*, **18**:3509-3517.
- Srivastava, R.K., Mi, Q.S., Hardwick, J.M., Longo, D.L., 1999. Deletion of the loop region of Bcl-2 completely blocks paclitaxel-induced apoptosis. *Proc. Natl. Acad. Sci. (USA)*, **96**(7):3775-3780. [doi:10.1073/pnas.96.7.3775]
- Stanelle, J., Stiewe, T., Theseling, C.C., Peter, M., Pützer, B.M., 2002. Gene expression changes in response to E2F1 activation. *Nucleic. Acids Res.*, **30**(8):1859-1867. [doi:10.1093/nar/30.8.1859]
- Stehlik, C., Demartin, R., Kumabashiri, I., Schmid, J.A., Binder, B.R., Lipp, J., 1998. Nuclear factor (NF)-kappaB-regulated X-chromosome-linked Iap gene expression protects endothelial cells from tumor necrosis factor alpha-induced apoptosis. *J. Exp. Med.*, **188**(1): 211-216. [doi:10.1084/jem.188.1.211]
- Stiewe, T., Pützer, B.M., 2000. Role of the p53-homologue p73 in E2F1-induced apoptosis. *Nat. Genet.*, **26**(4):464-469. [doi:10.1038/82617]
- Stroka, D.M., Badrichani, A.Z., Bach, F.H., Ferran, C., 1999. Overexpression of A1, an NF-kappaB-inducible anti-apoptotic Bel gene, inhibits endothelial cell activation. *Blood*, **93**:3803-3810.
- Strunnikova, M., Schagdarsurengin, U., Kehlen, A., Garbe, J.C., Stampfer, M.R., Dammann, R., 2005. Chromatin inactivation precedes de novo DNA methylation during the progressive epigenetic silencing of the RASSF1A promoter. *Mol. Cell Biol.*, **25**(10):3923-3933. [doi:10.1128/MCB.25.10.3923-3933.2005]
- Sugiyama, T., Shimizu, S., Matsuoka, Y., Yoneda, Y., Tsujimoto, Y., 2002. Activation of mitochondrial voltage-dependent anion channel by pro-apoptotic BH3-only protein Bim. *Oncogene*, **21**(32):4944-4956. [doi:10.1038/sj.onc.1205621]
- Sun, L.Q., Arcesi, R.J., 2005. Altered epigenetic patterning leading to replicative senescence and reduced longevity. A role of a novel SNF2 factor, PASG. *Cell Cycle*, **4**:3-5.
- Susin, S.A., Zamzami, N., Castedo, M., Hirsch, T., Marchetti, P., Macho, A., Daugas, E., Geuskens, M., Kroemer, G., 1996. Bcl-2 inhibits the mitochondrial release of an apoptogenic protease. *J. Exp. Med.*, **184**(4):1331-1341. [doi:10.1084/jem.184.4.1331]
- Susin, S.A., Lorenzo, H.K., Zamzami, N., Isabel Marzo, I., Snow, B.E., Brothers, G.M., Mangion, J., Jacotot, E., Costantini, P., Loeffler, M., et al., 1999a. Molecular characterization of mitochondrial apoptosis-inducing factor. *Nature*, **397**(6718):441-446. [doi:10.1038/17135]
- Susin, S.A., Lorenzo, H.K., Zamzami, N., Marzo, I., Brenner, C., Larochette, N., Prevost, M.C., Alzari, P.M., Kroemer, G., 1999b. Mitochondrial release of caspase-2 and -9 during the apoptotic process. *J. Exp. Med.*, **189**(2): 381-393. [doi:10.1084/jem.189.2.381]
- Suzuki, K., Mori, I., Nakayama, Y., Miyakoda, M., Komada, S., Watanabe, M., 2001a. Radiation-induced senescence-like growth arrest requires TP53 function but not telomere shortening. *Radiat. Res.*, **155**(1):248-253. [doi:10.1667/0033-7587(2001)155[0248:RISLGA]2.0.CO;2]
- Suzuki, Y., Imai, Y., Nakayama, H., Takahashi, K., Takio, K., Takahashi, R., 2001b. A serine protease, Htra2, is released from the mitochondria and interacts with Xiap, inducing cell death. *Mol. Cell*, **8**(3):613-621. [doi:10.1016/S1097-2765(01)00341-0]
- Takahashi, A., Ohtani, N., Yamakoshi, K., Iida, S., Tahara, H., Nakayama, K., Nakayama, K.I., Ide, T., Saya, H., Hara, E., 2006. Mitogenic signalling and the p16INK4a-Rb pathway cooperate to enforce irreversible cellular senescence. *Nat. Cell Biol.*, **8**(11):1291-1297. [doi:10.1038/ncb1491]
- Tamaru, H., Selker, E.U., 2001. A histone H3 methyltransferase controls DNA methylation in *Neurospora crassa*. *Nature*, **414**(6861):277-283. [doi:10.1038/35104508]
- Tamaru, H., Zhang, X., McMillen, D., Singh, P.B., Nakayama, J., Grewal, S.I., Allis, C.D., Cheng, X., Selker, E.U., 2003. Trimethylated lysine 9 of histone H3 is a mark for DNA methylation in *Neurospora crassa*. *Nat. Genet.*, **34**(1): 75-79. [doi:10.1038/ng1143]
- Tamatani, M., Che, Y.H., Matsuzaki, H., Ogawa, S., Okado, H., Miyake, S., Mizuno, T., Tohyama, M., 1999. Tumor necrosis factor induces Bcl-2 and Bcl-x expression through NF kappa B activation in primary hippocampal neurons. *J. Biol. Chem.*, **274**(13):8531-8538. [doi:10.1074/jbc.274.13.8531]
- Tan, K.O., Tan, K.M.L., Chan, S.L., Yee, K.S.Y., Bevert, M., Ang, K.C., Yu, V.C., 2001. MAP-1, a novel proapoptotic protein containing a BH3-like motif that associates with Bax through its Bcl-2 homology domains. *J. Biol. Chem.*, **276**(4):2802-2807. [doi:10.1074/jbc.M008955200]
- Tanaka, N., Ishihara, M., Lamphier, M.S., Nozawa, H., Matsuyama, T., Mak, T.W., Aizawa, S., Tokino, T., Oren, M., Taniguchi, T., 1996. Cooperation of the tumour suppressors Irf-1 and p53 in response to DNA damage. *Nature*, **382**(6594):816-818. [doi:10.1038/382816a0]
- Thomas, A., Elrouby, S., Reed, J.C., Krajewski, S., Silber, R., Potmteil, M., Newcomb, E.W., 1996. Drug-induced apoptosis in B-cell chronic lymphocytic leukemia: relationship between p53 gene mutation and Bcl-2/Bax proteins in drug resistance. *Oncogene*, **12**:1055-1062.

- Tollefsbol, T.O., Andrews, L.G., 1993. Mechanisms for methylation-mediated gene silencing and aging. *Med. Hypotheses*, **41**(1):83-92. [doi:10.1016/0306-9877(93)90040-W]
- Tsujimoto, Y., Finger, L.R., Yunis, J., Nowell, P.C., Croce, C.M., 1984. Cloning of the chromosome breakpoint of neoplastic B cells with the t(14:18) chromosome translocation. *Science*, **226**(4678):1097-1099. [doi:10.1126/science.6093263]
- Vail, M.E., Chaisson, M.L., Thompson, J., Fausto, N., 2002. Bcl-2 expression delays hepatocyte cell cycle progression during liver regeneration. *Oncogene*, **21**(10):1548-1555. [doi:10.1038/sj.onc.1205212]
- Vairo, G., Innes, K.M., Adams, J.M., 1996. Bcl-2 has a cell cycle inhibitory function separable from its enhancement of cell survival. *Oncogene*, **13**:1511-1519.
- van Vugt, M.A., Smits, V.A., Klompaker, R., Medema, R.H., 2001. Inhibition of Polo-like kinase-1 by DNA damage occurs in an ATM- or ATR-dependent fashion. *J. Biol. Chem.*, **276**(45):41656-41660. [doi:10.1074/jbc.M101831200]
- Vander Heiden, M.G., Thompson, C.B., 1999. Bcl-2 proteins: regulators of apoptosis or of mitochondrial homeostasis? *Nat. Cell Biol.*, **1**(8):E209-E216. [doi:10.1038/70237]
- Vander Heiden, M.G., Chandel, N.S., Williamson, E.K., Schumacker, P.T., Thompson, C.B., 1997. Bcl-X(L) regulates the membrane potential and volume homeostasis of mitochondria. *Cell*, **91**(5):627-637. [doi:10.1016/S0092-8674(00)80450-X]
- Vaux, D.L., Cory, S., Adams, J.M., 1988. Bcl-2 gene promotes haemopoietic cell survival and cooperates with c-myc to immortalize pre-B cells. *Nature*, **335**(6189):440-442. [doi:10.1038/335440a0]
- Verhagen, A.M., Ekert, P.G., Pakusch, M., Silke, J., Connolly, L.M., Reid, G.E., Moritz, R.L., Simpson, R.J., Vaux, D.L., 2000. Identification of DIABLO, a mammalian protein that promotes apoptosis by binding to and antagonizing IAP proteins. *Cell*, **102**(1):43-53. [doi:10.1016/S0092-8674(00)00009-X]
- Villunger, A., Michalak, E.M., Coultas, L., Müllauer, F., Böck, G., Ausserlechner, M.J., Adams, J.M., Strasser, A., 2003. p53- and drug-induced apoptotic responses mediated by BH3-only proteins Puma and Noxa. *Science*, **302**(5647):1036-1038. [doi:10.1126/science.1090072]
- Waldman, T., Lengauer, C., Kinzler, K.W., Vogelstein, B., 1996. Uncoupling of S phase and mitosis induced by anticancer agents in cells lacking p21. *Nature*, **381**(6584):713-716. [doi:10.1038/381713a0]
- Wang, K., Yin, X.M., Chao, D.T., Milliman, C.L., Korsmeyer, S.J., 1996. BID: a novel BH3 domain-only death agonist. *Genes. Dev.*, **10**:2859-2869.
- Wang, C.Y., Mayo, M.W., Korneluk, R.G., Goeddel, D.V., Baldwin, A.S., 1998. NF-kappa-B antiapoptosis: induction of TRAF1 and TRAF2 and c-IAP1 and c-IAP2 to suppress caspase-8 activation. *Science*, **281**(5383):1680-1683. [doi:10.1126/science.281.5383.1680]
- Wang, X.W., Zhan, Q.M., Coursen, J.D., Khan, M.A., Kontny, H.U., Yu, L., M. Hollander, M.C., O'Connor, P.M., Fornace, A.J.Jr, Harris, C.C., 1999a. GADD45 induction of a G(2)/M cell cycle checkpoint. *Proc. Natl. Acad. Sci. (USA)*, **96**(7):3706-3711. [doi:10.1073/pnas.96.7.3706]
- Wang, C.Y., Guttridge, D.C., Mayo, M.W., Baldwin, A.S.Jr, 1999b. NF-kappaB induces expression of the Bcl-2 homologue A1/Bfl-1 to preferentially suppress chemotherapy-induced apoptosis. *Mol. Cell Biol.*, **19**:5923-5929.
- Willis, S.N., Chen, L., Dewson, G., Wei, A., Naik, E., Fletcher, J.I., Adams, J.M., Huang, D.C.S., 2005. Proapoptotic Bak is sequestered by Mcl-1 and Bcl-xL, but not Bcl-2, until displaced by BH3-only proteins. *Genes. Dev.*, **19**(11):1294-1305. [doi:10.1101/gad.1304105]
- Willis, S.N., Fletcher, J.I., Kaufmann, T., van Delft, M.F., Chen, L., Czabotar, P.E., Ierino, H., Lee, E.H., Fairlie, W.D., Bouillet, P., et al., 2007. Apoptosis initiated when BH3 ligands engage multiple Bcl-2 homologs, not Bax or Bak. *Science*, **315**(5813):856-859. [doi:10.1126/science.1133289]
- Xie, S., Wu, H., Wang, Q., Cogswell, J.P., Husain, I., Conn, C., Stambrook, P., Jhanwar-Uniyal, M., Dai, W., 2001. Plk3 functionally links DNA damage to cell cycle arrest and apoptosis at least in part via the p53 pathway. *J. Biol. Chem.*, **276**(46):43305-43312. [doi:10.1074/jbc.M106050200]
- Yamamoto, K., Ichijo, H., Korsmeyer, S.J., 1999. Bcl-2 is phosphorylated and inactivated by an ASK1/Jun N-terminal protein kinase pathway normally activated at G(2)/M. *Mol. Cell Biol.*, **19**:8469-8478.
- Yang, E., Zha, J., Jockel, J., Boise, L.H., Thompson, C.B., Korsmeyer, S.J., 1995. Bad, a heterodimeric partner for Bcl-xL and Bcl-2, displaces Bax and promotes cell death. *Cell*, **80**(2):285-291. [doi:10.1016/0092-8674(95)90411-5]
- Yang, J., Liu, X., Bhalla, K., Kim, C.N., Ibrado, A.M., Cai, J., Peng, T.I., Jones, D.P., Wang, X., 1997. Prevention of apoptosis by Bcl-2: release of cytochrome C from mitochondria blocked. *Science*, **275**(5303):1129-1132. [doi:10.1126/science.275.5303.1129]
- Yang, A.N., Kaghad, M., Wang, Y.M., Gillett, E., Fleming, M.D., Dotsch, V., Andrews, N.C., Caput, D., McKeon, F., 1998. p63, a p53 homolog at 3q27-29, encodes multiple products with transactivating, death-inducing, and dominant-negative activities. *Mol. Cell*, **2**(3):305-316. [doi:10.1016/S1097-2765(00)80275-0]
- Yasuda, M., Han, J.W., Dionne, C.A., Boyd, J.M., Chinnadurai, G., 1999. BNIP3 alpha: a human homolog of mitochondrial proapoptotic protein BNIP3. *Cancer Res.*, **59**:533-537.
- Yin, X.M., Oltvai, Z.N., Korsmeyer, S.J., 1994. BH1 and BH2 domains of Bcl-2 are required for inhibition of apoptosis and heterodimerization with Bax. *Nature*, **369**(6478):321-323. [doi:10.1038/369321a0]
- Yu, J., Zhang, L., Hwang, P.M., Kinzler, K.W., Vogelstein, B., 2001. Puma induces the rapid apoptosis of colorectal cancer cells. *Mol. Cell*, **7**(3):673-682. [doi:10.1016/S1097-2765(01)00213-1]

- Yuan, Z.M., Shioya, H., Ishiko, T., Sun, X., Gu, J., Huang, Y.Y., Lu, H., Kharbanda, S., Weichselbaum, R., Kufe, D., 1999. p73 is regulated by tyrosine kinase c-Abl in the apoptotic response to DNA damage. *Nature*, **399**(6738):814-817. [doi:10.1038/21704]
- Zamzami, N., Kroemer, G., 2001. The mitochondrion in apoptosis: how Pandora's box opens. *Nat. Rev. Mol. Cell Biol.*, **2**(1):67-71. [doi:10.1038/35048073]
- Zamzami, N., Marchetti, P., Castedo, M., Decaudin, D., Macho, A., Hirsch, T., Susin, S.A., Petit, P.X., Mignotte, B., Kroemer, G., 1995. Sequential reduction of mitochondrial transmembrane potential and generation of reactive oxygen species in early programmed cell death. *J. Exp. Med.*, **182**(2):367-377. [doi:10.1084/jem.182.2.367]
- Zamzami, N., Susin, S.A., Marchetti, P., Hirsch, T., Gomez-Monterrey, I., Castedo, M., Kroemer, G., 1996. Mitochondrial control of nuclear apoptosis. *J. Exp. Med.*, **183**(4):1533-1544. [doi:10.1084/jem.183.4.1533]
- Zamzami, N., Brenner, C., Marzo, I., Susin, S.A., Kroemer, G., 1998. Subcellular and submitochondrial mode of action of Bcl-2-like oncoproteins. *Oncogene*, **16**(17):2265-2282. [doi:10.1038/sj.onc.1201989]
- Zeng, Y., Forbes, K.C., Wu, Z., Moreno, S., Piwnica-Worms, H., Enoch, T., 1998. Replication checkpoint requires phosphorylation of the phosphatase Cdc25 by Cds1 or Chk1. *Nature*, **395**(6701):507-510. [doi:10.1038/26766]
- Zhan, Q., Antinore, M.J., Wang, X.W., Carrier, F., Smith, M.L., Harris, C.C., Fornace, A.J.Jr., 1999. Association with Cdc2 and inhibition of Cdc2/Cyclin B1 kinase activity by the p53-regulated protein Gadd45. *Oncogene*, **18**(18):2892-2900. [doi:10.1038/sj.onc.1202667]
- Zhang, H., Holzgreve, W., de Geyter, C., 2001. Bcl2-L-10, a novel anti-apoptotic member of the Bcl-2 family, blocks apoptosis in the mitochondria death pathway but not in the death receptor pathway. *Hum. Mol. Genet.*, **10**(21):2329-2339. [doi:10.1093/hmg/10.21.2329]
- Zhang, R., Chen, W., Adams, P.D., 2007. Molecular dissection of formation of senescent associated heterochromatin foci. *Mol. Cell. Biol.*, **27**(6):2343-2358. [doi:10.1128/MCB.02019-06]
- Zinkel, S.S., Hurov, K.E., Ong, C., Abtahi, F.M., Gross, A., Korsmeyer, S.J., 2005. A role for proapoptotic BID in the DNA-damage response. *Cell*, **122**(4):579-591. [doi:10.1016/j.cell.2005.06.022]
- Zong, W.X., Edelstein, L.C., Chen, C.L., Bash, J., Gelinas, C., 1999. The prosurvival Bcl-2 homolog Bfl-1/A1 is a direct transcriptional target of NF-kappaB that blocks TNF alpha-induced apoptosis. *Genes. Dev.*, **13**:382-387.
- Zoratti, M., Szabo, I., 1995. The mitochondrial permeability transition. *Biochim. Biophys. Acta*, **1241**:139-176.

Editors-in-Chief: Wei YANG & Peter H. BYERS
ISSN 1673-1581 (Print); ISSN 1862-1783 (Online), monthly

Journal of Zhejiang University **SCIENCE B**

www.zju.edu.cn/jzus; www.springerlink.com

jzus@zju.edu.cn

JZUS-B focuses on "Biomedicine, Biochemistry & Biotechnology"

JZUS-B online in PMC: <http://www.pubmedcentral.nih.gov/tocrender.fcgi?journal=371&action=archive>

Welcome Contributions to JZUS-B