- All letters must be typed with double spacing and signed by all authors.
- No letter should be more than 400 words.
- For letters on scientific subjects we normally reserve our correspondence columns for those relating to issues discussed recently (within six weeks) in the *BMJ*.
- We do not routinely acknowledge letters. Please send a stamped addressed envelope if you would like an acknowledgment.
- Because we receive many more letters than we can publish we may shorten those we do print, particularly when we receive several on the same subject.

Searching published reports

SIR,—All learned journals require that their authors have satisfactorily reviewed published work. Over the past decade there have been several critical reviews of authors' use of statistics¹ and recommendations to try to tighten up the use and presentation of statistics.² Is it now time to do the same for the review of published reports? It sometimes seems as if authors have not read, or choose to ignore, previous reports.

Short reports in the *BMJ*, by the rules of submission, must have no more than five references. In their short report on patients' access to their own psychiatric records Morris Bernadt and colleagues make use of only three,³ one of which is to the Access to Health Records Act. They begin their discussion with the words, "The few reports that there are about the consequences of patients having access to their medical records. . . ." This assertion is not supported by examination of published work: two recent reviews cited a total of 31 papers which appeared to be reports of original research and included many with results of the consequences of patients having access to their medical records.⁴ For example, there are reports on the effects on patients' anxiety,⁵ on the number of records lost,⁶ and on corrections to the record.⁷

Later in their discussion Bernadt and colleagues claim, "No study has systematically examined the influence of demographic data and diagnosis." Yet, for example, there are reports looking at the influence of age, sex, and diagnoses on the censoring of patient held records.⁸ The authors also state, "We know of only one study which examined patients who requested to see their records as opposed to research recruits. . . ." In several of the studies reported previously—for example, that of Jones *et al.*⁹—records have routinely been given to all patients in a given category, and so it is inappropriate to call these patients research recruits.

In discussing the arguments for and against patients' access to records, Kirby opens with "Clinicians' reactions to the idea of greater patient access to case notes vary from lukewarm support to downright opposition. . . ." Kirby does not back this claim with evidence. In fact, several papers and letters from clinicians offer their enthusiastic support.¹⁰ Later Kirby argues, "There is no systematic evidence" (on the contrasting degrees of enthusiasm between hospital doctors and general practitioners), ignoring several British hospital studies of women receiving antenatal care holding their own records,¹¹ the issue of a computer produced summary in a hospital diabetic clinic,¹² and the copying of letters from hospital consultants to general practitioners to patients.¹³

Of the 31 papers listed in the two reviews,¹⁴ 25 were listed on MEDLINE and most of these would have been found by using the MESH headings (a) "medical records" and "truth disclosure";

(b) "medical records" and "patient education"; (c) "medical records" and "attitude to health"; (d) "medical records" and "patient participation"; or (e) "medical records" and "health education." (A list of 101 references on the subject of patients' access to records, of which 53 are to original research, is available.)

If scientific knowledge of medicine and health services is to progress we need to make full use of the work of others. In the United States MEDLINE is now being used effectively in routine clinical practice.¹¹ Jennett *et al* argue that the use of a bibliographic searching tool such as MEDLINE should be an element in the medical undergraduate curriculum.¹² Perhaps also research journals should require their authors to include details of their method of searching published work.

RAY JONES

Department of Public Health,
University of Glasgow,
Glasgow G12 8RZ

- 1 Gore SM, Jones IG, Rytter EC. Misuse of statistical methods: critical assessment of articles in the British Medical Journal from January to March 1976. *BMJ* 1977;ii:85-7.
- 2 Altman DG, Gore SM, Gardner MJ, Pocock SJ. Statistical guidelines for contributors to medical journals. *BMJ* 1983; 286:1489-93.
- 3 Bernadt M, Gunning L, Quenstedt M. Patients' access to their own psychiatric records. *BMJ* 1991;303:967. (19 October.)
- 4 McGhee SM, Hedley AJ, Jones RB, Cheng KK. Patient held records: their current status and implications for health care in Hong Kong. *Hong Kong Practitioner* 1991;13:1374-81.
- 5 Gilhooly MLM, McGhee SM. Medical records: practicalities and principles of patient possession. *J Med Ethics* 1991;17: 138-43.
- 6 Baldry M, Cheal C, Fisher B, Gillet M, Huet V. Giving patients their own medical records in general practice: experience of patients and staff. *BMJ* 1986;292:596-8.
- 7 Elbourne D, Richardson M, Chalmers I, Waterhouse I, Holt E. The Newbury maternity care study: a randomized controlled trial to assess a policy of women holding their own obstetric records. *Br J Obstet Gynaecol* 1987;94:612-9.
- 8 Jones RB, Hedley AJ, Allison SP, Tattersall RB. Censoring of patient-held records by doctors. *J R Coll Gen Pract* 1988;38: 117-8.
- 9 Kirby BJ. Patient access to medical records. *J R Coll Phys Lond* 1991;25:240-2.
- 10 Tattersall RB. Writing for and to patients. *Diabetic Med* 1990;7:917-9.
- 11 Lindberg DAB, Siegel ER. On assessing the impact of medical information: does MEDLINE make a difference? *Methods Inf Med* 1991;30:239-40.
- 12 Jennett PA, Edworthy SM, Rosenal TW, Maes WR, Yee N, Jardine PG. Preparing doctors for tomorrow: information management as a theme in undergraduate medical education. *Med Educ* 1991;25:135-9.

Hazards of pharmacological tests of growth hormone secretion

SIR,—A Shah and colleagues show courage and candour in reporting two deaths and serious neurological sequelae after tests of growth hormone secretion in children.¹ Although there is a system for reporting adverse medical events after the

administration of drugs, there is no equivalent for documenting problems after investigations and alerting the medical community of associated risks, and thus their report is valuable. It is well known among growth specialists, however, that insulin induced hypoglycaemia is potentially dangerous, and at least one child was previously known to have died after an insulin tolerance test.

Shah and colleagues highlight the danger of reversing the hypoglycaemia with excessive quantities of hyperosmolar glucose solutions, which probably caused or exacerbated cerebral oedema. But I challenge the necessity of using a test that carries such a risk, whatever the mechanism of cerebral oedema and death, to investigate short stature when other, safer tests are available.

The insulin tolerance test is still probably the most commonly used provocation test for growth hormone secretion and is regarded as the gold standard. When the Health Services Human Growth Hormone Committee investigated the relative value of several tests the insulin tolerance test performed as poorly as others, the rate of false positive results being 25%.² The technical value of the test is that it assesses hypothalamo-pituitary-adrenal status as well, but this information, adequate for clinical purposes, can be inferred from other tests. (At Manchester growth clinic we have diagnosed and categorised growth disorders and hypopituitarism in children without using the insulin tolerance test for the past 10 years.)

Diagnosing and defining isolated growth hormone deficiency and more generalised hypopituitarism in children will continue to be important so that they are managed with appropriate treatment regimens. Safe pharmacological and physiological stimuli of growth hormone secretion are available, and non-invasive investigations such as urinary excretion of growth hormone are under assessment.³ With the publication of Shah and colleagues' paper it seems unlikely that parents will give their informed consent readily, and the insulin tolerance test will find its rightful place in medical history.

D A PRICE

St Mary's Hospital,
Manchester M13 0JH

- 1 Shah A, Stanhope R, Matthew D. Hazards of pharmacological tests of growth hormone secretion in childhood. *BMJ* 1992;304:173-4. (18 January.)
- 2 Health Services Human Growth Hormone Committee. Comparison of the intravenous insulin and oral clonidine tests for growth hormone secretion. *Arch Dis Child* 1981;56:852-4.
- 3 Walker JM, Wood PJ, Williamson S, Betts PR, Evans AJ. Urinary growth hormone excretion as a screening test for growth hormone deficiency. *Arch Dis Child* 1990;65:89-92.

SIR—A Shah and colleagues' Lesson of the Week is a sharp and regrettable reminder to all clinicians, including gynaecologists, of the hazards of pituitary challenge tests.¹