resulting in inadequate monitoring of 21 patients. Secondly, urine output was not monitored in 34 patients.

Thirdly, shocked patients (blood pressure <100 mmHg, postural drop in blood pressure, thirst, tachycardia) were identified but not always monitored as closely as they should have been. Of the eight shocked patients, four did not have a urinary catheter, six received colloid in the first hour, and five received blood in the first four hours.

Fourthly, blood transfusion was judged to be late (>24 hours after admission) or insufficient owing to underestimation of blood loss in nine patients.

Fifthly, surgical assessment on admission was delayed by more than six hours in 24 patients.

Finally, abnormal clotting was not corrected in two patients.

It is salutary that the deficiencies in the management of these patients were largely due to the failure to apply simple clinical and nursing measures rather than any deficiency of high technology medicine. Careful haemodynamic monitoring and fluid and blood replacement have always been the key to managing patients with gastrointestinal bleeding successfully. This simple audit serves to re-emphasise this.

On the basis of the deficiencies identified by the audit the protocol has been modified. The revised protocol will again be given to all junior staff concerned in managing patients with gastrointestinal bleeding and will be incorporated into a handbook of protocols for common medical emergencies being compiled for junior staff.

M G S DUNNILL
S R GOULD

Department of Medicine,
Epsom General Hospital,
Epsom,
Surrey KT18 7EG

- 1 Daneshmend TK, Hawkey CJ, Langman MJS, Logan RFA, Long RG, Walt RP. Omeprazole versus placebo for acute upper gastrointestinal bleeding: randomised double blind controlled trial. *BMJ* 1992;304:143-7. (18 January.)
- 2 Madden MV, Griffiths GH. Management of upper gastrointestinal bleeding in a district general hospital. *J R Coll Phys Lond* 1986;20:212-5.
- 3 Management of gastrointestinal bleeding. *BMJ* 1981;283:456-7.

Psychological aspects of lower urinary tract infections in women

SIR,—In their article on the psychological aspects of lower urinary tract infections in women David Sumners and colleagues make the point that assessment of clinical symptoms alone was not helpful in distinguishing between true urinary tract infection and the urethral syndrome in women presenting with dysuria and frequency.¹ They also suggest that antibiotics might be prescribed more appropriately if significant bacteriuria could be predicted before the result of culture was available. We agree with both these points but suggest that this would be the case if microscopic examination of the urine specimen was undertaken in the practice straight after the sample was passed.

Several studies have shown the value of early microscopic examination in diagnosing urinary tract infection in both general practice and hospital.^{2,3} Not only can pyuria be detected in fresh, unspun urine but recent work has shown that bacteriuria can be recognised and is more specific for diagnosing urinary tract infection than pyuria alone.⁴ Uninfected samples can be recognised and discarded, which saves laboratory expense, unnecessary antibiotics, and costly investigation.

Obtaining the equipment to perform urine microscopy need not be prohibitively expensive.

We have found that a mirrored counting chamber costing under £50 (Hawksley and Sons, Lancing, Sussex) gives good definition and allows bacteria to be recognised easily even with a cheap light microscope. Phase contrast microscopy gives optimum definition, and a relatively inexpensive model is now available—namely, the updated McArthur microscope (K W Kirk and Sons, Milton, Cambridge), which has been around as a light microscope since 1974. This has the advantage of measuring only 10×9×5 cm and being very robust. Coltman comprehensively documented its usefulness in general practice.²

Our experience has been that microscopic examination of urine is not a difficult skill to acquire, and the time saved in writing out laboratory forms and chasing up results more than compensates for the short time it takes to look at the specimen, in addition to the satisfaction of making an instant diagnosis for the patient.

A F MELLON
M G COULTHARD

Department of Child Health,
Royal Victoria Infirmary,
Newcastle upon Tyne NE1 4LP

J SHAW

Ethel Street Practice,
Benwell,
Newcastle upon Tyne

J MILLAR

Grove Practice,
Gosforth,
Newcastle upon Tyne NE3 1NU

- 1 Sumners D, Kelsey M, Chait I. Psychological aspects of lower urinary tract infections in women. *BMJ* 1992;304:17-9. (4 January.)
- 2 Coltman KMD. A new technique for detecting urinary tract infection. *Practitioner* 1978;221:243-6.
- 3 Ditchburn RK, Ditchburn JS. A study of microscopical and chemical tests for the rapid diagnosis of urinary tract infections in general practice. *Br J Gen Pract* 1990;40:406-8.
- 4 Robins DG, White RHR, Rogers KB, Osman MS. Urine microscopy as an aid to detection of bacteriuria. *Lancet* 1975;i:476-8.
- 5 Vickers D, Ahmad T, Coulthard MG. Diagnosis of urinary tract infection in children: fresh urine microscopy or culture? *Lancet* 1991;338:767-70.

SIR,—In their study of the psychological aspects of lower urinary tract infections in women David Sumners and colleagues attempt to differentiate between urinary tract infection and the urethral syndrome by using “case notes, clinical symptoms, or psychiatric state.”¹ Their criteria for diagnosing the urethral syndrome, however, are unclear, and they seem not to have included infection with organisms such as *Neisseria gonorrhoeae*, *Chlamydia trachomatis*, and *Trichomonas vaginalis*.²

STEVE HIGGINS

Department of Genitourinary Medicine,
St Thomas's Hospital,
London SE1 7EH

- 1 Sumners D, Kelsey M, Chait I. Psychological aspects of lower urinary tract infections in women. *BMJ* 1992;304:17-9. (4 January.)
- 2 Brumfitt W, Hamilton-Miller JMT, Gillespie WA. The mysterious “urethral syndrome.” *BMJ* 1991;303:1-2.

SIR,—David Sumners and colleagues' paper on the psychological aspects of lower urinary tract infections in women would have been more informative if they had paid greater attention to detail.¹ In their comparison of women with and without urinary tract infections presenting to general practitioners one of the two significant differences that they were able to show between the two groups was in the prevalence of nocturia. Regrettably, they did not define nocturia. It is generally accepted that voiding once in the night is acceptable but twice or more times is regarded as abnormal. The other significant difference was in the dysuria score between the two groups. The

authors state that the difference in the means yielded a *t* statistic of 1.998, which is significant. Calculation of this statistic from their presented means and standard deviations, however, results in *t*=1.8, which is not significant.

The authors have assumed that patients presenting with dysfunction of the lower urinary tract who do not have a urinary tract infection have the urethral syndrome. This is obviously incorrect as they have ignored the possibility of various other abnormalities of the urinary tract, such as detrusor instability. The authors do not define the urethral syndrome but state that its aetiology has been the subject of considerable debate.² This is hardly surprising as the syndrome is thought to occur in young women as well as in postmenopausal women, and the treatments suggested for these two groups are completely different.³ It would be better to drop the term the urethral syndrome as it is not clearly defined and does not help clarify the therapeutic options for a problem that is already difficult to manage.

EBOO VERSI

Division of Obstetrics and Gynaecology,
St Thomas's Hospital,
London SE1 7EH

- 1 Sumners D, Kelsey M, Chait I. Psychological aspects of lower urinary tract infections in women. *BMJ* 1992;304:17-9. (4 January.)
- 2 Brumfitt W, Hamilton-Miller HMT, Gillespie WA. The mysterious “urethral syndrome.” *BMJ* 1991;303:1-2.
- 3 Versi E, Cardozo LD. Oestrogens and lower urinary tract function. In: Studd JWW, Whitehead MI, eds. *The menopause*. Oxford: Blackwell Scientific, 1988:76-84.

SIR,—Unfortunately, the data in David Sumners and colleagues' study of the psychological aspects of lower urinary tract infections in women do not support their conclusion that “patients with the urethral syndrome are no more neurotic than those with significant bacteriuria.”¹ Once again, the reason is that there are too few patients in the study groups.

Wisely, O'Dowd *et al* did not claim a significant difference between the eight out of 40 patients with the urethral syndrome who had received tranquillisers and the one out of 46 patients with urinary tract infection who had received tranquillisers in their study.² Statistical tests on groups with such small numbers can be misleading. If 10% of patients with urinary tract infection had neurosis and 20% of patients with the urethral syndrome had neurosis 307 patients in each group would be needed to reach significance (95% level with power 80%) (Epi-Info computer program).

I am at present analysing the results of a study to evaluate prospectively a method of diagnosing urinary tract infection based on a bayesian probability score which was described in 1987.³ Seven patients had received a psychoactive drug in the previous two years out of 66 patients with urinary tract infection, as compared with eight out of 60 patients with the urethral syndrome. This study is also too small to decide whether there really is a difference in prescribing of psychoactive drugs between the groups.

FRANK DOBBS

Drumcliffe Family Practice,
Drumcliffe,
County Sligo,
Irish Republic

- 1 Sumners D, Kelsey M, Chait I. Psychological aspects of lower urinary tract infections in women. *BMJ* 1992;304:17-9. (4 January.)
- 2 O'Dowd TC, Smail JE, West RR. Clinical judgment in the diagnosis and management of frequency and dysuria in general practice. *BMJ* 1984;288:1347-9.
- 3 Dobbs FF, Fleming DM. A simple scoring system for evaluating symptoms, history and urine dipstick testing in diagnosis of urinary tract infection. *J R Coll Gen Pract* 1987;37:100-4.

AUTHORS' REPLY,—We are pleased with the interest generated by our paper. We did not intend our study to provide diagnostic criteria but to

investigate the aetiology. We agree with A F Mellon and colleagues about the usefulness of urine microscopy as an aid to diagnosing infection, but we however advise caution when dealing with children,¹ patients with a self induced diuresis, and patients with vaginitis or cervicitis.²

Steve Higgins and Eboo Versi preach, wisely, that patients should have other causes of urethral symptoms investigated, and we do not disagree. We, however, took a pragmatic approach, realising that this was probably a low risk population³ and that a sexual history could be used as a screening instrument. The definition of the urethral syndrome was one of exclusion and was consistent with the Medical Research Council's definition.⁴ The finding that the urethral syndrome in our patients was negatively correlated with sexual activity reaffirms our belief, and that of others,⁵ that most cases of the urethral syndrome are not related to sexually acquired disease. With regard to nocturia, we defined it as voiding in excess of a patient's normal habit.

Frank Dobbs's comments on the power of the study are interesting. Regrettably, all research is a balance between what is feasible and what is desirable. Given the size of our sample, we did fail to disprove the null hypothesis that there was no difference in psychiatric disease between the two groups. We found that 17% (5/29) of the women with urinary tract infection and 25% (10/40) of those with the urethral syndrome had psychiatric disease. Had 17 (43%) patients with the urethral syndrome been psychiatrically ill this would have been significant at the 5% level ($\chi^2 = 3.84$, 1 df) and the sample size no larger. We had no previous work on which to judge our sample size and must leave it to others to confirm or refute our findings.

We thank Versi for pointing out an error in table I, which gives the mean (SD) dysuria score of patients with the urethral syndrome and urinary tract infection: the correct figures are 4.69 (2.1) and 5.27 (1.8), respectively. This difference is not significant.

MICHAEL C KELSEY

Whittington Hospital,
London N19 5NF

D SUMNERS

Napsbury Hospital,
London Colney,
Hertfordshire AL2 1AA

I CHAIT

Royal Free Hospital,
London NW3 2QG

- 1 Working Group of the Research Unit, Royal College of Physicians. Guidelines for the management of acute urinary tract infection in childhood. *J R Coll Physicians Lond* 1991; 25:36-42.
- 2 Kelsey MC, Fillin B. Gonorrhoea presenting as "sterile" pyuria. *BMJ* 1980;281:1009.
- 3 Kelsey MC, Mann GK, Bangham AM, Milnthorpe J. Non-specific (anaerobic) vaginitis: relevance of clinical and laboratory studies in a practice population. *J R Coll Gen Pract* 1987;37:56-8.
- 4 Members of the Medical Research Council Bacteriuria Committee. Recommended terminology of urinary-tract infection. *BMJ* 1979;iii:717-9.
- 5 Gillespie WA, Henderson EP, Linton KB, Smith PJB. Microbiology of the urethral (frequency and dysuria) syndrome. A controlled study with 5 year review. *Br J Urol* 1989;64:270-4.

Clinical trials in community pharmacies

SIR,—Steve Chaplin and Alison Blenkinsopp's editorial on clinical trials in community pharmacies states that great ignorance exists of the use, efficacy, and safety of over the counter medicines.¹ The profile of safety and efficacy has been well established by the time these drugs are licensed; that the medicine is available without prescription indicates that there were no major fears about safety. We need to learn more about use in the community, but it is incorrect to imply that inadequately researched drugs are available through pharmacies.

The authors state that pharmacy based trials conforming to the European Commission's guidelines for good clinical practice are possible. Compliance with some of the stipulations of the guidelines, however, would be problematic. For example, it is necessary to check case record forms with source documents, such as patients' notes.² This ensures that the record form accurately represents the facts. Unless pharmacists keep separate notes on individual patients such validation would be impossible.

The guidelines state that the investigator is medically responsible for subjects during the trial and must ensure that appropriate medical care is maintained subsequently. Clearly, this cannot apply if the investigator is not medically qualified.

According to the guidelines, investigators should immediately notify the sponsor, and possibly the ethics committee and authorities, of serious adverse events. These include death, life threatening events, and events resulting in admission to hospital, but the pharmacist might not be aware of these if the patient presented to his or her general practitioner or hospital. This affects the validity of any study.

The study that compared astemizole and terfenadine was unsatisfactory in that inappropriate conclusions were drawn. It was claimed that patients' assessments of control of symptoms showed comparability in the onset of action of the two drugs.³ It was subsequently pointed out that the study had not been performed in a way that made this inference possible.⁴ The first follow up data were obtained some 11 hours after dosing, which would have failed to detect the actual onset of action of the drugs (expected after about one hour with terfenadine).

Though we do not deny that community pharmacists could have a valuable role in clinical research, the process may not be straightforward and, for the present, the proposition should be viewed with caution.

H C MASHETER
E G BROWN

Marion Merrell Dow,
Uxbridge,
Middlesex UB11 1BE

- 1 Chaplin S, Blenkinsopp A. Clinical trials in community pharmacies. *BMJ* 1991;304:63-4. (11 January.)
- 2 Committee for Proprietary Medicinal Products Working Party on Efficacy of Medicinal Products. *Good clinical practice for trials on medicinal products in the European Community*. Brussels: Commission of the European Communities, 1988. (No 111/3976/88-EN Final.)
- 3 Quick JM, Blenkinsopp A. A community pharmacy study to compare the control of hay fever symptoms by astemizole 10 mg and terfenadine 120 mg. *Pharmaceutical Journal* 1991;246:272-5.
- 4 Masheter HC. Control of hay fever symptoms. *Pharmaceutical Journal* 1991;246:410.

Apnoea monitors and accidental strangulation

SIR,—J L Emery and colleagues draw attention to the risk of accidental strangulation associated with apnoea monitors, referring to an event that occurred when a Graseby MR10 monitor was used.¹

I wish to point out that every monitor leaves Graseby Medical's factory with an instruction leaflet. This leaflet explains where to position the sensor and that the tubing should be brought down the baby's legs, preferably underneath the clothing, so that accidental strangulation is not possible. Also, every respiration sensor comes in a pouch bearing an explanatory diagram.

As the company is aware that these issues cannot be emphasised enough it is currently reviewing the packaging of the sensor. Explanatory text will be added in six languages, as well as a different and clearer diagram. Thus every time a new sensor is used both the diagram and the instructions can be

referred to, giving optimal and instantaneous information.

When these sensors and monitors go to the community it is paramount that those who hand them out fully educate the users in their use. These moves should prevent further events similar to that reported by Emery and colleagues.

JASPERIEN MEINHARDT

Graseby Medical,
Watford,
Hertfordshire WD2 4LG

- 1 Emery JL, Taylor EM, Carpenter RG, Waite AJ. Apnoea monitors and accidental strangulation. *BMJ* 1992;304:117. (11 January.)

Not Anon

SIR,—In *The Penguin Book of Comic and Curious Verse*, published in 1952, J M Cohen, who had selected the verses, expressed his debt of gratitude¹ to W H Auden for supplying the verse "The Common Cormorant."^{2,4} Cohen noted in the foreword to the book that he had always suspected it to be a composition of Auden's "fathered by him on that broad shouldered old reprobate, Anon," whose authorship it was published under.¹

JIM MCCrackEN

Surgery on Rise Park,
Nottingham NG5 5EB

- 1 Cohen JM, ed. *The Penguin book of comic and curious verse*. Harmondsworth: Penguin, 1952:10.
- 2 Smith T. On lightning. *BMJ* 1991;303:1563. (21-28 December.)
- 3 Isherwood C. The common cormorant. In: Grigson G, ed. *Faber book of nonsense verse*. London: Faber, 1979:292.
- 4 Gordon R. Not Anon. *BMJ* 1992;304:257. (25 January.)

Junior doctors on the warpath

SIR,—Though I can understand the frustration of junior doctors that led to the call for a ballot on industrial action, I think that they will find that it was an ill judged decision which they will come to regret. Be that as it may, I cannot understand how the *BMJ*, which has such scrupulously high standards for publishing scientific articles, could accept Luisa Dillner's editorial discussing the issue,¹ which is inaccurate, misleading, and, I believe, irresponsible.

To say that "in 12 months little or nothing has changed" is flying in the face of the facts. In my own region of Wessex 201 posts were identified last year where junior doctors were working in excess of 83 hours a week in hard pressed specialties. A star chamber meeting between the regional task force and local working parties was held in November, and as a result of action after that meeting there are currently 80 posts that require in excess of 83 hours, but measures identified within each unit will reduce this number during 1992 to about seven. The regional task force will now be concentrating on those posts requiring over 73 hours.

To accuse some of the task forces of "incompetence" is both insulting and demoralising. Of course, some have been more effective than others. It is for this reason that a national meeting of the chairmen of regional task forces is to be held shortly so that best practice can be disseminated widely.

I suggest that the progress that those of us who were party to the agreement to reduce junior doctors' hours of work wish to see is more likely to follow regular factual reports than ill informed editorials.

A P J ROSS

Chairman, Joint Consultants Committee,
London WC1H 9JP

- 1 Dillner L. Junior doctors on the warpath. *BMJ* 1992;304:270. (1 February.)