that the road to the concentration camp, like the road to hell, is paved with good intentions.

DEREK MYERS

Brisbane,
Australia 4000

1 Widgery D. Desert storms. *BMJ* 1992;304:125. (11 January.)

FHSA medical advisers

SIR,—Tom Walley and John Bligh's editorial on the role of medical advisers to family health services authorities—of which I am one—seems paradoxical.¹ On the one hand we are described as the general practice equivalent of hospital clinical directors, which suggests a powerful managerial role. On the other hand we are described as bridge builders filling "the gap between primary care and the wider health service," which suggests passive mediators. Neither accurately defines the important nature of the role.

Both the main political parties agree that the future direction of development is for district health authorities and family health services authorities to merge into joint commissioning authorities. The medical adviser is the key to determining not only the priorities for developing the health service but, especially, the means by which to achieve them.

It is essential that the academic and planning skills of public health medicine are interlocked with the clinical and executive skills of medical advisers if we are incisively to influence the development of health services in primary care. In this way we can develop meaningful needs assessment in the community and provide better intelligence to the commissioning authority for use in purchasing. We will then be in a powerful position both to determine minimum standards and to ensure that consumers' interests are more objectively weighed alongside providers' priorities.

Currently, medical advisers are acting as agents of change but have neither the power nor the desire to direct clinically, as in a provider unit, the enormously varied range of work in general practice. But we will not be effective by merely filling gaps and building bridges. Forward looking family health services authorities are developing their capacity for research and development, led by medical advisers. This will enable them to break down the barrier between primary and secondary care by facilitating innovative practice that substitutes care in hospital for more appropriate and often cheaper care in the community. As purchasers we will have the power to redirect resources to develop services from the secondary sector in primary care. This seems to me to mean creating some gaps and burning some bridges.

ANDREW HARRIS

Lambeth, Southwark, and Lewisham Family
Health Services Authority,
London SE1 7NT

1 Walley T, Bligh J. FHSA medical advisers: friends or foes? *BMJ* 1992;304:133-4. (18 January.)

SIR,—Tom Walley and John Bligh's editorial discusses the role of medical advisers to family health services authorities.¹ I hope that family health services authorities and local medical committees will review this role, which should be to be a friend of general practice rather than a foe. These posts are provided for in regulation 33A, which states that "the FHSA shall appoint a doctor to assist it in the exercise of its functions pursuant to . . . enquiries about prescriptions and referrals." Many medical advisers now have a far wider remit than that, and there is often much more evidence of their activity in these other fields than in those for which they were originally appointed.

Family health services authorities are free to

seek medical advice from whom they choose, but not all medical advisers will have the necessary experience to provide advice on public health, community paediatrics, and audit. Advice on planning, resource allocation, and service development has traditionally come from local medical committees, and they must now ensure that they are not sidelined if family health services authorities obtain their advice from their own medical adviser. The value of advice from a local medical committee is that it is a distillate of the views of many general practitioners from a wide variety of different practices. The medical adviser may well have a much narrower experience and be less able to offer a balanced broad view.

Some medical advisers are active in medical audit advisory groups and have made valuable contributions to the establishment of audit and its support structures. Local medical committees should ask whether involving an employee of a family health services authority in audit is appropriate, given the profession's clear view that audit is not a "management" exercise.

Family health services authorities have a responsibility to consult widely and to take advice from representatives of the profession. This must include the local medical committees, which should ensure that communications with the authorities have been adapted to meet the requirements of the authorities' new responsibilities, particularly in relation to planning and monitoring the provision of general medical services. Local medical committees and individual general practitioners should establish relationships with their medical adviser and work with him or her, ensuring that the adviser is well acquainted with developments in general practice and with its needs and aspirations. Medical advisers must show that they can be of benefit to individual general practitioners and to general practice as a whole in their locality. If they are to assume responsibilities beyond the monitoring of prescribing habits and referral patterns they must develop close links with local medical committees, where they will have every opportunity to show that they are indeed friends of general practice.

D E PICKERSGILL

Birchwood Surgery,
North Walsham,
Norfolk NR28 0BQ

1 Walley T, Bligh J. FHSA medical advisers: friends or foes? *BMJ* 1992;304:113-4. (18 January.)

Distinction awards

SIR,—I have recently seen the Advisory Committee on Distinction Awards' analysis of distinction awards as at 31 December 1990.¹ Yet again I am struck by the sheer number of awards—to nearly 36% of all eligible practitioners.

Apart from the impossibility of distinguishing merit at the level of the C award fairly, surely awarding selective supplementary income so widely does nothing to promote adequate remuneration for the profession as a whole. Is it not time to scrap this anachronism?

M ASHLEY-MILLER

Nuffield Provincial Hospitals Trust,
London NW1 7SP

1 Department of Health. Distinction awards: analysis by type of award, specialty and percentage distribution at 31st December 1990—England and Wales. *Health Trends* 1991;23:127.

Vincent van Gogh's illness

SIR,—It is a fundamental truth in clinical medicine that common illnesses occur more commonly, and we think that this admonition against diagnosing

rare medical disorders when a common disease fits the course and symptoms of an illness equally well or better is particularly apt in the case of Vincent van Gogh.¹

The prevalence of manic-depressive illness is 1%; that of acute intermittent porphyria is less than 0.008%. When the classic course and symptoms of manic-depressive illness are present its diagnosis is extremely reliable.² Vincent van Gogh's premorbid personality, cyclic pattern of affective episodes, lucidity between attacks, absence of cognitive deterioration, seasonal exacerbations, bursts of productivity, altered sleep and energy patterns, hyperreligiosity, volatile irritability and violence, visual and auditory hallucinations, and alcohol misuse (also a common cause of gastrointestinal complaints) are the hallmarks of manic-depressive illness. His convulsive disorder may have been precipitated by misuse of absinthe or due to a coexisting complex partial seizure disorder.

Vincent's affective symptoms clearly predated any history of seizures, and his family history is more consistent with manic-depressive illness. His brother Theo also suffered from an affective disorder, and their sister, Wilhelmina, spent many years in a psychiatric institution. Vincent and his youngest brother, Cor, both killed themselves, an outcome that they share with 15-20% of people with untreated manic-depression³; moreover, 70-80% of all suicides are associated with affective disorder.² Suicide in acute intermittent porphyria, on the other hand, is rare. Clustering of affective illness and suicide is common in families of people with manic-depression³; in contrast, only one first degree relative of 13 probands with acute intermittent porphyria had any form of mental illness (it was not manic-depressive illness).⁴ Finally, manic-depressive illness is far more common in highly creative artists and writers than in the general population^{2,5}; acute intermittent porphyria shows no such relation. In fact, acute intermittent porphyria has been associated with neuropsychiatric impairment and extremely poor manual dexterity.⁶

It is incorrect to dismiss manic-depressive illness because of the "facile reversibility and relatively short duration of Vincent's crises." Short duration and reversibility are perfectly consistent with manic-depressive illness, particularly early in its course.²

KAY REDFIELD JAMISON

Johns Hopkins University School of Medicine,
Baltimore,
Maryland,
USA

RICHARD JED WYATT

Neuropsychiatry Branch,
National Institute of Mental Health,
Washington, DC 20032,
USA

- 1 Loftus L, Arnold W. Vincent van Gogh's illness: acute intermittent porphyria? *BMJ* 1991;303:1589-91. (21-28 December.)
- 2 Goodwin FK, Jamison KR. *Manic-depressive illness*. New York: Oxford University Press, 1990.
- 3 Egeland JA, Sussex JN. Suicide and family loading for affective disorder. *JAMA* 1985;254:915-8.
- 4 Ackner B, Cooper JE, Gray CH, Kelly M. Acute porphyria: a neuropsychiatric biochemical study. *J Psychosom Res* 1962;6:1-24.
- 5 Jamison KR. *Touched with fire: manic-depressive illness and the artistic temperament*. New York: Free Press, 1992.
- 6 Tishler PV, Woodward B, O'Connor J, Holbrook DA, Seidman LJ, Hallet M, et al. High prevalence of intermittent acute porphyria in a psychiatric patient population. *Am J Psychiatry* 1985;142:1430-6.

Correction

Deaths from haemolytic disease of the newborn in 1990

A printer's error occurred in this letter by Ruth Hussey and Cyril Clarke (15 February, p 444). Throughout the letter anti-C should have been printed as anti-c.