and the Department of Health with accident compensation statistics is clearly necessary if adequate data are to be used in preventing accidents.

PATRICK J MORRISON

Department of Medical Genetics,
Queen's University of Belfast,
Belfast City Hospital,
Belfast BT9 7AB

- 1 Teanby D. Underreporting of pedestrian road accidents. *BMJ* 1992;304:422. (15 February.)
- 2 Morrison PJ, Kjellstrom T. A comparison of hospital admissions data and official government statistics of serious traffic accident injuries. *NZ Med J* 1987;100:517-20.

Home visits by psychiatrists

SIR,—Christine Hardy-Thompson and colleagues report general practitioners' views on psycho-geriatric domiciliary consultations.¹ We recently submitted for publication a report of a study evaluating the cost of psychiatric outpatient consultations and what the costs of home consultations would be if all such outpatients were seen at home. The cost of psychiatric outpatient consultations in England for 1989 was estimated as £205 882 529 and that for home consultations as £215 326 723. The main reason for the costs being similar despite the extra time spent travelling for home visits was the high non-attendance rate (up to 38%²) in the outpatient clinic.

Assessing patients at home has several advantages. The consultation rate is almost 100%. The patient's home environment can be observed at first hand rather than information on it obtained from the patient or an informant at the clinic. Unnecessary social work and occupational therapy assessments, which may otherwise be requested by the outpatient clinic are avoided. The psychiatrist has ready access to the patient's medication, which is often not brought to the clinic. This draws attention to polypharmacy, poor compliance, possible interaction with non-psychotropic drugs, hoarding for overdose, and dependence on benzodiazepines. The stigma of attending a psychiatric clinic is reduced.

Given these advantages and that home consultations are only marginally more expensive than outpatient consultations, we should be looking at placing greater emphasis on home consultations. Some psycho-geriatric services have almost done away with outpatient clinics in favour of home visits. This to some extent will reduce the importance of the indications for domiciliary consultations.

AJIT SHAH

Section of Epidemiology and General Practice,
Institute of Psychiatry,
London SE5 8AF

- 1 Hardy-Thompson C, Orrell MW, Bergmann K. Evaluating a psycho-geriatric domiciliary visit service: views of general practitioners. *BMJ* 1992;304:421-2. (15 February.)
- 2 Shah AK, Lynch S. The characteristics of patients attending a psychiatric follow-up clinic. *Psychiatric Bulletin* 1990;14:153-4.

Fundamental principles of the NHS

SIR,—A consensus is now emerging that general practitioners can be effective commissioners of secondary care, and there is concern as to how the commissioning role can be efficiently extended to the whole of general practice. The fact is that fundholders in general practice, by redefining their relationship with their consultant colleagues, exploiting the flexibility that their small volumes of work allow them, and combining their knowledge of patients' needs with their knowledge of pro-

viders' capabilities, can be more effective commissioners of care than health authorities.

With further expansion of the fundholding scheme several serious issues need considering. How can contractors cope with multiple purchasers? Is the increased management cost justified? How will family health services authorities and regional health authorities effectively monitor an expanded scheme? Will all practices perform efficiently as commissioners of care? What will the effect be on district health authorities and purchasing consortiums? How can general practitioners' purchasing plans be integrated within the strategic framework developed by health authorities? What is the role of the general practitioner to be?

These questions are complex, and the profession needs to develop answers as, no matter what the political environment, the issues will remain. Fundholding has shown that general practitioners, acting as the patients' agent, can deliver an improved service. How can this be spread to the rest of our colleagues?

The stark choice facing us is between rationing by restricting the range of services provided by the NHS or improving the efficiency of the service, as J B L Howell points out.¹ We should all consider how those efficiency gains can be achieved.

N H L STAREY

Burnham Surgery,
Burnham on Crouch,
Essex CM0 8SJ

- 1 Howell JBL. Re-examining the fundamental principles of the NHS. *BMJ* 1992;304:297-9. (1 February.)

SIR,—Disconcertingly, in J B L Howell's article¹ the duties of the minister of health specified in the NHS Act of 1946 tailed off into a row of dots just before the word "rehabilitation" occurred in the original. Sir Donald Acheson has recently reminded us that Beveridge's three legged stool of the NHS specified prevention, treatment, and rehabilitation. It is surely a fundamental principle of three legged stools that the shorter one leg gets in comparison with the others, the more uncomfortable the stool becomes. There is a great deal of catching up to be done in the development of our rehabilitation services before we can claim to be back in alignment with the fundamental principles of the NHS.

D L McLELLAN

Rehabilitation Research Unit,
University of Southampton,
Southampton SO9 4XY

- 1 Howell JBL. Re-examining the fundamental principles of the NHS. *BMJ* 1992;304:277-9. (1 February.)

Prescribing at the hospital-general practice interface

SIR,—My recent research leads me to concur with Roger Jones and Michael Rawlins that communication is singularly important for quality of care at the interface between hospitals and general practitioners.¹

I conducted qualitative research, interviewing 21 general practitioners from nine local non-fundholding practices. My objective was to identify the real concerns of general practitioners with regard to prescribing issues at the interface. My findings in order of importance as perceived by general practitioners are as follows.

Firstly, the promptness and quality of communication from hospitals to general practitioners about prescribing was criticised. Though patients received seven days' supply of drugs on discharge, they often attended the surgery well within this time to request continuation of treatment. Thus in many cases the general practitioner was being

asked to prescribe without adequate information. The clarity and amount of information were said to need attention. General practitioners wanted to know the intended duration of treatment and the reasons for interventions.

Secondly, the interviewees were concerned that clinical responsibility was unclear. Views varied widely, including, "The hospital should provide the general practitioner with information to prescribe complex drug regimens" and, "The hospital must share clinical responsibility if the general practitioner prescribes for the hospital."

Thirdly, there was a general feeling that it was unfair to levy a prescription charge for two weeks' treatment. Some "more experienced" patients go direct from the outpatient department to the general practitioner, knowing that they may obtain one month's supply.

Fourthly, loss leaders to the hospital service were frequently mentioned, though this was considered comparatively less contentious.

Fifthly, the general practitioners often mentioned that an intervention in treatment while the patient was in hospital had not been explained to them.

Finally, some of the general practitioners considered that the system of returning patients' own drugs, taken into hospital for a drug history, was not working adequately, leading to potential wastage.

These concerns are now being addressed locally; I believe that general practitioners in other areas may have similar concerns. I thus recommend that those concerned in compiling hospital drug protocols should involve local general practitioners both through liaison committees and by a direct personal approach. Patients should be informed of the communication process and its timescale.

I believe that the recent NHS reforms and an executive letter^{2,3} now give us the opportunity to improve the quality of care across the interface between hospitals and general practitioners. There is no need to wait for any merging of district health authorities and family health services authorities or changes allowing virement between budgets before starting initiatives.

R DAVIES

Wrexham Maelor Hospital,
Wrexham,
Clwyd LL13 7TD

- 1 Jones R, Rawlins MD. Prescribing at the interface between hospitals and general practitioners. *BMJ* 1992;304:4-5. (4 January.)
- 2 NHS Management Executive. *Responsibility for prescribing between hospitals and GPs*. London: Department of Health, 1991. (EL(91)127.)
- 3 Welsh Office. *Responsibility for prescribing between hospitals and GPs*. Cardiff: Welsh Office, 1991. (WHC(91)94.)

One car down

SIR,—R S Bhopal's article on car ownership was thoughtful and provocative,¹ but its focus was on what individuals can do by and for themselves. Another set of questions must be asked about what health authorities, NHS trusts, and directly managed provider units could do to discourage unnecessary car use and so contribute to a greener NHS.

Possible ways to discourage use of cars include (1) increasing cycle mileage rates to near those for cars. This would encourage staff to use cycles for work by allowing them to make a small profit. (2) Giving free on site car parking spaces to staff (regardless of status) only if they need to use their car during working time. Many hospitals automatically give consultants free car parking spaces when they need their car only for travel to and from work. (3) Where on site car parking is charged for, to do so on the basis of engine size. (4) Providing secure cycle racks at the expense of car parking spaces. (5) Giving loans for public