sible for the condition) is likely. Indeed Levitt *et al* reported on three North American families with malignant hyperthermia in which linkage studies excluded the locus for the malignant hyperthermia gene from the 19q 13.1 region, where the gene for the ryanodine receptor is situated.¹² Similarly, the European Malignant Hyperthermia Group has found that malignant hyperthermia does not cosegregate with a gene in the 19q 13.1 region.¹³

Even in families in which the gene for the ryanodine receptor may be implicated the actual mutation may vary in different family members. Although Fujii *et al* found that a single point mutation in the porcine ryanodine receptor gene was associated with porcine malignant hyperthermia in five major breeds,¹⁴ a similar mutation has been discovered in only one of 35 human pedigrees.¹⁵

Although the DNA research provides an exciting new dimension to laboratory study of malignant hyperthermia, the importance of the clinical condition should not be forgotten. If a DNA based test for malignant hyperthermia became available family members would be more easily screened than at present with the in vitro muscle contracture test. One of the problems, however, is to identify and locate members of extended families. From our own work we know of two British families in which a second proband was discovered, and both died of malignant hyperthermia. Had the anaesthetist known that a family member had been recognised as being susceptible to malignant hyperthermia the trigger agent for the condition would not have been administered. The reason for these tragedies was not the lack of a suitable screening test but lack of communication between family members.

Currently, detection of clinical episodes of malignant hyperthermia will remain largely the responsibility of the anaesthetist in the operating theatre. But detecting susceptibility to malignant hyperthermia in members of known malignant hyperthermia pedigrees is also essential to avoid further anaesthetic tragedies.

F RICHARD ELLIS

Reader in Anaesthesia, St James's University Hospital, Leeds LS9 7TF

- 1 MacLennan DH, Duff C, Zorato F, Fujii J, Phillips M, Korneluk RG, *et al*. Ryanodine receptor gene is a candidate for predisposition to malignant hyperthermia. *Nature* 1990;343:559-61.
- 2 McCarthy TV, Healy S, Heffron JJA, Lehane M, Deufel T, Lehmann-Horn F, *et al*. Localisation of malignant hyperthermia susceptibility locus to human chromosome 19q12-13.2. *Nature* 1990;343:562-4.
- 3 Ellis FR, Heffron JJA. Clinical and biochemical aspects of malignant hyperthermia. *Recent Advances in Anaesthesia and Analgesia* 1985;15:173-207.
- 4 Kalow W, Britt BA, Terreau ME, Haist C. Metabolic error of muscle metabolism after recovery from malignant hyperthermia. *Lancet* 1970;ii:895-8.
- 5 Ellis FR, Harriman DGF, Keaney NP, Kyei-Mensah K, Tyrrell JH. Halothane-induced muscle contracture as a cause of hyperpyrexia. *Br J Anaesth* 1971;43:721-2.
- 6 European Malignant Hyperpyrexia Group. A protocol for the investigation of malignant hyperpyrexia (MH) susceptibility. *Br J Anaesth* 1984;56:1267-9.
- 7 Larach MG (for the North American Malignant Hyperthermia Group). Standardization of the caffeine halothane muscle contracture test. *Anesth Analg* 1989;69:511-5.
- 8 Renwick DS, Chandraker A, Bannister P. Missed neuroleptic malignant syndrome. *BMJ* 1992;304:831-2.
- 9 Adnet PJ, Krivosic-Horber RM, Adamantidis MM, Handecouer G, Adnet-Bonte CA, Saulnier F, *et al*. The association between the neuroleptic malignant syndrome and malignant hyperthermia. *Acta Anaesthesiol Scand* 1989;33:676-80.
- 10 Campkin NTA, Davies UM. Another death from Ecstasy. *J R Soc Med* 1992;85:61.
- 11 Healy JMS, Heffron JJA, Lehane M, Bradley DG, Johnson K, McCarthy TV. Diagnosis of susceptibility to malignant hyperthermia with flanking DNA markers. *BMJ* 1991;303:1225-8.
- 12 Levitt RC, Nouri N, Jedlicka AE, McKuride VA, Marks AR, Shutack JG, *et al*. Evidence for genetic heterogeneity in malignant hyperthermia susceptibility. *Genomics* 1991;11:543-7.
- 13 Deufel T, Meitinger T, Golla A, Johnson K, MacLennan DH, Lehmann-Horn F. Two recombinations between malignant hyperthermia susceptibility (MHS) and human ryanodine receptor (RYR1) in a single family: Another gene for MHS? *Cytogenet Cell Genet* 1992 (in press).
- 14 Fujii J, Otsu K, Zorato F, de Leon S, Khanna VK, Weiler JE, *et al*. Identification of a mutation in porcine ryanodine receptor associated with malignant hyperthermia. *Science* 1991;253:448-51.
- 15 Gillard EF, Otsu K, Fujii J, Khanna VK, de Leon S, Derdemezi J, *et al*. A substitution of cysteine for arginine 614 in the ryanodine receptor is potentially causative of human malignant hyperthermia. *Genomics* 1991;11:751-5.

The end of scientific journals?

No longer so far away

Francis Fukuyama's book *The End of History and the Last Man* has excited much debate.¹ He argues disturbingly that history is fizzling out into a world where there are no alternatives to liberal democracy and free markets and where the last man may want nothing more than to be left alone. Even more disturbing for the scientific, technical, and medical publishers who gathered in force earlier this month in Amsterdam is the thought that the last scientist may not even be reading their journals. Rather he will access everything through his computer, bringing to an end 350 years of paper journals.

Publishers have, of course, heard this before. The paperless society has been predicted for 30 years, and publishers who have grown older and richer during this time are understandably cynical about further predictions of their end. But it may now be close. The United States Office of Technology Assessment has predicted that by the end of the century research scientists will use their computers to gather information from each other through electronic mail and bulletin boards and from supercomputers, data archives, digital libraries, and electronic journals.

The Amsterdam conference was different from its many predecessors on the end of journals in that "the dogs who are going to eat the publishers' dinner" were there. They are people who are offering what have been called (by David Brown, a consultant who spoke at the conference) CASIAS systems—Current Awareness Services and Individual Article Supply. These services communicate with users through

computer networks and tell them about new relevant articles. They then aim at supplying users with copies of articles they want within an hour. No company has a full system ready yet, but out in front is an organisation called CARL (Colorado Alliance of Research Libraries). It has a network that spans the United States with about 5000 terminals that the public can access. It covers some 12 000 different journals and has two million articles in its database with another 600 000 being added annually. It can supply articles by fax within 24 hours and can supply within an hour any articles that have been electrocopied (that is, scanned and stored electronically). Users pay a set fee of \$6.50 for each article supplied plus a copyright fee to the publisher. Several other organisations are about to begin such services, and many libraries are also beginning to offer their users such a service.

What terrifies the publishers is that many libraries might stop subscribing to journals and subscribe to CASIAS organisations instead. This might lead to the disappearance of many journals unless they could keep themselves solvent by selling individual articles through the CASIAS systems. The problem is that survival might depend on charging a great deal more than the \$10-15 that it is thought the average reader will be willing to pay for each article.

These possibilities are opening up now because economic, technological, and cultural changes have come together, said Maurice Long, international sales manager of the BMJ Publications Group, at the conference. Libraries around the world are suffering severe financial cuts and as a consequence

are having to reduce their journals lists. Thus a sample of 10 libraries in the United States have cut 175 journals each a year for the past three years but calculate that they will have to cut over 500 each in 1992. Then, culturally, librarians have stopped thinking of themselves as archivists and come to see themselves as information scientists providing readers with the information they need as quickly as possible; in addition, their users—increasingly members of the “Nintendo generation”—are willing to find their information through a computer. Finally, technology has developed to make CASIAS systems easy to run. In particular, computer networks have increased rapidly (from 600 worldwide in 1988 to 3000 now) and fax and optical character readers have become cheap and widely available.

Another factor driving the rapid change—but not mentioned much at the publishers’ conference—is the realisation that many journals in a library are consulted only once or twice a year.² When subscriptions may be \$500 a year it is obviously much cheaper to get that one article faxed through at perhaps \$15. These findings by librarians fit with the increasing realisation by information scientists that only a few articles in most journals are scientifically sound and of direct use to most readers.^{3,4}

This development in CASIAS systems comes at a time when electronic journals are also appearing.^{5,6} These journals should have various advantages over their paper counterparts: potentially faster peer review systems because of electronic transmission of manuscripts backwards and forwards; immediate release once the manuscript has been accepted; the

potential to include many more data than is possible in a paper journal; the possibility of attaching structured abstracts or even the full papers of references; and the chance to make corrections on the stored article.

The appearance of these alternatives to paper journals is driving publishers and editors to think about just what value they do add to the process of disseminating science and whether they are needed. Peer review is one added ingredient, but editors need to be sure that they do it speedily and well. Technical editing is also important (as readers of unedited manuscripts will testify), but the most important added value may be to group together different sorts of material—not only original science but also comment, reviews, news, correspondence, and articles—in a way that makes a journal satisfying enough to read in bed.

The final result of these rapid changes in how science is disseminated might thus be that general journals will remain in paper form, whereas specialist journals will eventually cease and instead individual articles will be available electronically. But none of us should be complacent: the Romans assumed that their empire would last forever.

RICHARD SMITH

Editor, BMJ

- 1 Fukuyama F. *The end of history and the last man?* London: Hamish Hamilton, 1992.
- 2 McSean A. Managing periodicals subscriptions: improving cost effectiveness. *Serials* (in press).
- 3 Williamson JW, Goldschmidt PG, Colton T. The quality of medical literature: an analysis of validation assessments. In: Bailar JC, Mosteller F. *Medical use of statistics*. Waltham, Mass: NEJM Books, 1986.
- 4 Lock S. Introduction. In: Lock S, ed. *The future of medical journals*. London: BMJ, 1991.
- 5 Smith J. New electronic journal aims to beat NEJM. *BMJ* 1991;303:945.
- 6 Kassirer JP. Journals in bits and bytes. *N Engl J Med* 1992;326:195-6.

Vertebral fractures

How large is the silent epidemic?

The ascendancy of osteoporosis to its present status as a leading health problem has led, finally, to the critical study of vertebral fractures. Although these have long been attributed to osteoporosis, only recently have epidemiological data confirmed that low bone mineral density is their most important determinant.^{1,2} Large prospective studies have suggested that the risk of sustaining a new vertebral fracture more than doubles with each decrease of one standard deviation in bone mineral density of the lumbar spine.^{3,4}

Falls, so prominent in the pathogenesis of fractures of the hip and distal forearm, do not play such a large part in causing vertebral fractures because the spine is subjected to substantial loads during daily activities such as bending forward, lifting objects, and climbing stairs; vertebral fractures result uniquely from such loading.^{5,6} Despite the strength of the association between bone mineral density and vertebral fracture the overlap between bone density in patients with vertebral fractures and that in control subjects without fractures is sufficiently large for vertebral fracture to be diagnosed reliably only from radiographs. Densitometry is more appropriately used to assess the future risk of fracture.⁷

Assessment of the impact of vertebral fractures has been hampered by the absence of formal criteria for identifying fractures in radiographs of the thoracolumbar spine. Even in early case series three patterns were recognised: wedge, crush, and end plate (biconcave) fractures. Since then the means of defining vertebral fractures have evolved through several stages. Initial methods, relying on subjective radiological assessment,⁸ gave way to morphometric measurement

of vertebral height with fractures defined according to fixed cut off values.⁹ As each vertebral body in the spinal column has unique dimensions¹⁰ recent analyses have focused on deriving the distribution of vertebral dimensions at each spinal level and calculating cut off values from these.^{11,12}

The most widely adopted thresholds for defining and grading fractures denote moderate (or grade 1) fractures as deformities that fall between three and four standard deviations from the mean values specific to each vertebra, and severe (or grade 2) fractures as those that fall four standard deviations or more from this mean.¹¹ When morphometric studies are done without reference to clinical presentation the abnormalities found are usually referred to as deformities rather than fractures.

The application of recently developed morphometric techniques to various population samples in the United States has permitted estimation of the incidence of new vertebral fractures in the general population. One recent estimate of the age adjusted incidence among white American women aged 50 and over was 15.1 per 1000 person years. This is more than twice the incidence of hip fracture (6.2 per 1000 person years).

It has long been clear, however, that some vertebral fractures do not reach clinical attention, although the size of this fraction was unknown.¹³ Recently, the age adjusted incidence of clinically ascertained vertebral fractures was estimated at 5.3 per 1000 person years among white American women aged 50 and over, or 35% of the total figure.¹⁴ The incidence of clinically diagnosed vertebral fractures in this