and that we have as much to learn from Europe as vice versa.

The Hospital Doctors' Association has campaigned for two years for a single training grade for all branches of specialist medicine, in which the trainee would spend a shorter time and be more closely supervised than at present. This idea is now being examined by the Royal College of Obstetricians and Gynaecologists and piloted by the Royal College of Surgeons. I believe that Brearley is mistaken in seeking to extend the numbering system currently operated by the Royal College of Surgeons. It cannot be right to have two people in essentially the same job on the same rota, with one preselected for promotion and accreditation and the other denied this. Such a system will eventually fall foul of the courts in the United Kingdom or Brussels in the same way that the practices of the Joint Committee on Higher Medical Training have done already.

We must stop separating the issues of manpower, training, accreditation, and hours of work. They are all intimately linked. We need wholehearted reform of our practices of work and training, and the Hospital Doctors' Association has called for the royal colleges, the General Medical Council, the Department of Health, and the representative associations of the profession to meet and take up the challenge.

DAVID WREDE

Chairman, Hospital Doctors' Association,
Ascot,
Berkshire SL5 7EN

1 Brearley S. Accreditation after Goldstein. *BMJ* 1992;304:518-9. (29 February.)

Visiting registrars

SIR,—I congratulate Stephen Brearley on his editorial on accreditation.¹ A system riddled with anomalies was bound to have its days numbered. He goes on to say, however, that accreditation is a purely British phenomenon and designed to fulfil a uniquely British need. Does he mean the use of overseas doctors (alias the politically correct "visiting registrar") in district general hospitals, where the bulk of NHS work is done?

He recommends using visiting registrars for unrecognised senior registrar posts. Is this because they lack the necessary intelligence or because they do not deserve better? Once more the visiting registrar becomes the stop gap for unrecognised locum senior registrar posts. Justice is long overdue here too.

S BORGES

Caithness General Hospital,
Wick KW1 5LA

1 Brearley S. Accreditation after Goldstein. *BMJ* 1992;304:518-9. (29 February.)

Renal protective effect of enalapril in diabetic nephropathy

SIR,—We feel that Staffan Björck and colleagues¹ are not justified in concluding that enalapril reduces the rate of decline in kidney function in diabetic nephropathy more than metoprolol, given a similar antihypertensive effect. The accompanying editorial by Carl Erik Mogensen² points to some disagreement between their results and those of similar studies.^{3,4} Their results therefore need detailed scrutiny.

The question at issue is whether Björck *et al* did in fact achieve an equivalent antihypertensive effect with the enalapril and metoprolol based regimens. They emphasised the similarity in mean arterial pressure but ignored important differences in supine blood pressure. In absolute terms supine

diastolic pressure was 6 mm Hg higher ($p<0.005$) with metoprolol, and the antihypertensive effect from baseline values was substantially smaller (11/1 mm Hg metoprolol; 17/12 mm Hg enalapril). One might argue about the relative importance of diastolic, systolic, or mean arterial pressure, but what is clear is that a mean difference in diastolic pressure of 6 mm Hg cannot be dismissed. For example, such a difference is associated with a 40% reduction in the incidence of stroke in intervention and epidemiological studies.⁵ It is quite conceivable that the blood pressure differences observed could be responsible for the apparent benefit of enalapril on renal function.

There is no doubt that angiotensin converting enzyme inhibitors reduce proteinuria, a response observed also in non-diabetic nephropathy even when renal function tends to decline more rapidly than with other treatment.⁶ However, the association between proteinuria and progression to end stage renal failure hinges predominantly on the relation observed in untreated patients,⁷ which may not hold in the treated state. The hypothesis that angiotensin converting enzyme inhibitors are renoprotective is likely to be answered convincingly only by much larger studies than that of Björck *et al*, and preferably by studies measuring real rather than surrogate endpoints.

W W YEO

L E RAMSAY

P R JACKSON

Sheffield Hypertension Clinic,
University Department of Medicine and Pharmacology,
Royal Hallamshire Hospital,
Sheffield S10 2JF

- 1 Björck S, Mulec H, Johnsen SA, Norden G, Aurell M. Renal protective effects of enalapril in diabetic nephropathy. *BMJ* 1992;304:339-43. (8 February.)
- 2 Mogensen CE. Angiotensin converting enzyme inhibitors and diabetic nephropathy. *BMJ* 1992;304:327-8. (8 February.)
- 3 Parving HH, Smidt UM, Mathiesen ER, Hommel E. Ten years' experience with antihypertensive treatment in diabetic nephropathy. *Diabetologia* 1991;34 (suppl 2):A38.
- 4 Gronhagen-Riska C, Honkanen E, Metsarinne K, Rosenlof K, Tikkanen I, Fyhrquist F. ACE-inhibition versus conventional antihypertensive treatment (beta-blockade) in diabetic nephropathy. *Am J Hypertens* 1990;3:67A.
- 5 Collins R, Peto R, MacMahon S, Herbert P, Fiebach NH, Eberlein KA, *et al*. Blood pressure, stroke, and coronary heart disease. 2. Short-term reductions in blood pressure: overview of randomised drug trials in their epidemiological context. *Lancet* 1990;335:827-38.
- 6 Apperloo AJ, de Zeeuk D, Sluiter HG, de Jong PE. Differential effects of enalapril and atenolol on proteinuria and renal haemodynamics in non-diabetic renal disease. *BMJ* 1991;303:821-4.
- 7 Williams PS, Fass G, Bone JM. Renal pathology and proteinuria determine progression in untreated mild/moderate chronic renal failure. *Q J Med* 1988;252:343-54.

SIR,—Staffan Björck and colleagues claim to have shown a specific beneficial effect, independent of blood pressure, of enalapril on the decline of glomerular filtration rate in diabetic nephropathy.¹ We would like to draw attention to several problems with this study that lead us to question the validity of the conclusions.

In both the interim report of their study and in the final paper Björck *et al* reported significantly lower supine diastolic blood pressure values in the enalapril group than in the metoprolol group.^{1,2} Since even small differences in blood pressure above the normotensive range may have a substantial effect on the progression of diabetic nephropathy³ the effect of enalapril cannot be attributed to an action of the drug independent of blood pressure.

Differences in the decline of glomerular filtration rate between the groups were only significant when the measurements of the rate before randomisation of patients were included in the evaluation; differences were not significantly different during the drug treatment period (table III).¹

Deterioration of renal function over time was analysed by linear regression. This method of analysis can be used only if the decline of renal function is linear—but this was not the case in the

enalapril group. After six months of treatment glomerular filtration rate deteriorated significantly in only the enalapril group, as stated in the interim report and in the final report (fig 3).¹ Since the decline of glomerular filtration rate in the enalapril group was not linear no linear regression analysis can be calculated for this group. A transient drop in glomerular filtration rate when enalapril was started may have artificially "reduced" the slope of the decline of glomerular filtration rate in a linear regression analysis. Unfortunately the authors do not give the final values for the glomerular filtration rate in each group and they do not state whether the rate was different between the groups at the end of the study. Most probably it was not.

Patients receiving drugs whose renal function deteriorated rapidly should have been excluded in both treatment groups (patient No 23) and not only in the enalapril group (patient No 21).

Patients with a follow up of only six months should have been excluded as this follow up period allowed glomerular filtration rate to be measured only once during drug treatment. The lack of further measurements renders any regression analysis of the decline of glomerular filtration rate under the influence of the investigated drugs impossible.

In conclusion, the study does not provide convincing evidence for a specific beneficial effect of enalapril on the decline of glomerular filtration rate in diabetic nephropathy.

PETER T SAWICKI

MICHAEL BERGER

Department of Nutrition and Metabolism,
Heinrich-Heine University,
W-4000 Düsseldorf 1, Germany

- 1 Björck S, Mulec H, Johnsen SA, Norden G, Aurell M. Renal protective effect of enalapril in diabetic nephropathy. *BMJ* 1992;304:339-42.
- 2 Björck S, Mulec H, Johnsen SA, Nyberg G, Aurell M. Contrasting effects of enalapril and metoprolol on proteinuria in diabetic nephropathy. *BMJ* 1990;300:904-7.
- 3 Sawicki PT, Didjurgeit U, Mühlhauser I, Heinemann L, Berger M. Near normotension and near normoglycaemia in blind type 1 diabetic patients with overt diabetic nephropathy. *J Diabetic Complications* 1990;4:179-83.

AUTHOR'S REPLY,—It was a new finding in our study that the enalapril group tended to have a lower blood pressure than the metoprolol group in the supine position and that the reverse occurred in the standing position. This phenomenon cannot be handled by an increased sample size. The mean of the supine and the standing blood pressure is the only measure that can be used for the evaluation because of this unexpected difference in antihypertensive action. If we had used more aggressive antihypertensive treatment to reduce supine blood pressure more in the metoprolol treated patients the two groups certainly would not have been comparable, with a much lower standing blood pressure in the metoprolol treated patients. The difference of 1 mm Hg that we obtained in mean arterial blood pressure between the two groups is very small.

Even without a control group the result of enalapril treatment can be seen to be very good and is the best reported in this type of patient. After the initial half year the fall in glomerular filtration rate was only 0.4 (6.9) ml/min/year. For the six patients followed for three years the corresponding figure was 0.4 (1.4) ml/min/year, which is remarkable in patients with diabetic nephropathy and a 50% reduction in renal function. The figure is the same as the normal, age dependent fall in glomerular filtration rate.¹

Another conclusion from our data is that studies of angiotensin converting enzyme inhibitors should focus on the renal effects after renal function has been allowed to stabilise for some months. Therefore, even though ours is one of the larger prospective studies in diabetic nephropathy, new studies comparing other types of antihypertensive

drugs should aim at an even larger number of patients and continue longer. The planning of this study started in 1985 after our first results¹ were available; at this time there were no human data suggesting a renal protective effect of angiotensin converting enzyme inhibitors.

Peter T Sawicki and Michael Berger seem to recommend a comparison of enalapril and metoprolol treatments during the follow up period from six months onwards, and that only patients with follow up times of one year or more be included. A total of 18 patients treated with enalapril and 13 treated with metoprolol are evaluable in this respect. During this period the mean (SD) rate of change in glomerular filtration rate was 0.8 (4.7) ml/min/year in the enalapril treated and -3.3 (4.2) ml/min/year in the metoprolol treated patients ($p=0.016$). The mean arterial blood pressures (mean of supine and standing pressures) were 102 (5) mm Hg and 103 (4) mm Hg, respectively. Multiple regression analysis shows that enalapril or metoprolol treatment was correlated to the rate of fall in glomerular filtration rate ($p=0.02$) but mean arterial blood pressure was not ($p=0.8$). If supine or standing mean arterial blood pressure is used, neither result is significant ($p=0.26$ and 0.4 , respectively).

Thus, in our enalapril treated patients there was no fall in glomerular filtration rate after the first six months. Blood pressure was less important for the prognosis of renal function than type of anti-hypertensive treatment in these patients with well controlled blood pressure.

STAFFAN BJÖRCK

Department of Nephrology,
Sahlgrenska Hospital, University of Gothenburg,
S-413 45 Gothenburg,
Sweden

1 Granerus G, Aurell M. Reference values for Cr-EDTA clearance as a measure of glomerular filtration rate. *Scand J Clin Lab Invest* 1981;41:611-6.

2 Björck S, Nyberg G, Mulec H, Granerus G, Herlitz H, Aurell M. Beneficial effects of angiotensin converting enzyme inhibition on renal function in patients with diabetic nephropathy. *BMJ* 1986;293:467-70.

Rehabilitation of amputees

SIR,—S J D Chadwick and John H N Wolfe rightly point out that early referral of a patient with critical ischaemia of the lower limb is essential and that if salvage is not possible sufficient time will be available to prepare the patient for amputation.¹ They fail to emphasise, however, that rehabilitation must start immediately, not postoperatively.

Most amputations should be planned procedures that allow time for full medical and social assessment, improvement in general condition, adequate control of pain, counselling, and psychological help. Early assessment allows patients to collaborate in setting realistic, achievable goals and may prevent the disappointment, disillusion, and despair resulting from overoptimistic promises about future capabilities.

Patients who are referred late to prosthetic centres for assessment and measurement, only to be advised that wheelchair independence is a more realistic goal, inevitably plead for the chance to walk. In this forlorn struggle ritual visits for physiotherapy may ensue and amputees and professionals alike become increasingly frustrated. Repeated attendances at clinics focus on the discomfort of prolonged periods sitting in a wheelchair wearing the leg, its weight, and requests for a lighter leg as the grieving process continues—fuelled by failure to extinguish memories of youthful vigour and cope with the harsh reality of loss of the limb. This more common result of above knee or Gritti-Stokes amputation emphasises that preserving a functional below knee stump as a new organ of locomotion is probably the single most important physical factor in maximising functional ability.

Rehabilitation has major consequences for effective use of resources. The rehabilitation team can have a major impact on reducing hospital stay and repeated outpatient physiotherapy visits and improving use of prostheses.² The importance of specialist interdisciplinary, hospital based teams dedicated to rehabilitating amputees working in close cooperation with prosthetic centres cannot be emphasised too greatly. Some districts have already achieved this, but others lag far behind in affording amputees the special consideration they require. Some changes recommended by McColl have now been implemented.³ The Disablement Services Authority was finally integrated into the NHS in April last year, and most prosthetic centres are now led by consultants in rehabilitation. As Chadwick and Wolfe point out, new elderly amputees require extensive support and supervision if they are to return to the community. Therefore, rehabilitation consultants should coordinate the support and specialist services required to enable efficient and effective integration back into the community.

J A LINDSAY

Regional Rehabilitation Centre,
Birmingham B29 6JA

1 Chadwick SJD, Wolfe JHN. Rehabilitation of the amputee. *BMJ* 1992;304:373-6. (8 February.)

2 Ham R, Regan JM, Roberts VC. Evaluation of introducing the team approach to the care of the amputee: the Dulwich study. *Prosthet Orthot Int* 1987;11:25-33.

3 McColl I. *Review of artificial limb and appliance centre services*. London: HMSO, 1986.

SIR,—In their article on rehabilitation of amputees S J D Chadwick and John H N Wolfe make the point that an amputee who can walk is the main aim.¹ This is a good general rule but perhaps not one to be pursued at all costs.

For a considerable proportion of elderly patients who have had above knee (and especially bilateral) amputations, persevering in the attempt to obtain prosthetic ambulation is futile and even dangerous.^{2,3} The energy cost of ambulation with one above knee prosthesis is roughly twice that for someone with two normal legs. For someone whose cardiac or respiratory system is compromised that might be too much. The energy costs of wheelchair independence are less, and, provided that the patients do not have any problems with their arms, this might be the preferred option.

Much disappointment on the part of both the patient and the doctor can be averted if, for these selected few patients, wheelchair independence (as against prosthetic mobility) is opted for in the first instance, with support from the occupational therapist and social worker to adapt the home. Foam "trouser fillers" to restore the body image are required for some of these patients. Some above knee amputees use the prosthesis only as a prop for transferring themselves, and this may be a legitimate use of a prosthesis.

S MUTHU

Donald Tod Rehabilitation Centre,
Fazakerley Hospital,
Liverpool L9 7AL

1 Chadwick SJD, Wolfe JHN. Rehabilitation of the amputee. *BMJ* 1992;304:373-6. (8 February.)

2 Muthu S. Limb fitting and survival in the dysvascular double above-knee amputee. *J R Coll Surg Edinb* 1983;28:157-9.

3 Muthu S. Amputation of both legs in the elderly. *BMJ* 1983;287:1225.

Morbidity due to asthma

SIR,—K P Jones and colleagues have confirmed previous reports of the level of morbidity among patients with asthma.¹ I am concerned, however, about their analysis of the correlates of morbidity and the claims made for their morbidity index.

I assume that subjects were asked to summarise their asthma over the previous year (this is the

interval used by Hilton *et al*, from whose work the questionnaire was developed, and is the interval noted in table II).² I suspect that having to review morbidity over such a long period may result in considerable overreporting. I have observed some overreporting with my own questionnaire, which reviews only the previous six months.³ The usefulness of such lengthy reporting in subjects under 15 years old is even more doubtful as symptoms are more likely to come and go in children.⁴ Subjects aged under 15 comprised more than a third of the subjects in Jones and colleagues' study.

The diary of peak flow was completed in the week after the interview and reflects the state of asthma then. The questionnaire, on the other hand, described the asthma over the previous year. Though the peak flow recorded in the diary will probably be related to the reporting of morbidity because they were both reported by the same patients, it is not reasonable to claim that the index constructed from symptoms over the past year "relates to lung function" as lung function was not measured at the same time.

I believe that the authors have given an important signal to doctors treating asthma. We should not fail to assess the long term impact of asthma in our history taking. In testing the value of our strategies to improve care in asthma, however, we must continue to refine our measurement of morbidity in patients with asthma.

PATRICK T WHITE

Department of General Practice and Primary Care,
King's College School of Medicine and Dentistry,
London SE5 9PJ

1 Jones KP, Bain DJG, Middleton M, Mulec MA. Correlates of asthma morbidity in primary care. *BMJ* 1992;304:361-4.

2 Hilton S, Sibbald B, Anderson HR, Freeling P. Evaluating health education in asthma—developing the methodology: preliminary communication. *J R Soc Med* 1982;75:625-30.

3 White PT, Pharoah CA, Anderson HR, Freeling P. Randomised controlled trial of small group education on the outcome of chronic asthma in general practice. *J R Coll Gen Pract* 1989;39:182-6.

4 Anderson HR, Bland JM, Patel S, Peckham C. The natural history of asthma in childhood. *J Epidemiol Community Health* 1986;40:121-9.

Occupational causes of disorders in the upper limbs

SIR,—N J Barton and colleagues miss the point in their review of occupational causes of disorders in the upper limb.¹ They are guilty of the crime of diagnostic greed, for they want clear cut physical signs and histological evidence as well, presumably in individual cases. How else can any condition be prescribed as an industrial disease if it is to be unambiguously defined, both clinically and pathologically?

We suggest a simpler approach. History taking concerning the onset and relief of pain may provide clear clues in individual cases. Pain occurring later during working days and later on during the week suggests that the complaint may be related to work. Pains that improve over the weekend or remit during holidays or during intercurrent illness add weight to a clinical suspicion that symptoms are related to work. If the patient's job is analysed and ergonomic measures taken and if symptoms respond then a "therapeutic test" has supported the hypothesis that the condition is related to work. This is a reasonable approach in less acute cases. In cases of the so called repetitive strain syndrome, however, it may be too late to apply such a therapeutic test.

European Community legislation adds weight to the views of ergonomists. Simply because pain is an everyday part of working life does not mean that it should not be prevented or relieved, and legislation and the recognition of "prescribed disorders" are inevitable. The only approach that will work, particularly where diagnostic entities overlap so frequently and the measurement of disability is so