legitimate work of public health medicine and happen daily.

FRADA ESKIN

Yorkshire Health,
Harrogate HG1 5AH

1 Whitty P, Jones I. Public health heresy: a challenge to the purchasing orthodoxy. *BMJ* 1992;304:1039-41. (18 April.)

SIR,—Paula Whitty and Ian Jones's article questioning the role of public health doctors in the purchasing of health care services is challenging.¹ It lacks, however, a broader perspective on the problems faced by the specialty of public health medicine in procuring health gain.

Firstly, the authors mention just briefly the impact of fundholding by general practitioners on the organisation and planning of health care in a population. With post-election hindsight (and a new secretary of state for health, whose ambivalence towards trusts and whose dogged determination over fundholding is well publicised), this "wild card" in the NHS reforms deserves greater and urgent attention.² Public health doctors with a purchasing role in their district health authority have generally ignored the purchasing power of fundholders, especially those practices operating in consortiums. In so doing they may have tipped the delicate balance between the need to remain influential in the district health authority and the need to retain independent professional responsibility for the health of the whole population too far in the former direction. In addition, there are no incentives for fundholders to take regard of departments of public health medicine with a view to obtaining rational advice on purchasing.

Secondly, the authors assert that NHS services are poor at promoting health gain in comparison with fundamental changes in socioeconomic conditions. Historically most NHS resources have been focused on secondary health care at the expense of primary health care services. But primary health care services are rarely regarded as potential providers of cost effective, integrated, and locally responsive health care; rather they are considered merely to be efficient gatekeepers. Although, like secondary health care, primary health care remains essentially unevaluated, the reorientation of medical care towards preventing disease, promoting health, and primary health care is an inevitable but, hopefully, worthwhile strategic trend.³

Thirdly, the purchasing role of family health services authorities has at best been marginal (in terms of absolute money available) yet their potential for influencing health gain is unrealised. Some faltering steps are being made by a few public health doctors in health care planning and developing strategy in primary health care. The relationship between medical advisers or directors and public health medicine is of paramount importance, yet there is little evidence that it is being managed proactively. I suspect that the future success of public health medicine depends more on its central involvement in primary health care than it does on any other purchasing role.

COLIN POLLOCK

Carlton,
Goole,
North Humberside DN14 9QP

1 Whitty P, Jones I. Public health heresy: a challenge to the purchasing orthodoxy. *BMJ* 1992;304:1039-41. (18 April.)

2 Scheuer MA, Robinson R. A wild card in the pack? *Health Service Journal* 1991;101:18-9.

3 European Office of the World Health Organisation. *Regional strategy for attaining health for all by the year 2000*. Copenhagen: WHO/EURO, 1981. (EUR/RC 3018 rev 1.)

SIR,—Paula Whitty and Ian Jones's article on public health medicine is misleading and over-pessimistic.¹ The authors fail to acknowledge the growth throughout the 1980s of the new public health movement, which seeks to influence from

within the NHS as well as operating more widely through structures such as the healthy cities movement and external organisations such as the Public Health Alliance.^{2,4}

It is erroneous to attribute the current division of working time of those in public health to the NHS and Community Care Act. Since 1974 public health has been anchored in the NHS management structure, and much of the time and energy of public health physicians has been devoted to health care services. The authors produce no evidence that more time has been spent on health care services and less on health promotion since the implementation of the act.

Though there are serious problems with the NHS and Community Care Act, we see particular advantages to the split between purchasing and provision. Health authorities are now responsible for the health of a defined population, and, as partners in purchasing teams, public health physicians have a major role in the distribution of resources. We are now able to require information on outcomes and effectiveness as part of service contracts.

The fear that assessment of health care needs will not improve the public's health derives from the work of McKeown⁵ and assumes that the only important outcome measures are reductions in mortality. We suggest that a reduction in distress and disability due to the activities of health care services is equally important. Indeed, Whitty and Jones are in danger of belittling the daily contribution that the one million employees of the NHS make in ameliorating inequalities and relieving distress, disability, and disease. Not only do they adduce no evidence to support their assertion that public health skills are "presently redundant" in the measurement of outcomes but they later state that public health has an accepted role in evaluating the benefit of health services.

Like Whitty and Jones, we are concerned about the mergers of health authorities, impact of fundholding, effects of relatively diminishing resources, and serious lack of reliable information on activity and effectiveness of health services. Our particular skills in advocacy and analytical interpretation of data and the ability to take an overview of the population will help to meet the challenges within the NHS as well as in a wider local context.

Now is an exciting time to work in public health, as the quantity and quality of our new recruits attest. Our work is reflective and analytical, and we are more prone than other specialties to question our role. In that case we should take care to give the positive aspects as wide publicity as the negative ones.

ALLYSON POLLOCK   MARY BOOTHROYD-BROOKS
AILEEN CLARKE   JIA KANI
NISH CHATURVEDI   MARK MCCARTHY
IAN BASNETT   JANET MURRAY

Department of Public Health Medicine,
Bloomsbury and Islington Health Authority,
London NW1 2LJ

1 Whitty P, Jones I. Public health heresy: a challenge to the purchasing orthodoxy. *BMJ* 1992;304:1039-41. (18 April.)

2 Connelly J, Kelleher C, Morton S, St George D, Roderick P, eds. *Housing or homelessness: a public health perspective*. London: Faculty of Public Health Medicine, 1991.

3 Public Health Alliance. *Health on the move: policies for health promoting transport. The policy statement of the Transport and Health Study Group*. Birmingham: Public Health Alliance, 1991.

4 Ashton J, Seymour H. *The new public health*. Milton Keynes: Open University Press, 1988.

5 McKeown T. *The role of medicine: dream, mirage or nemesis?* Oxford: Blackwell, 1989.

Vertebral fractures

SIR,—Cyrus Cooper and L Joseph Melton's editorial brings together several of the current thoughts on spinal deformity.¹ We disagree, however, with the conclusion that the "clinically

manifest burden [of vertebral fractures] is considerable." The burden of a vertebral fracture must be the continued pain, disability, or deformity that arises plus the prospect of further fractures in the future.

Any quantification of fractures should closely relate structural deformation to symptoms. The authors barely touched on the problem of grading deformities let alone relating them to symptoms. The plethora of grading systems indicates the problem. No system rises appreciably above all others in its ability to relate to symptoms.

In a study that we carried out in Nottingham subjective assessment was no worse than any of three grading systems that were compared by cross correlations of deformity, and none of the grading systems related significantly to pain or disability (unpublished findings). The study also showed that even the measurements of vertebral dimensions that the grading systems use are open to subjective interpretation.

We are surprised that Cooper and Melton do not quote a study by Cooper himself with colleagues.² This cross sectional epidemiological study of vertebral fractures showed, like ours, that there was no significant difference in the prevalence of back pain between groups with and without fractures. A study by Ross *et al* did show a correlation of back pain with limitation of activity and vertebral fractures but only for those in whom lateral radiography showed a loss of 35% of the area of at least one vertebra.³ This was the most severe grade and is above the grades 1 and 2 that Cooper and Melton quote. Ross *et al* found no correlation of fractures of Cooper and Melton's grades 1 and 2 with back pain unless there were multiple deformations.

So do we need grading at all? Why not define vertebral fractures in population studies as occurring above some level of deformity—say, a 35% loss of area?

From Cooper and Melton's figures for the frequency of vertebral fractures in the United Kingdom, we would expect 114 patients a year to be admitted with painful vertebral fractures in Nottingham. Hospital Activity Analysis data, however, show that only 18 patients were admitted with pain. Though we acknowledge that coding problems and underreporting occur with Hospital Activity Analysis data, we suggest (based on admissions to hospital) that the clinical problem is probably smaller than the authors suggest.

The subtitle of the editorial asked how large the silent epidemic is. Do we have a silent epidemic or not? Until a relevant definition of a vertebral fracture is found further research work will be hampered.

ALISON L ARMSTRONG
GARETH THOMAS
W ANGUS WALLACE

Department of Orthopaedics,
University Hospital,
Nottingham NG7 2UH

DES GREEN

Department of Radiology,
City Hospital,
Nottingham NG5 1PB

1 Cooper C, Melton LJ. Vertebral fractures. *BMJ* 1992;304:793-4. (28 March.)

2 Cooper C, Shah S, Hand DJ, Adams J, Compston J, Davie M, *et al*. Screening for osteoporosis using individual risk factors. *Osteoporosis Int* 1991;2:48-53.

3 Ross PD, Ettinger B, Davis JW, Melton LJ III, Wasnich RD. Evaluation of adverse health outcomes associated with vertebral fractures. *Osteoporosis Int* 1991;1:134-40.

Routine weighing during antenatal visits

SIR,—The tone of the editorial on routine weighing during antenatal visits was disappointing.¹ The authors seem to overlook the fact that routine antenatal care is no more than a screening process. Thus the tests applied during antenatal care should