occurred more than seven weeks after his admission for self cutting. During five days of inpatient observation he was wary but coherent and rational. Repeated examination found no psychotic signs. With the ward team's agreement he was discharged into intensive community care, was seen repeatedly at his home and at the daily living programme team's base, managed well, and found a job. Two days after last being seen he suddenly killed a child and revealed previously concealed paranoid delusions. Had he remained an inpatient he could have absconded, avoided outpatient follow up, and become someone for whom district services could disclaim responsibility. Community care teams like the daily living programme team which do assertive outreach, however, continue their responsibility for patients and thus their vulnerability to blame for adverse events.

Conclusions from our study await our final report. They will have to balance the clinical and social outcomes, patients' and relatives' satisfaction, risk assessment, and an economic analysis.

MATT MUIJEN

ISAAC MARKS

JOSEPH CONNOLLY

Institute of Psychiatry and Maudsley Hospital,
London SE5 8RZ

Public opinion, the NHS, and the media

SIR,—The annual survey of public opinion on the NHS, attributed by Ken Judge and colleagues to the National Association of Health Authorities and Trusts,¹ is in fact commissioned jointly by the *Health Service Journal* and NAHAT. Its findings have been reported extensively in the *Health Service Journal* with an accompanying interpretation each year since we began polling in 1985.

The journal reached the same conclusion as your authors—that use of the NHS and medical coverage affect attitudes towards the service—as long ago as 1987. Across our seven polls, the number of people who have been satisfied with hospital treatment has fluctuated by only 4% and has averaged 86.5%. By contrast, those recording a good opinion of the NHS locally—who would include many who had not used the service—had fluctuated by 18% and averaged 72.5%.

The survey has down the years collected data on a wide range of attitudes to the NHS, including opinions on the level of funding; sources of extra revenue—including willingness to pay higher taxes—and priority areas for spending it; attitudes to curbing the availability of some treatments on the NHS; willingness to travel for treatment and to pay for "extras" during a hospital stay; the extent of understanding of the NHS reforms; aspects of hospital treatment most liked and disliked; difficulties finding a general practitioner; satisfaction with dental treatment; and the use of alternative medicine.

Reports of past years' results can be found in issues of the *Health Service Journal* for 6 June 1985, 15 May 1986, 2 April 1987, 19 May 1988, 22 June 1989, 2 August 1990, and 14 November 1991.

PETER DAVIES

Health Service Journal,
London WC2R 3LF

1 Judge K, Solomon M, Miller D, Philo G. Public opinion, the NHS, and the media: changing patterns and perspectives. *BMJ* 1992;304:892-5. (4 April.)

Duplicate publication

SIR,—The Universities Funding Council's current research assessment exercise is of such a nature that despite Lock's request eight years ago¹ and

Tony Waldron's recent evidence about duplicate publication² "Doctors are now under such pressure to publish—to impress appointments and grants committees—that the temptation to milk as many publications as possible out of a piece of research is strong."³ Moreover, regrettably, this may be unavoidable as the inadequate system that the Universities Funding Council uses in assessing research has necessarily converted the phrase "publish and be damned" into "publish or be damned."

There is, however, one way by which journals and, indeed, reviewers themselves can help. It requires a personal computer, modem, communications software, and telephone. When these are used to connect to Compuserve there is easy access to PaperChase 24 hours a day, seven days a week. PaperChase is an online information system that requires no special formats or terms and no experience or training to search successfully; saves searches automatically for up to six months; and searches both the continually updated MEDLINE database (over 6.5 million biomedical references to articles published in 4000 international journals) and the continually updated Health Planning and Administration (HEALTH) database (over 500 000 references on issues to do with the delivery of health care).

Two recent examples may be helpful. Firstly, when I checked on one paper PaperChase produced (within six minutes and ready for local printing) six references, five of which included the same author; the abstracts of three of these five contained the same statistical data. Secondly, and within seven minutes, five independent references were found to cover almost identical work but had not been included in a paper being considered for publication. The latest uncited reference had been published in a leading international journal in February this year.

As journals seek to maintain not only their integrity but also that of the scientific community, the use of this system by journals and reviewers may encourage potential authors to be more critical before seeking publication. This simple screening approach has the added advantage that, while Medical Subject Headings (MeSH) can be accessed quickly, papers can also be identified by using authors' names. Furthermore, my most recent search (on 13 April) confirmed that PaperChase was so comprehensive as to be ahead in time. It covered "the Entire MEDLINE Database from January 1966 Thru June 1992 [sic] Update, Part 1; and the Entire Health Planning and Administration Database, from January 1975 thru the April Update." Potential duplicating authors beware.

BRIAN LIVESLEY

Charing Cross and Westminster Medical School,
London W6 8RP

- 1 Lock S. Repetitive publication: a waste that must stop. *BMJ* 1984;288:661-2.
- 2 Waldron T. Is duplicate publishing on the increase? *BMJ* 1992;304:1029. (18 April.)
- 3 Lowry S, Smith J. Duplicate publication. *BMJ* 1992;304:999-1000. (18 April.)

Salami publication

SIR,—Tony Waldron's paper and the accompanying editorial regard the issue of salami publication from a biased stance.¹² When much time, effort, and money have been invested in carefully designed and executed large or long term prospective trials it is understandable and ethically defensible that all potentially important results should eventually be published. The American coronary artery surgery study of early surgery versus medical treatment is a well known case in point. Multicentre trials may have a publications committee to oversee the proper release of results. Few megatrials can afford to investigate only one hypothesis at a time.

More often, several different aspects need to be investigated in the same study.

Also, it may be impossible to accommodate all the interesting results in only one paper, especially as the trend in prestigious general journals seems to favour concise and pithy offerings. Indeed, in much the same way that salami is better served sliced than consumed whole, some readers may prefer to digest various key messages delivered separately in a series of short papers, whether in one journal or different ones. Editors should not forget that in writing and submitting several papers from one major study authors also spend more time and energy than they do in simply preparing one all embracing report.

Finally, when debate arises after the preliminary presentation of findings at scientific meetings or publication of the definitive paper there may be a case for further analysis of the data and further publication of substantial findings that support or generate important hypotheses requiring prospective study. In our view, clinical science has a better chance of advancement if interesting papers are allowed the opportunity to stand the test of peer review and time than if research is stifled by a blinkered view of salami publication.

M J BENNIE

C W LIM

Liverpool L7 3QH

- 1 Waldron T. Is duplicate publishing on the increase? *BMJ* 1992;304:1029. (18 April.)
- 2 Lowry S, Smith J. Duplicate publication. *BMJ* 1992;304:999. (18 April.)

Listen to the patient

SIR,—Maurice Wooldridge's note about a patient who changed his practice¹ reminds me of a lesson I learnt early in my career.

As a young researcher doing one clinic in rheumatology a week, I saw a woman at The London Hospital who complained that every time she walked she had a strange unsteadiness and tended to fall over, particularly when pegging out the washing. In those days everything at The London Hospital was viewed through a haze of syphilis thanks to the teaching of a distinguished series of directors of the "special department."

To my frustration the Wassermann reaction was negative (twice). I was unable to cure the woman despite diligent examination and sent her to the neurologists, who wrote back to me mentioning cervical spondylosis (Lord Brain was then head of the department). Dissatisfied with this explanation, I sent her to a colleague, the ever meticulous and sensible Duncan Vere, who wrote me the following illuminating letter.

Dear Tony,

Thank you for asking me to see this lady who has a strange sensation as if walking on sorbo-rubber.

On examination she had sorbo-rubber insoles, the removal of which produced an immediate improvement in her symptoms.

It was an incident that warned me (fortunately, very early on) of the dangers of taking the history one would like to hear rather than listening to what the patient is saying.

J A HICKLIN

Crawley Hospital,
Crawley,
West Sussex RH11 7DH

1 Wooldridge M. Beware the repeat prescription. *BMJ* 1992;304:824. (28 March.)

Advice to authors

Priority will be given to letters that are less than 400 words long and are typed with double spacing