notice. This would accommodate changes in personal or professional circumstances. Family health services authorities could match skills to jobs. Job sharing and mobility would be facilitated. The whole exercise may be construed as a global job share. "Ghettoisation" for either sex would become voluntary rather than imposed by consumers.

Family health services authorities' skill banks would offer a wider distribution of services and facilitate planning. Benefits would accrue to consumers and practitioners through a wider choice of service and variety of experience, respectively. Referral to secondary care would probably fall.

Modular contracting would allow specialisation within the primary health care team. Specialisation by general practitioners could be extended into "primary consultant" posts: obstetrics, minor surgery, dermatology, psychiatry, rheumatology, diabetes, audit, and geriatrics are obvious candidates. Hospital, Department of Social Security, and industrial appointments could also attract recognition.

Modular contracts would offer improved provision of services, easy planning, flexibility, rewards for effort and skill, accreditation, some independence, career planning, and equality of opportunity.

STEVEN FORD

Haydon Bridge,
Northumberland NE47 1HJ

Goya's living skeleton

SIR,—I am grateful to Richard and Maureen Park for providing contemporary descriptions of the feet and hands of Claude Seurat, the "living skeleton."^{1,2} They do not exclude a diagnosis of fibrodysplasia ossificans progressiva as valgus deformities of the great toes and clinodactyly are common features of the condition.³ Ossification within the paraspinal and cervical muscles, so well seen in the drawings of Seurat, is invariable but does not always cause scoliosis or torticollis. About 70% of patients have involvement of the jaw muscles, which would explain why Seurat could eat only small fragments of food.

As his bones cannot be examined the diagnosis of Seurat's illness must be speculative, but fibrodysplasia ossificans progressiva remains a strong possibility.

GEOFFREY HOOPER

Princess Margaret Rose Orthopaedic Hospital,
Edinburgh EH10 7ED

1 Park R, Park M. Goya's living skeleton. *BMJ* 1992;304:844. (28 March.)

2 Park RHR, Park MP. Goya's living skeleton. *BMJ* 1991;303:1594-6. (21-28 December.)

3 Connor JM, Evans DAP. Fibrodysplasia ossificans progressiva. *J Bone Joint Surg [Br]* 1982;64:76-83.

BUPA and consultants over 70

SIR,—British United Provident Association (BUPA) has announced that by the end of this year it proposes to make no further payments to insured patients in respect of consultants' fees if the consultant is aged 70 or over. I believe this to be a mean cost cutting exercise dressed up in a mantle of spurious righteousness. I speak as a doctor who has subscribed to BUPA since the day it was born.

At the moment none of the other private medical contributory schemes are wielding the axe to consultants aged over 70. I suggest that medical practitioners who are under 70 and are currently insured with BUPA should start shopping around and switch their personal insurance to whichever company offers the best deal.

I have two specific complaints, one general and

one personal in so far as it concerns my practice. I have patients who have been under my care for 30 years. I think it a bit much to have to say to them that, because of my age, if they belong to BUPA, from the end of this year they will not be refunded what they pay me. They don't wish to go to a younger man; if they did they would have gone long ago.

In the United Kingdom I am the only person who corrects severe myopia by inserting an implant into the anterior chamber of the eye without disturbing the crystalline lens, which has a high minus optic. I have done 40 such operations over the past two and a half years; I think I have a duty to keep doing them to benefit patients and so that I can accumulate experience to lay before my colleagues to help them take an informed decision on whether they should start doing this operation themselves. Consultants' potential contributions to the welfare of their patients and their specialty should be considered individually.

D P CHOYCE

London W1M 7DG

*We sent Mr Choyce's letter to BUPA, whose reply is given below.

SIR,—In declining to pay benefits to members for fees from specialists over the age of 70 BUPA is bringing itself into line with private hospitals and the NHS. Most private hospitals withdraw admitting privileges at the age of 70. Consultants over 70 are permitted to undertake locum posts in the NHS only with the specific permission of the Department of Health. Since the NHS requires specialists to retire at 65 it could be argued that BUPA should have followed suit. The motive for introducing this change has nothing to do with cost containment but is to promote the highest possible quality and most up to date medical advice to BUPA's members.

It would be invidious for BUPA to make individual assessments of the practice of consultants over 70. It would take only one untoward incident in which the advice given to a member could not be supported by current medical opinion for the whole system of specialist medical advice to be called into question.

Much of the clinical work that individual consultants over 70 have described in correspondence with BUPA seems to be long term follow up and surveillance of patients whose conditions have long since ceased to be "acute episodes of illness or injury." Treatment falling outside this definition is ineligible for reimbursement under the terms of the contract between BUPA and its subscribers.

I am unable to comment on the specific procedure that D P Choyce describes other than to note that it seems to be experimental; as such, neither Choyce's services nor the hospital costs are eligible for reimbursement under BUPA's rules.

HUGH THELWALL-JONES

Group medical director, BUPA,
London WC2R 3AX

Predictors of general practitioners' workload

SIR,—We agree with R Balarajan and colleagues that a scheme for compensating general practitioners for the effects of social deprivation should be based on an objective measure of workload such as consultation rates.¹ The use of the general household survey for this purpose, however, has serious limitations. The data on workload are based only on consultations over two weeks, so that even with a large sample there are wide confidence intervals around the odds ratios for the associations of socioeconomic variables with consultation. No distinction is made between consultations for illness and consultations for preventive pro-

cedures, which are likely to be used by the least deprived groups and for which general practitioners are usually remunerated separately. All consultations are counted equally, the workload generated by a telephone consultation being equated with that generated by a home visit, which people in households without access to a car are especially likely to require. For these reasons the analysis is likely to underestimate the effects of deprivation on workload.

We have also examined the effects of deprivation on workload by using consultations from the third national morbidity survey.² This dataset, linking general practice consultations with the census records of 140 050 subjects aged over 1 year, overcomes the limitations above. Our results, like those of Balarajan and colleagues, emphasise the importance of housing tenure and car ownership as predictors of workload. These two studies provide a basis for a more rational system for compensating general practitioners for the effects of deprivation than the Jarman underprivileged area index.

YOAV BEN-SHLOMO

IAN WHITE

Department of Epidemiology and Public Health,
University College and Middlesex School of Medicine,
London WC1E 6EA

PAUL MCKEIGUE

Department of Epidemiology and Population Sciences,
London School of Hygiene and Tropical Medicine,
London WC1E 7HT

1 Balarajan R, Yuen P, Machin D. Deprivation and general practitioner workload. *BMJ* 1992;304:529-34. (29 February.)

2 Ben-Shlomo Y, White I, McKeigue PM. Prediction of general practice workload from census-based social deprivation scores. *J Epidemiol Community Health* (in press).

Doctors and control of major releases of chemicals

SIR,—Though they are rare, major disasters leading to the release of poisonous gases have occurred. That health authorities should plan for disasters has been stated several times in the recent past.^{1,3} Chemical disasters, however, pose additional problems to those posed by other disasters in that poisonous gases can drift, causing morbidity and death over several kilometres.⁴ Prompt control of a release of toxic gas and the protection of the surrounding population demand the coordinated work of several authorities and emergency services. Recently we have participated, as medical officers, in a planning exercise to control a hypothetical major release of chemicals.

The incident medical officer has a dual role in such exercises. He or she will communicate with the local health authority and also be the source of medical advice to the incident control team. He or she will be the person best able to assess the possible health care needs of victims involved in the incident and then communicate this to the relevant health authority, hospitals, and general practitioners.

In addition, the medical officer will be the only site based source of medical skill for the incident control team. Whether or not to evacuate populations depends ultimately on the health gain or loss resulting from evacuation compared with no evacuation. This decision tends to be taken by the senior police officer or senior executive of the district local authority, neither of whom is likely to have the training or skill to make such judgments on his or her own. Deciding when to evacuate a population is not easy and requires a strong medical input.

That public health doctors should have a central role in planning for, and managing, disasters has been emphasised.^{4,5} Assessment of the health risks to the general population in releases of chemicals requires the skills of public health doctors. We suggest that in the United Kingdom the most appropriate public health doctor is the consultant

in communicable disease control. The control of major outbreaks has many similarities to the control of chemical incidents. There is often a similar sense of urgency and public alarm. Also, consultants in communicable disease control still often act as advisers on environmental health and toxicological issues to local authorities. These consultants should participate in planning for, and controlling, major releases of chemicals. Improving training in this subject for consultants in communicable disease control should be a high priority.

PAUL R HUNTER
PHILIP T MANNION

Public Health Laboratory,
Chester, Cheshire CH2 3EG

1 Bliss AR. Major disaster planning. *BMJ* 1984;288:1433-4.
2 Nancekivell DG. Disaster management: practice makes perfect. *BMJ* 1989;298:477.
3 Baxter PJ, Davies PC, Murray V. Medical planning for toxic releases into the community: the example of chlorine gas. *Br J Ind Med* 1989;46:277-85.
4 Korn DA, McIntyre L, Conway J. The emerging role of public health in community emergencies and disasters. *Can J Public Health* 1988;79:12-5.
5 Cormie K, Edwards J, Howell J, Jones D, Mills K, Ready H. The Edmonton tornado disaster: the role of the public health department. *Can J Public Health* 1988;79:6-10.

Economics and the fundamental principles of the NHS

SIR,—In the first of a series of articles addressing questions raised by *Leading for Health: a BMA Agenda for Health*.¹ J B L Howell expresses two fears. The first concerns the reliability of explicit methods of rationing health care resources such as the Oregon experiment and the ranking of treatments according to cost per quality adjusted life year (QALY). Howell argues that the management of waiting lists is “morally and practically the more acceptable.”²

Waiting lists provide a partial view of need. They say little about the urgency, cost, or effectiveness of the treatment required by those waiting or about the relative burdens borne by families, the community, and social services because of any delay in treatment. Nor does everyone in need get on to a waiting list. “The general conclusion to be drawn from published waiting list statistics is that it is very difficult to tell what is going on.”³ Furthermore, the recent scramble, at considerable cost, to remove from waiting lists by 31 March patients who had been waiting over two years was a questionable use of resources aimed at an essentially political target.

Howell’s second fear is that explicit methods of rationing legitimise denying care to some patients. Health economics is concerned not with whether to use resources but with how to use them.⁴ It makes explicit the health benefits that would be gained by increasing or changing patterns of expenditure. By advocating a system that does not attempt to quantify these opportunity costs explicitly Howell does not make the choices between different uses of health care resources any easier. Nor does he manage to assume away the need to make such decisions.

The Oregon experiment highlighted these stark choices. The final cut off below which treatments would not be funded was drawn after an expansion of funds allocated to health care.⁵ The exercise made explicit how much people’s tax liabilities would have to rise to fund most of the treatments defined as being critical to basic health care.⁶ It does not strike me that such an exercise rationalises denying patients care.

I agree with Howell that as yet we cannot have much faith in lists of treatments prioritised on the basis of QALYs. QALYs are in their infancy and are crude. Compiling estimates of costs per QALY, however, provides useful insights into the cost effectiveness of different forms of care.^{7,8}

I do not share Howells’s lack of faith in value judgments made by the public. In the Oregon experiment certain rankings of treatments were overturned because of their unacceptability. This highlights criteria on which society wants to base its decisions to allocate health care resources. If the ranking of treatment in a QALY league table and the cut off below which treatment will not be funded prove unacceptable, our values regarding the correct level and basis of resource allocation in the NHS are forced into the public arena. We are forced to justify them and are made aware of their cost in terms of the health benefits that we lose by pursuing them. These are the goals of health economics. To caricature it as a system for legitimising denying people care is simply to misunderstand it.

NEIL CRAIG

Newcastle upon Tyne NE12 9BH

1 BMA. *Leading for health: a BMA agenda for health*. London: BMA, 1991.
2 Howell JBL. Re-examining the fundamental principles of the NHS. *BMJ* 1992;304:297-9. (1 February.)
3 Radical Statistics Health Group. *NHS reforms: The First Six Months—proof of progress or a statistical smokescreen?* *BMJ* 1992;304:705-9. (14 March.)
4 Henry D. Economic analysis as an aid to subsidisation decisions. The development of Australian guidelines for pharmaceuticals. *Pharmacoeconomics* 1992;1:54-67.
5 Maynard A. Priorities: follow the Oregon trail? *Medical Audit News* 1992;2:20-2.
6 Steinbrook R, Lo B. The Oregon Medicaid demonstration project—will it provide adequate care? *N Engl J Med* 1992;326:340-4.
7 Bryan S, Parkin D, Donaldson C. Chiropody and the QALY: a case study in assigning categories of disability and distress to patients. *Health Policy* 1991;18:169-85.
8 Donaldson C, Atkinson A, Bond J, Wright K. QALYs and long-term care for elderly people in the UK: scales for assessment of quality of life. *Age Ageing* 1988;17:379-87.

Consultants’ hours

SIR,—We have now sorted out juniors’ hours moderately well. Is it not time that some consideration was given to sorting out the hours and conditions of service of consultants, especially in high profile specialties with low numbers of consultants such as accident and emergency, and often paediatrics, where absence on leave can result in consultants being expected to provide rather more cover than is permissible for juniors? Certainly in accident and emergency the absence of one consultant in a department with two consultants can result in the other being expected to be available on a one in one basis for a considerable time if no locum is provided, as is the case in so many places nowadays.

It is surely time that the BMA addressed this problem, which lasts for 30 years in many people’s careers, as well as juniors’ problems, which last for only a short time during their total career.

C S FLOWERS

Cheltenham General Hospital,
Cheltenham,
Gloucestershire GL53 7AN

Private insurers not passing the buck

SIR,—Jonathan Boyce’s editorial on private practice, subtitled “Troubled insurers prepared to pass the buck,” is misleading.¹

The reasons for the steeply rising costs of medical insurance claims are well known (more treatments, more sophisticated methods of treatment, the increasing complexity of private sector work, an increasingly elderly population, etc), but private medical insurers are not just sitting back and complacently passing the buck.

Private Patients Plan has been working, in partnership with hospitals and doctors, for several years to find ways of controlling costs of claims

without jeopardising the quality of care. This work, which has already had some success, is continually breaking new ground, to everyone’s benefit.

One example is Private Patients Plan’s pilot study over the past few years of whether the concept of managed care can be successfully introduced in the United Kingdom. Working closely with a large employer, doctors, and a hospital in the north of England, Private Patients Plan has been encouraged by the findings so far and has embarked on the next phase of the study. Another example is Private Patients Plan’s psychiatric benefit protection scheme, which has been developed with the enthusiastic support and participation of virtually all providers of private psychiatric care in Britain. Again, early results are encouraging and the scheme promises to enable Private Patients Plan to maintain cost controlled cover in this important aspect of care.

Additionally, to streamline administration procedures and thereby speed up payment of claims Private Patients Plan has worked with British United Provident Association to develop a common schedule of procedures with standard codings for use by doctors, hospitals, and insurers. We hope that this will soon become a standard for the industry, which will again be of widespread benefit.

Thus, far from complacently passing the buck, Private Patients Plan strongly believes in working in partnership with others to solve common problems and will continue to battle against steeply rising costs.

CHRISTOPHER JOHNSON

Private Patients Plan,
London WC1H 9LJ

1 Boyce J. Private practice. *BMJ* 1992;304:458-9. (22 February.)

Weightings used in distributing NHS resources

SIR,—E S Williams and colleagues observe that the standardised mortality ratio is negatively correlated with the percentage of people aged 65 and over living in English district health authorities and conclude that any distribution of NHS funds based on a positive weighting for the proportion of elderly people is therefore unfair.¹ Two issues should be emphasised.

Firstly, the correlation that they showed is significant. The prime determinant of the significance of any correlation or simple linear regression slope, however, is the number of points contributing to it. The authors did not give the scattergram of the 190 contributing district health authorities or the coefficient of determination (a measure of the strength of the association as opposed to its significance). The figure may explain why. The correlation may be significant but is not strong. The coefficient of determination, r^2 , is low at 0.11. Indeed, a second linear regression excluding the

Simple linear regression of population of district health authorities aged over 65 and standard mortality ratio. $y = -1.2x + 120.5$, $r^2 = 0.11$