the patients were lost to follow up. This would be unacceptable in any trial in acute myocardial infarction or heart failure, in which I have considerable experience; I cannot understand why a study in hypertension would be different. This loss of patients indicates that the investigators have had serious problems monitoring their patients. Furthermore, it is known that patients lost to follow up differ from patients who remain in trials.

Secondly, the number of patient years during which patients received treatment as a proportion of the total number of patient years in the subgroups was 69% in the diuretic group, 55% in the β blocker group, and 69% in the placebo group. Obviously, if close to 50% of the assigned treatment is not administered it will not be possible to observe any effect.

Thirdly, the number of withdrawals from β blocker treatment in this single blind study is in contrast to the experience in the Swedish trial in old patients with hypertension (STOP trial): 177 patients treated with β blockers in the Medical Research Council's trial were withdrawn because of bradycardia compared with none in the STOP trial (B Dahlöf, personal communication).

Fourthly, the power of the trial was calculated to study active treatment versus placebo. There is clearly insufficient power to permit conclusions regarding lack of effects between the treatment subgroups and placebo group—even more so considering my second point above. Wisely, the STOP trial did not present its subgroup results.

None of these important flaws is discussed by the authors. I suggest that the trial should be discussed more in the context of pitfalls and flaws than in the context of treatment of hypertension.

KARL SWEDBERG

Department of Medicine,
Östra Hospital,
S-416 85 Göteborg,
Sweden

- 1 MRC Working Party. Medical Research Council trial of treatment of hypertension in older adults: principal results. *BMJ* 1992;304:405-12. (15 February.)
- 2 Dahlöf B, Lindholm LH, Hansson L, Schersten B, Ekbom T, Wester P-O. Morbidity and mortality in the Swedish trial in old patients with hypertension (STOP-hypertension). *Lancet* 1991;338:1281-5.

EDITOR,—All physicians are asked to advise on the management of blood pressure and often feel lonely when competing with drug company promotion. This feeling is intensified when a report carrying the imprimatur of the Medical Research Council draws a misleading conclusion.¹ In this single blind study the only unexpected finding was the death rate—an end point not subject to the same influences as the others. Of the 4396 patients in the study, 823 died—409 in the treatment group and 414 in the placebo group, a difference of five deaths. Expressed in another way, there were five additional deaths in 25355 patient years of observation. About half of the patients were lost from the study, so the beneficial effect needs to be halved—one additional death in 10000 patient years.

Regardless of statistical games, can such marginal benefit warrant the time and trouble for these elderly patients and their enthusiastic doctors, whose additional time is at least financially rewarded? Similar sentiments expressed by others in reaction to the report relieve my feeling of isolation.² Drugs and matching placebo were kindly donated by drug companies. These “free” drugs are even more effective than free lunches in promoting sales.

Another paper on the management of elderly patients with sustained hypertension analyses the MRC's report and five others.³ None of the six studies were designed to detect a change in overall mortality, and five of the six failed to do so. Despite this failure of treatment to improve survival, treatment is advised because of the reported

improvement in cardiovascular morbidity and mortality.

But what treatment? For young independent people I have no hesitation in recommending weight reduction, exercise, and stopping smoking. In geriatric patients strictures on food and tobacco may be enforced by the carer but encouraging exercise is more difficult. I advise the avoidance of excess salt and alcohol, which is quite different from advising a moderate reduction; is there any convincing evidence that this helps in managing blood pressure?

And so to drugs. If “there is little to choose between individual thiazides” then the cheapest should surely be mentioned at its optimum low dose (bendrofluazide 2.5 mg daily).⁴ The safety of these inexpensive agents is emphasised in subsequent correspondence.⁵ The older the patient the more the reduction in renal function and the more the dangers of hyperkalaemia with triamterene or amiloride. β Adrenoceptor blocking drugs, both hydrophilic and lipophilic, have been used in trials that claimed benefit,⁶ but they are not much favoured because of unwanted effects.

“None of the newer agents have been subject to definitive controlled trials . . . nevertheless such drugs may be favoured on a number of theoretical and practical grounds.” The support of Bristol Myers Squibb is acknowledged. Which of the newer agents will be given to these elderly people, now patients, before proof of efficacy?

A B S MITCHELL

North London Postgraduate Medical Centre,
St Ann's Hospital,
London N15 3TH

- 1 Medical Research Council Working Party. MRC trial of treatment of hypertension in older adults: principal results. *BMJ* 1992;304:405-12. (15 February.)
- 2 Hand CH. Treatment of hypertension in older adults. *BMJ* 1992;304:639. (7 March.)
- 3 Beard K, Bulpitt C, Mascie-Taylor H, O'Malley K, Sever P, Webb S. Management of elderly patients with sustained hypertension. *BMJ* 1992;304:412-6. (15 February.)
- 4 Carlsen JE, Kober L, Torp-Pedersen C, Johansen P. Relation between dose of bendrofluazide, antihypertensive effect, and adverse biochemical effects. *BMJ* 1990;300:975-8.
- 5 Johnston GD. Treatment of hypertension in older adults. *BMJ* 1992;304:639. (7 March.)
- 6 Kendall MJ. Treatment of hypertension in older adults. *BMJ* 1992;304:639. (7 March.)

EDITOR,—The results of the Medical Research Council's trial of treatment of hypertension in older adults raise at least two important questions that are not discussed.¹

Firstly, the trial reported that about a quarter of the patients were lost to follow up. About half of the patients stopped taking their randomised treatment and many patients received diuretic or β blocker treatment in addition to their original randomised treatment. The findings from this trial cannot be generalised to the elderly hypertensive population at large.

Secondly, total mortality and high mortality from coronary heart disease in hypertensive patients can be substantially reduced only if the risk of sudden coronary death can be reduced. β Blockers have prevented coronary death in secondary prevention trials, mainly by reducing the risk of sudden death.^{2,3} Possibly the therapeutic effects in this respect differ between β blockers. Benefits of β blockade on sudden death in patients after myocardial infarction have been shown only for the lipophilic β blockers. Data from several primary prevention studies indicate a lower risk for hypertensive men taking lipophilic β blockers than for those taking diuretics.⁴ In the metoprolol atherosclerosis prevention in hypertensives trial the risk of coronary events was 24% lower in hypertensive men receiving β blockade than in men receiving diuretics.⁴ Also, total mortality was reduced with the β blocker regimen due to a reduction in sudden cardiovascular deaths.⁵

Clinical and experimental data indicate that not

only ischaemia and increased sympathetic tone but also a low vagal tone (influenced by brain mechanism) play an important part in the development of atherosclerosis and in reduced cardiac electric stability.^{2,3,6-10} Experimental data with propranolol and metoprolol suggest that β blockade within the brain can decrease the risk of ventricular fibrillation and sudden death by increasing cardiac vagal tone.^{9,10} In one study spontaneous ventricular fibrillation was reduced with metoprolol but not with the hydrophilic β blocker atenolol.¹⁰ Possibly, relatively lipophilic β blockers, which easily pass the blood-brain barrier, more readily increase vagal tone and reduce the risk of ventricular fibrillation than hydrophilic β blockers, which pass the blood-brain barrier to only a small degree. Thus data obtained with one particular antihypertensive drug may not be generalised to other similar drugs as the effects (other than their blood pressure lowering effect) may differ even within one class of drugs. The results from this study should not be used to judge the relative importance of different antihypertensive drugs in preventing coronary heart disease.

JAAKKO TUOMILEHTO

National Public Health Institute,
SF-00510 Helsinki,
Finland

- 1 MRC Working Party. Medical Research Council trial of treatment of hypertension in older adults: principal results. *BMJ* 1992;304:405-12. (15 February.)
- 2 Hjalmarson Å, Olsson G. Myocardial infarction. Effects of beta-blockade. *Circulation* 1991;84(suppl VI):101-7.
- 3 Wikstrand J, Berglund G, Tuomilehto J. Beta-blockade in the primary prevention of coronary heart disease in hypertensive patients. *Circulation* 1991;84(suppl VI):93-100.
- 4 Wikstrand J, Warnold J, Tuomilehto J, Olsson G, Barber H, Eliasson K, et al. Metoprolol versus thiazide diuretics in hypertension. Morbidity results from the MAPHY study. *Hypertension* 1991;17:579-88.
- 5 Olsson G, Tuomilehto J, Berglund G, Elmfeldt D, Warnold I, Barber H, et al. Primary prevention of sudden cardiovascular death in hypertensive patients: mortality results from the MAPHY study. *Am J Hypertens* 1991;4:151-8.
- 6 Eckberg DL. Beta-adrenergic blockade may prolong life in post-infarction patients in part by increasing vagal cardiac inhibition. *Med Hypotheses* 1984;15:421-32.
- 7 Hayano J, Yamada A, Mukai F, Sakakibara Y, Yamada M, Ohte N, et al. Severity of coronary atherosclerosis correlates with the respiratory component of heart rate variability. *Am Heart J* 1991;121:1070-9.
- 8 Predictors of sudden death after myocardial infarction [editorial]. *Lancet* 1991;338:727-8.
- 9 Parker GW, Michael LH, Hartley CJ, Skinner JE, Entman JE. Central beta-adrenergic mechanisms may modulate ischemic ventricular fibrillation in pigs. *Circ Res* 1990;66:259-70.
- 10 Åblad B, Björö T, Björkman J-A, Edström T, Olsson G. Role of central nervous beta-adrenoceptors in the prevention of ventricular fibrillation through augmentation of cardiac vagal tone (abstract). *J Am Coll Cardiol* 1991;17:165A.

AUTHORS' REPLY,—We think that Karl Swedberg has not fully appreciated the practical demands and clinical implications of long term trials as distinct from the shorter term examples he cites.

As was pointed out in our paper, we used the NHS central register for notification of deaths; such notifications were provided whether or not we were still in touch with participants. Ascertainment of fatal events was therefore virtually complete. The duration of follow up averaged 5.8 years and, for some patients, was eight years. Clearly, some loss to follow up over these extended periods is inevitable: the proportion was 25% in each of the diuretic, β blocker, and placebo groups (considerably less than A B S Mitchell suggests), so biased ascertainment of non-fatal events is unlikely to have occurred.

Our trial was based on 226 general practices throughout the United Kingdom, and, to deal with Jaakko Tuomilehto's first point, its results indicate the extent to which supplementation of treatment and changes in treatment are likely to be needed in day to day clinical practice. Our “on treatment” analyses, in which departures from randomised treatment were allowed for, gave results similar to those of the intention to treat analyses. We have not seen the data on bradycardia in the Swedish trial in old patients with hypertension (STOP trial)

referred to in the personal communication to Swedberg, but in our trial withdrawals for this reason were according to prespecified criteria and in line with usual clinical practice.

The size of our trial was indeed based on a comparison of active with placebo treatment, but, as we pointed out, an assessment of the two active agents was an important secondary objective from the outset. The advantage of the diuretic over the β blocker was clear, particularly for coronary events, and is also suggested by other recent results.¹ Swedberg is surely not suggesting that findings of such immediate clinical relevance should remain unreported.

STANLEY PEART

Hunterian Institute,
Royal College of Surgeons,
London WC2A 3PN

T W MEADE

MRC Epidemiology and Medical Care Unit,
Wolfson Institute of Preventive Medicine,
London EC1M 6BQ

S J POCOCK

Department of Epidemiology and Medical Sciences,
Medical Statistics Unit,
London School of Hygiene and Tropical Medicine,
London WC1E 7HT

1 Beard K, Bulpitt C, Mascie-Taylor H, O'Malley K, Sever P, Webb S. Management of elderly patients with sustained hypertension. *BMJ* 1992;304:412-6.

Life events and breast cancer prognosis

EDITOR,—As Jennifer Barraclough and colleagues¹ point out, the findings of their prospective study of life events and breast cancer prognosis differ from those of a cross-control study undertaken at Guy's Hospital.² Possible reasons for these apparently discrepant results should be sought before any influence of severe life experiences on relapse of breast cancer is dismissed.

At Guy's Hospital cases and controls were carefully matched for the major factors that influence outcome, including histological grade and number of involved nodes. At Southampton this information was not available. Thus important imbalances could have been present between patients with and without severe life experiences.

Clearly, knowledge of relapse status introduces the possibility of bias, but this is possible in any life events study as events can be ascertained only retrospectively. In the Southampton study the patients were not blind to their disease status at the time of an interview, and it must have been difficult for the interviewer to remain so. Bias in the rating of severity of events can be minimised if this is done by a panel of judges who are unaware of disease status, as was the case at Guy's Hospital.

At Guy's Hospital all interviews were conducted with the patients themselves, whereas for 11 of the 47 patients who relapsed at Southampton a final interview was conducted only with next of kin and for two other patients who relapsed no interview was performed. This could have led to under-reporting of severe life experiences in a most important group.

The apparently high incidence of severe "own health" experiences not related to breast cancer observed at Southampton, affecting 21 patients during the 42 month follow up, was surprising. No such severe health experiences were recorded at Guy's Hospital. We were therefore interested in the finding that severe own health experiences had a significant adverse effect on outcome ($p=0.01$). The authors comment that it makes intuitive sense that patients with breast cancer in poor general health are more vulnerable to relapse. This is unproved.

The power of the Southampton study was diminished by the fact that 42 patients declined to participate and that these patients had a higher

death rate (and presumably higher relapse rate) than those interviewed. We estimate that with 204 patients entered and a 23% relapse rate the study had about a 50% chance of detecting a difference between the groups if the true relative risk attributable to serious life experiences is 2.0.

Differences in the use of tamoxifen between the two studies may be important. At Southampton most postmenopausal women with positive nodes received tamoxifen. None of the patients at Guy's Hospital received tamoxifen (all patients presented initially before 1987). An intriguing possibility arises that the beneficial effect of tamoxifen could be partly mediated by counteracting any adverse effect of severe events. This benefit might be most pronounced in hormone sensitive tumours. In the Guy's Hospital study the influence of severe events on relapse was most apparent among the patients with tumours positive for oestrogen receptors.³

Recurrence free survival in women with breast cancer who had a severe event in first two years of follow up compared with other women in study (patients followed up to January 1991)

In our view the question of whether serious life experiences influence relapse is still unresolved. We are conducting a prospective study similar to that reported by Barraclough and colleagues, in which we are also assessing the impact of severe events on immunological and endocrinological variables. We would also encourage Barraclough and colleagues to re-examine the outcome for their patients after a longer follow up. This would not entail collection of further data on life events as we found that severe events occurring within the first two years after diagnosis were largely responsible for the positive outcome of our study (figure).

A J RAMIREZ J P WATSON
M A RICHARDS I S FENTIMAN
W M GREGORY R D RUBENS
T J K CRAIG

Imperial Cancer Research Fund Clinical Oncology Unit
and Division of Psychiatry,
Guy's Hospital,
London SE1 9RT

- 1 Barraclough J, Pinder P, Cruddas M, Osmond C, Taylor I, Perry M. Life events and breast cancer prognosis. *BMJ* 1992;304:1078-81. (25 April.)
- 2 Ramirez AJ, Craig TJK, Watson JP, Fentiman IS, North WRS, Rubens RD. Stress and relapse of breast cancer. *BMJ* 1989;298:291-3.
- 3 Ramirez AJ, Richards MA, Gregory WM, Craig TJK. Psychological correlates of hormone receptor status in breast cancer. *Lancet* 1990;335:1408.

EDITOR,—Jennifer Barraclough and colleagues report the lack of an association between psychosocial stress and relapse in breast cancer.¹ The results of their prospective study contrast with those of two studies showing a positive association between adverse life events and the onset and relapse of breast cancer.^{2,3} The results of all three studies are based on data obtained with the same interview (the life events and difficulties schedule), suggesting that Barraclough and colleagues' opposite conclusions arise from differences in the design and analysis of their study.

Given the slow evolution of many breast cancers, the period of assessment before relapse may need

to be several years. Geyer found an excess of adverse life events in the eight years before diagnosis in patients with breast cancer compared with controls.² The study of Ramirez *et al* covered life events over a median of 30.5 months before relapse.³ A comparable figure is not given in Barraclough and colleagues' paper, but as the maximum period of assessment was 42 months the average period covered by the life events and difficulties schedule was probably shorter.

Barraclough and colleagues' patients were older (54.3 years) than those in the studies of Ramirez *et al* (49.5 years) and Geyer (45.5 years). These age differences may be clinically important as they are likely to be associated with differences in menstrual status, pathology, and the types of adverse life events experienced. Sixty per cent of Barraclough and colleagues' patients were postmenopausal compared with 31% in the study of Ramirez *et al*. As it may be hypothesised that psychosocial stress influences the onset or progress of cancer through indirect hormonal effects, these differences between study samples may be important in explaining contradictory results.

It would also be relevant to know what proportion of patients in both studies received tamoxifen, and their hormone receptor status. Allied to this, certain combinations of medical and surgical interventions may conceivably conceal the limited effects of psychosocial adversity on the disease process. Differences in treatment may also account for the lack of an association between life events and prognosis in certain studies.

These studies make the questionable assumption that all patients react similarly to adverse events. Yet, clearly, some people become more upset than others after the death of a spouse, loss of employment, or other major events. The link between adverse events and physical health may become clearer if an attempt is made to understand the way in which psychosocial stress interacts with personality and coping style to influence physiological processes.

The possibility of an association between psychological factors and the course of physical disease is a fascinating but clinically unimportant issue in the assessment of patients with cancer. Although future studies may confirm that adverse life events do not shorten the life span of such patients, psychological interventions can result in a significant improvement in quality of life for many of them.⁴

CHWEN CHENG CHEN
THOMAS FAHY

Department of Psychological Medicine,
King's College Hospital and Institute of Psychiatry,
London SE5 9RS

- 1 Barraclough J, Pinder P, Cruddas M, Osmond C, Taylor I, Perry M. Life events and breast cancer prognosis. *BMJ* 1992;304:1078-81. (25 April.)
- 2 Geyer S. Life events prior to manifestation of breast cancer: a limited prospective study covering eight years before diagnosis. *J Psychosom Res* 1991;35:355-63.
- 3 Ramirez AJ, Craig TJK, Watson JP, Fentiman IS, North WRS, Rubens RD. Stress and relapse of breast cancer. *BMJ* 1989;298:291-3.
- 4 Lovestone S, Fahy TA. Psychological factors in breast cancer. *BMJ* 1991;302:1219-20.

Psychological influences on cancer and ischaemic heart disease

EDITOR,—Anthony J Pelosi and Louis Appleby summarise recent research by Eysenck and Grossarth-Maticek about psychosocial influences on cancer and ischaemic heart disease.¹ They raise several critical questions; among them they suggest that "one is left to speculate whether the authors have made the mistake, during reanalysis of their data, of reassigning individuals to personality types after causes of death were known."

In an issue of *Psychological Inquiry*² a target