and as much a part of that nurse's initial assessment as deciding what observations to carry out or whether to lay the patient down.

I am also concerned about the wide discrepancies between the triage assessment of priority and that of the consultants. Either the consultants benefited from hindsight or there had been depressingly little prior communication in the department. Clear guidelines on assessing urgency were drawn up in my department by consultation between medical and nursing staff. New staff are properly trained in their use, and the accuracy is regularly monitored, consistently reaching over 90%.

The use of median waiting times is questionable; centile times convey a clearer picture. A median waiting time of 26 minutes for patients in priority category 1, however, is alarming. We define this group as patients who cannot safely wait more than 15 minutes for medical attention, and 97% of them receive attention within this time.

Triage should not be expected to cut waiting times. It does not after all reduce the amount of work to be done, it simply redistributes waiting times away from the more seriously ill towards those who will not suffer unduly from delays. A reduction in waiting times requires some patients to be treated or discharged by the assessing nurse—that is, it requires a nurse practitioner to be introduced. This is an entirely separate concept, and one of the references quoted by George and colleagues refers to a nurse practitioner system and not to triage.²

Prioritisation is clearly vital in the effective running of our seriously overloaded accident and emergency departments, and I am concerned that the often hesitant progress towards improving such prioritisation will be threatened by this well intentioned but, in my view, seriously flawed study.

Southend Hospital,
Westcliff on Sea,
Essex SS0 0RY

1 George S, Read S, Westlake L, Williams B, Fraser-Moodie A, Priddy P. Evaluation of nurse triage in a British accident and emergency department. *BMJ* 1992;304:876-8. (4 April.)
2 Maller J, Woolwich C. Triage in accident and emergency departments. *J Adv Nurs* 1990;15:1443-51.

SIR,—Steve George and colleagues have raised several important issues about using triage nurses in accident and emergency departments, particularly the possibility that triage might increase waiting times for all patients.¹ It is difficult, however, to determine from their paper the precise cause of the additional delays experienced by patients when triage was in use.

Only 48% of patients were actually assessed by a triage nurse during the triage periods, yet the median waiting times for all patients in the triage group were longer than those for the patients in the no triage group. Were the delays caused by the time taken for triage itself or by subsequent queuing problems as patients proceeded through the accident and emergency department? Perhaps one way of clarifying this would be to analyse further the extended waiting times during the periods when triage was being done. The waiting times of the 48% who were assessed by a triage nurse could be compared with the waiting times of the 52% of attenders during the same periods who were not. This might indicate whether the triage itself or subsequent queuing caused the delays.

The authors also analysed waiting times according to a retrospective judgment of urgency, made by a consultant in accident and emergency, rather than the prospective assessment of urgency made by the triage nurse. Given that the prospective assessment is intended to be a major determinant of waiting times (indeed, it is the whole purpose of triage), why did the authors not conduct such a prospective analysis?

The authors conclude that some form of priori-

tisation will benefit patients in most urgent need of care, but they question whether this needs to be formalised and called nurse triage. What system do they therefore recommend? Are they not simply calling for a more efficient form of nurse triage rather than none at all?

Royal Free Hospital,
London NW3 2QG

1 George S, Read S, Westlake L, Williams B, Fraser-Moodie A, Priddy P. Evaluation of nurse triage in a British accident and emergency department. *BMJ* 1992;304:876-8. (4 April.)

AUTHORS' REPLY,—Jeremy Dale and John Bache note that only 48% of patients attending during a triage period saw a triage nurse and conclude that outcome measurements for our triage group cannot be used to deduce the effects of triage itself and that our study is invalid. They misunderstand the principle of intention to treat trials. We made the decisions to redesignate shifts at the beginning of each day, not at a moment's notice, and we ensured that triage ran during each session designated. The periods in which nurse triage ran and did not run were entirely comparable and were not dictated by workload. In real life triage nurses may leave their stations to undertake clinical duties elsewhere, but this applied equally to both arms of our study. Ours is a valid study of the effectiveness of nurse triage in real life, whereas Dale and Bache propose a study of the efficacy of nurse triage under ideal circumstances.

John Heyworth and M Pledge suggest that urgency should be assessed retrospectively by nurses rather than doctors. This is irrelevant. So long as both groups are judged by the same assessors the assessors may be nurses, doctors, or, indeed, others. All have equal validity.

J E Porter questions the nature of the difference between formal triage and informal prioritisation in our study. Formal triage is a documented assessment governed by strict protocols, whereas the informal assessment is undocumented, intuitive, and dynamic and fits well with Porter's description of what happens in her department.

Triage ran for five years preceding our study. All triage nurses attend six lectures, are tested on theory, and undertake supervised practice before starting triage. Their collective skills and experience are considerable. Dale and Porter are correct, therefore, when they state that informal assessment was performed by nurses familiar with triage. But the question "Does prioritisation of patients by degree of urgency produce benefits compared with their being seen in strict order of arrival?" is artificial. It is nonsense to suggest that any accident and emergency department in the United Kingdom functions entirely in this fashion. At its simplest level, the urgent attention awarded a patient arriving by emergency ambulance is prioritisation, and the relevant question is "Does formal nurse triage produce benefits when compared with an existing scheme of informal prioritisation?" This was our question.

David St George requests an analysis of waiting times by prospective urgency. This would be ethically unacceptable as it would entail every patient being assessed for triage and then the

decision being ignored in half the cases. To answer his question regarding the cause of delays, however, we reanalysed waiting times, deducting the time taken for patients to see the triage nurse before booking in at reception. The results are essentially unchanged, showing that queuing problems after triage cause delays, rather than the triage process itself (table). Our and other studies show that nurses consistently give patients a higher priority than do clinicians.^{1,5} We must consider whether this benefits patients and, if not, how best it can be modified.

Our conclusion is to make triage more efficient rather than abandon it. Prioritisation by degree of urgency obviously benefits patients whose condition is most urgent, and this principle should be instilled in all who work in accident and emergency departments. It is the method by which that prioritisation is to be accomplished that is for debate.

STEVE GEORGE LINDA WESTLAKE
SUSAN READ BRIAN WILLIAMS

Medical Care Research Unit,
Department of Public Health Medicine,
University of Sheffield Medical School,
Sheffield S10 2RX

ALISTAIR FRASER-MOODIE PAUL PRITTY
Derbyshire Royal Infirmary,
Derby DE1 2QY

1 Russo RM, Gururaj VJ, Bunye AS, Kim YH, Ner S. Triage abilities of nurse practitioner vs pediatrician. *Am J Dis Child* 1975;129:673-5.
2 Zwicke DL, Bobzein WF, Wagner EG. Triage nurse decisions: a prospective study. *Journal of Emergency Nursing* 1982;8:132-8.
3 Albin S, Wassertheil-Smoller S, Jacobsen S, Bell B. Evaluation of emergency room triage performed by nurses. *Am J Public Health* 1975;65:1063-8.
4 Mills J, Webster AL, Wofsy CB, Harding P, D'Acuti D. Effectiveness of nurse triage in ED of an urban county hospital. *Journal of the American College of Emergency Physicians* 1976;5:877-82.
5 Rivara FP, Wall HP, Worley P, James KD. Pediatric nurse triage: its efficacy, safety and implications for care. *Am J Dis Child* 1986;140:205-20.

In search of the unknown primary

SIR,—Christopher Bradley and Peter Selby succinctly identify the dilemmas that challenge all clinicians who must decide how best to manage patients presenting with metastases from an unknown primary.¹ An exhaustive search for the primary is clearly clinically unjustifiable once malignancies that can be therapeutically influenced have been excluded. While such possibilities are being investigated, however, the time required for imaging and to obtain specialised histopathological results and measurements of serum markers should not be allowed to delay adequate palliation in symptomatic patients.

Many patients labelled as having an unknown primary present with pain from bone metastases. Palliative radiotherapy given in one day or over one week will relieve pain promptly in over 80% of patients without either delaying any essential investigation or compromising the ability to give adequate cytotoxic chemotherapy subsequently.

Patients with disseminated malignancy from an unknown primary are stressed by uncertainty

Median waiting time by study group and retrospective priority category. Figures given for triage group are exclusive (and inclusive) of time taken to see triage nurse

Retrospective priority category	Median waiting time (minutes)			
	Triage group (n=2515)	No triage group (n=2522)	Difference (triage minus no triage)	95% Confidence interval for difference*
1 (Most urgent)	26 (26.5)	16	10	1.5 to 13
2	44 (46)	37.5	6.5	1 to 11
3	56 (58)	55	1	0 to 2
4 (Least urgent)	62.5 (66)	62	0.5	-4 to 4

*Calculated using method described by J P Nicholl, meeting of International Society of Clinical Biostatistics, Maastricht, September 1989.